

Title: Quasi-central elements and p -nilpotence of finite groups

Author(s): Huaquan Wei, Yanming Wang and Guohua Qian

Let G be a finite group and let P be a Sylow p -subgroup of G . An element x of G is called quasi-central in G if $\langle x \rangle \langle y \rangle = \langle y \rangle \langle x \rangle$ for each $y \in G$. In this paper, it is proved that G is p -nilpotent if and only if $N_G(P)$ is p -nilpotent and, for all $x \in G \setminus N_G(P)$, one of the following conditions holds: (a) every element of $P \cap P^x \cap G^{\mathcal{N}_p}$ of order p or 4 is quasi-central in P ; (b) every element of $P \cap P^x \cap G^{\mathcal{N}_p}$ of order p is quasi-central in P and, if $p = 2$, $P \cap P^x \cap G^{\mathcal{N}_p}$ is quaternion-free; (c) every element of $P \cap P^x \cap G^{\mathcal{N}_p}$ of order p is quasi-central in P and, if $p = 2$, $[\Omega_2(P \cap P^x \cap G^{\mathcal{N}_p}), P] \leq Z(P \cap G^{\mathcal{N}_p})$; (d) every element of $P \cap G^{\mathcal{N}_p}$ of order p is quasi-central in P and, if $p = 2$, $[\Omega_2(P \cap P^x \cap G^{\mathcal{N}_p}), P] \leq \Omega_1(P \cap G^{\mathcal{N}_p})$; (e) $|\Omega_1(P \cap P^x \cap G^{\mathcal{N}_p})| \leq p^{p-1}$ and, if $p = 2$, $P \cap P^x \cap G^{\mathcal{N}_p}$ is quaternion-free; (f) $|\Omega(P \cap P^x \cap G^{\mathcal{N}_p})| \leq p^{p-1}$. That will extend and improve some known related results.

Address:

Huaquan Wei
School of Mathematics Science
Guangxi Teachers Education University
Nanning, 530001
China

Address:

Yanming Wang
Lingnan College
Zhongshan University
Guangzhou, 510275
China

Address:

Guohua Qian
Department of Mathematics
Changshu Institute of Technology
Changshu, 215500
China