

On a characterization theorem on Abelian groups

By GENNADIY FELDMAN (Kharkov) and MARGARYTA MYRONYUK (Kharkov)

Abstract. Let $\xi_1, \xi_2, \dots, \xi_n$, $n \geq 2$ be independent identically distributed random variables. It is well known that if $\bar{\xi} = \frac{1}{n} \sum_{j=1}^n \xi_j$ and $\mathbf{v} = (\xi_1 - \bar{\xi}, \xi_2 - \bar{\xi}, \dots, \xi_n - \bar{\xi})$ are independent, then all ξ_j are Gaussian. We give a complete description of second countable locally compact Abelian groups for which a group analogue of this characterization theorem holds true.

1. Introduction

It is well known that if $\xi_1, \xi_2, \dots, \xi_n$, $n \geq 2$ are independent identically distributed Gaussian random variables, then $\bar{\xi} = \frac{1}{n} \sum_{j=1}^n \xi_j$ and $\mathbf{v} = (\xi_1 - \bar{\xi}, \xi_2 - \bar{\xi}, \dots, \xi_n - \bar{\xi})$ are independent. Assume now that $\xi_1, \xi_2, \dots, \xi_n$, $n \geq 2$ are independent identically distributed random variables such that $\bar{\xi}$ and $\mathbf{v}$ are independent. Then $\bar{\xi}$ and $s^2 = \frac{1}{n} \sum_{j=1}^n (\xi_j - \bar{\xi})^2$ are also independent. This implies by Geary's theorem ([6], [11], [10], [13]) that the random variables ξ_j are Gaussian. Thus, a Gaussian measure on the real line is characterized by the independence of $\bar{\xi}$ and $\mathbf{v}$.

The article deals with a generalization of this characterization theorem to the case when independent random variables take values in a locally compact Abelian group. Since an arbitrary Abelian group generally is not a group with unique division by n , instead of $\bar{\xi}$ and $\mathbf{v}$ we consider $S = \sum_{j=1}^n \xi_j$ and $\mathbf{V} = (n\xi_1 - S, \dots, n\xi_n - S)$.

We will use in the article the standard results on structure of locally compact Abelian groups and the duality theory ([7]). Agree on notation. For an arbitrary

Mathematics Subject Classification: 60B15, 62E10, 39B99.

Key words and phrases: Gaussian measure, locally compact Abelian group.

locally compact Abelian group X denote by $Y = X^*$ its character group, and by (x, y) the value of a character $y \in Y$ at an element $x \in X$. If H is a closed subgroup of the group Y , then denote by $A(X, H) = \{x \in X : (x, y) = 1 \text{ for all } y \in H\}$ its annihilator. Denote by c_X the connected component of zero of the group X , and by b_X the set of all compact elements of X . Let n be a natural number. Put $X_{(n)} = \{x \in X : nx = 0\}$ and $X^{(n)} = \{x \in X : x = n\tilde{x}, \tilde{x} \in X\}$. Denote by $\mathbb{Z}$ the group of integers, by $\mathbb{R}$ the group of real numbers and by $\mathbb{T}$ the circle group (the one-dimensional torus). Let Y be an arbitrary Abelian group, $f(y)$ be a function on Y , h be an element of Y . Denote by Δ_h the finite difference operator

$$\Delta_h f(y) = f(y + h) - f(y).$$

A function $f(y)$ on Y is called a polynomial if for a nonnegative integer m $f(y)$ satisfies the equation

$$\Delta_h^{m+1} f(y) = 0, \quad y, h \in Y.$$

The minimal m for which this equality holds is called the degree of the polynomial $f(y)$.

We will assume in the article that X is a second countable locally compact Abelian group. Denote by $M^1(X)$ the convolution semigroup of probability distributions on X . For $\mu \in M^1(X)$ denote by

$$\hat{\mu}(y) = \int_X (x, y) d\mu(x)$$

its characteristic function. Note that if ξ is a random variable with values in X and distribution μ , then the characteristic function of the distribution μ is the mathematical expectation

$$\hat{\mu}(y) = \mathbf{E}[(\xi, y)].$$

Denote by E_x the degenerate distribution concentrated at the point $x \in X$. The set of all degenerate distributions on the group X denote by $D(X)$. For $\mu \in M^1(X)$ define the distribution $\bar{\mu} \in M^1(X)$ by the formula $\bar{\mu}(B) = \mu(-B)$ for any Borel set B . Denote by $\sigma(\mu)$ the support of a distribution μ .

A probability measure γ on the group X is called Gaussian (in the sense of PARTHASARATHY) ([12, Ch. 4.6]), if its characteristic function can be represented in the form

$$\hat{\gamma}(y) = (x, y) \exp\{-\varphi(y)\}, \quad (1)$$

where $x \in X$, and $\varphi(y)$ is a continuous nonnegative function on Y satisfying the equation

$$\varphi(u + v) + \varphi(u - v) = 2[\varphi(u) + \varphi(v)], \quad u, v \in Y. \quad (2)$$

Taking into account that in the article we will deal only with Gaussian measures in the sense of Parthasarathy we will name them Gaussian. Denote by $\Gamma(X)$ the set of Gaussian measures on the group X . Denote by m_K the normalized Haar measure of a compact subgroup K of the group X , and by $I(X)$ the set of shifts of such measures.

We will say that a distribution $\mu \in \Gamma_n(X)$ if there exist independent identically distributed random variables ξ_j , $j = 1, 2, \dots, n$, $n \geq 2$ with values in the group X and a distribution μ such that S and $\mathbf{V}$ are independent.

It is not difficult to verify that for a group X the inclusion $\Gamma(X) \subset \Gamma_n(X)$ holds. In §2 we completely describe groups X which have the following property: if $\mu \in \Gamma_n(X)$ and the characteristic function $\hat{\mu}(y)$ does not vanish, then $\mu \in \Gamma(X)$. We apply the results of §2 in §3 to give the complete description of locally compact Abelian groups X for which any distribution $\mu \in \Gamma_n(X)$ is invariant with respect to a compact subgroup K of the group X and under the natural homomorphism $X \mapsto X/K$ induces on the factor-group X/K a Gaussian measure. We can consider the obtained class of groups as the widest subclass of locally compact Abelian groups on which an analogue of the theorem about characterization of a Gaussian measure on the real line by the independence of $\bar{\xi} = \frac{1}{n} \sum_{j=1}^n \xi_j$ and $\mathbf{v} = (\xi_1 - \bar{\xi}, \xi_2 - \bar{\xi}, \dots, \xi_n - \bar{\xi})$ holds.

The above mentioned problems are reduced to solving of a functional equation in the class of continuous positive definite functions on the group $Y = X^*$.

2. The characteristic function $\hat{\mu}(y)$ does not vanish

Lemma 1. *A distribution $\mu \in \Gamma_n(X)$ if and only if its characteristic function $\hat{\mu}(y)$ satisfies the equation*

$$\begin{aligned} \prod_{j=1}^n \hat{\mu}(u + nv_j - (v_1 + \dots + v_n)) \\ = \hat{\mu}^n(u) \prod_{j=1}^n \hat{\mu}(nv_j - (v_1 + \dots + v_n)), \quad u, v_1, \dots, v_n \in Y. \end{aligned} \quad (3)$$

PROOF. Note that S and $\mathbf{V}$ are independent if and only if the equality

$$\mathbf{E}[(S, u)(\mathbf{V}, (v_1, \dots, v_n))] = \mathbf{E}[(S, u)]\mathbf{E}[(\mathbf{V}, (v_1, \dots, v_n))] \quad (4)$$

holds for all $u, v_1, \dots, v_n \in Y$. Taking into account that the random variables

$\xi_1, \dots, \xi_n$ are independent, we transform the left-hand side of (4):

$$\begin{aligned} \mathbf{E}[(S, u)(\mathbf{V}, (v_1, \dots, v_n))] &= \mathbf{E}[(\xi_1 + \dots + \xi_n, u)((n\xi_1 - (\xi_1 + \dots + \xi_n), \\ &\quad \dots, n\xi_n - (\xi_1 + \dots + \xi_n), (v_1, \dots, v_n))] \\ &= \mathbf{E} \left[\prod_{j=1}^n (\xi_j, u + nv_j - (v_1 + \dots + v_n)) \right] = \prod_{i=1}^n \widehat{\mu}(u + nv_i - (v_1 + \dots + v_n)). \end{aligned}$$

Analogously we transform the right-hand side of (4):

$$\begin{aligned} \mathbf{E}[(S, u)]\mathbf{E}[(\mathbf{V}, (v_1, \dots, v_n))] &= \mathbf{E}[(\xi_1 + \dots + \xi_n, u)]\mathbf{E}[(n\xi_1 - (\xi_1 + \dots + \xi_n), \\ &\quad \dots, n\xi_n - (\xi_1 + \dots + \xi_n), (v_1, \dots, v_n))] \\ &= \prod_{j=1}^n \mathbf{E}[(\xi_j, u)]\mathbf{E} \left[\prod_{j=1}^n (\xi_j, nv_j - (v_1 + \dots + v_n)) \right] \\ &= \prod_{j=1}^n \mathbf{E}[(\xi_j, u)] \prod_{j=1}^n \mathbf{E}[(\xi_j, nv_j - (v_1 + \dots + v_n))] \\ &= \widehat{\mu}^n(u) \prod_{i=1}^n \widehat{\mu}(nv_i - (v_1 + \dots + v_n)). \quad \square \end{aligned}$$

Suppose that $\gamma \in \Gamma(X)$ and the characteristic function $\widehat{\gamma}(y)$ has representation (1). By the function $\varphi(y)$ we can construct a symmetric 2-additive function by the formula

$$\psi(u, v) = \frac{1}{2}[\varphi(u + v) - \varphi(u) - \varphi(v)].$$

Then $\varphi(y) = \psi(y, y)$. Using this representation for the function $\varphi(y)$ one can check directly that the characteristic function $\widehat{\gamma}(y)$ satisfies equation (3). Hence, by Lemma 1 the inclusion

$$\Gamma(X) \subset \Gamma_n(X) \quad (5)$$

holds. The main result of this section is the complete description of groups X which have the property: if $\mu \in \Gamma_n(X)$ and the characteristic function $\widehat{\mu}(y)$ does not vanish, then $\mu \in \Gamma(X)$. The following proposition is valid.

Proposition 1. *Assume that $\mu \in \Gamma_2(X)$ and the characteristic function $\widehat{\mu}(y)$ does not vanish. This implies that $\mu \in \Gamma(X)$ if and only if the group X contains no subgroup topologically isomorphic to the two-dimensional torus $\mathbb{T}^2$. Assume that $\mu \in \Gamma_n(X)$, where $n \geq 3$. This implies that $\mu \in \Gamma(X)$ if and only if the group X contains no subgroup topologically isomorphic to the circle group $\mathbb{T}$.*

We need some lemmas to prove Proposition 1.

Lemma 2. *Let $X = \mathbb{T}$ and $n \geq 3$. Then there exists a distribution $\mu \in \Gamma_n(X)$ such that the characteristic function $\hat{\mu}(y)$ does not vanish and $\mu \notin \Gamma(X)$.*

PROOF. Let $n = 3$. Consider on the group $\mathbb{Z}$ the function

$$l(k) = \begin{cases} 1 & \text{if } k \in \mathbb{Z}^{(3)} \\ \exp\left\{\frac{2\pi i}{9}\right\} & \text{if } k \in 1 + \mathbb{Z}^{(3)} \\ \exp\left\{-\frac{2\pi i}{9}\right\} & \text{if } k \in 2 + \mathbb{Z}^{(3)}. \end{cases} \quad (6)$$

Obviously, (6) implies that

$$l^3(k) = \exp\left\{\frac{2\pi ki}{3}\right\}, \quad k \in \mathbb{Z}. \quad (7)$$

Taking into account (7) and the fact that $l(k+3p) = l(k)$, $k, p \in \mathbb{Z}$, we can verify directly that the function $l(k)$ satisfies equation (3) for $n = 3$.

Take $\sigma > 0$ in such a way that the inequality

$$\sum_{k \in \mathbb{Z}, k \neq 0} \exp\{-\sigma k^2\} < 1 \quad (8)$$

holds. Put

$$\rho(t) = 1 + \sum_{k \in \mathbb{Z}, k \neq 0} l(k) \exp\{-\sigma k^2 - ikt\}, \quad t \in \mathbb{R}.$$

Since $l(-k) = \overline{l(k)}$, $|l(k)| = 1$, $k \in \mathbb{Z}$, in view of (8) the inequality

$$\rho(t) > 0, \quad t \in \mathbb{R}$$

is valid. We also have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \rho(t) dt = 1.$$

Let μ be the distribution on the group $\mathbb{T}$ with density $r(e^{it}) = \rho(t)$ with respect to $m_{\mathbb{T}}$. By the construction the characteristic function of the distribution μ is of the form

$$\hat{\mu}(k) = l(k) \exp\{-\sigma k^2\}, \quad k \in \mathbb{Z}.$$

Since the function $l(k)$ satisfies equation (3) for $n = 3$, the characteristic function $\hat{\mu}(k)$ also satisfies equation (3) for $n = 3$. By Lemma 1 $\mu \in \Gamma_3(X)$. On the other hand, since the function $l(k)$ is not a character of the group $\mathbb{Z}$, we have $\mu \notin \Gamma(X)$.

Let $n > 3$. If $n = 2p - 1$ we put

$$l(k) = \begin{cases} 1 & \text{if } k \in q + \mathbb{Z}^{(n)}, \quad q = 0, 1, \dots, p-2, p+1, \dots, n-1, \\ \exp\left\{\frac{2\pi i}{n}\right\} & \text{if } k \in p-1 + \mathbb{Z}^{(n)} \\ \exp\left\{-\frac{2\pi i}{n}\right\} & \text{if } k \in p + \mathbb{Z}^{(n)}. \end{cases}$$

If $n = 2p$ we put

$$l(k) = \begin{cases} 1 & \text{if } k \notin p + \mathbb{Z}^{(n)} \\ -1 & \text{if } k \in p + \mathbb{Z}^{(n)}. \end{cases}$$

Next we argue as in the case when $n = 3$. □

Remark 1. Let $n \geq 3$. It is easily seen that the function $l(k)$ constructed in the proof of Lemma 2 is the characteristic function of a signed measure on the group $\mathbb{T}$ concentrated in the subgroup $\mathbb{Z}(n)$ (the multiplicative group of n th roots of unity).

Lemma 3. Assume that $\gamma \in \Gamma_n(X)$, and the characteristic function $\hat{\gamma}(y) > 0$ for $y \in Y$. Then $\gamma \in \Gamma(X)$, and the function $\hat{\gamma}(y)$ can be represented in the form (1), where $x = 0$.

PROOF. Put $\psi(y) = -\log \hat{\gamma}(y)$. By Lemma 1 the characteristic function $\hat{\gamma}(y)$ satisfies equation (3). Taking the logarithm of both sides of (3), we get

$$\begin{aligned} \sum_{j=1}^n \psi(u + nv_j - (v_1 + \dots + v_n)) \\ = n\psi(y) + \sum_{j=1}^n \psi(nv_j - (v_1 + \dots + v_n)), \quad u, v_j \in Y. \end{aligned} \quad (9)$$

We use the finite difference method to solve equation (9). Let h_1 be an arbitrary element of the group Y . Substitute $u + h_1$ for u and $v_j + h_1$ for v_j , $j = 1, 2, \dots, n$ in equation (9). Subtracting equation (9) from the resulting equation we obtain

$$\sum_{j=1}^n \Delta_{h_1} \psi(u + nv_j - (v_1 + \dots + v_n)) = n\Delta_{h_1} \psi(u), \quad u, h_1, v_j \in Y. \quad (10)$$

Let h_2 be an arbitrary element of the group Y . Substitute $u + h_2$ for u and $v_1 + h_2$ for v_1 in equation (9). Subtracting equation (10) from the resulting equation we get

$$\Delta_{nh_2} \Delta_{h_1} \psi(u + nv_1 - (v_1 + \dots + v_n)) = n\Delta_{h_2} \Delta_{h_1} \psi(u), \quad u, h_1, h_2, v_j \in Y. \quad (11)$$

Let h_3 be an arbitrary element of the group Y . Substitute $u + h_3$ for u and $v_2 + h_3$ for v_2 in equation (9). Subtracting equation (11) from the resulting equation we find

$$n\Delta_{h_3}\Delta_{h_2}\Delta_{h_1}\psi(u) = 0, \quad u, h_1, h_2, h_3 \in Y. \quad (12)$$

Put in (12) $h_1 = h_2 = h_3 = h$. We get

$$\Delta_h^3\psi(u) = 0, \quad u, h \in Y, \quad (13)$$

i.e. $\psi(y)$ is a continuous polynomial of degree ≤ 2 . It is easy to see that each polynomial of degree ≤ 2 , in particular $\psi(y)$, can be represented in the form

$$\psi(y) = \varphi(y) + l(y) + c, \quad y \in Y, \quad (14)$$

where the function $\varphi(y)$ satisfies equation (2), the function $l(y)$ satisfies equation

$$l(u + v) = l(u) + l(v), \quad u, v \in Y,$$

and $c = \text{const}$. Since $\hat{\gamma}(0) = 1$, we can assume that $c = 0$. Since the function $\hat{\gamma}(y)$ is real-valued, we have $\hat{\gamma}(-y) = \overline{\hat{\gamma}(y)} = \hat{\gamma}(y)$. Hence, $\psi(-y) = \psi(y)$. This implies that in (14) $l(y) = 0$, $y \in Y$. So, $\psi(y) = \varphi(y)$, $y \in Y$. This proves the lemma. $\square$

A distribution $\mu \in M^1(X)$ is called a Gaussian measure in the sense of Bernstein ([9, 5.3]) if μ has the following property: if ξ_1 and ξ_2 are independent identically distributed random variables with values in X and distribution μ , then their sum and difference are independent. We denote by $\Gamma_B(X)$ the set of Gaussian measures in the sense of Bernstein on the group X .

Lemma 4 ([9, 5.3]). *A distribution $\mu \in M^1(X)$ belongs to the class $\Gamma_B(X)$ if and only if the characteristic function $\hat{\mu}(y)$ satisfies the equation*

$$\hat{\mu}(u + v)\hat{\mu}(u - v) = \hat{\mu}^2(u)|\hat{\mu}(v)|^2, \quad u, v \in Y. \quad (15)$$

Lemma 5. $\Gamma_2(X) = \Gamma_B(X)$.

PROOF. Let $\mu \in \Gamma_2(X)$. By Lemma 1 the characteristic function $\hat{\mu}(y)$ satisfies equation (3) which takes the form

$$\hat{\mu}(u + (v_1 - v_2))\hat{\mu}(u - (v_1 - v_2)) = \hat{\mu}^2(u)|\hat{\mu}(v_1 - v_2)|^2, \quad u, v_1, v_2 \in Y. \quad (16)$$

Substituting $v_1 = v$, $v_2 = 0$ into (16), we obtain that the characteristic function $\hat{\mu}(y)$ satisfies equation (15). Hence, by Lemma 4 $\mu \in \Gamma_B(X)$. Lemmas 1 and 4 also imply that if $\mu \in \Gamma_B(X)$, then $\mu \in \Gamma_2(X)$. $\square$

PROOF OF PROPOSITION 1. Let $n = 2$. Applying Lemma 5 we reduce the proof of Proposition 1 to the proof of the corresponding statement for distributions from the class $\Gamma_B(X)$, but for such distributions this statement was proved in [2] (see also [5], Lemmas 9.6 and 9.7).

Let $n \geq 3$ and $\mu \in \Gamma_n(X)$. Put $\nu = \mu * \bar{\mu}$. It follows from Lemma 1 that $\nu \in \Gamma_n(X)$. Since $\widehat{\nu}(y) = |\widehat{\mu}(y)|^2 > 0$, by Lemma 3 $\nu \in \Gamma(X)$. If a group X contains no subgroup topologically isomorphic to $\mathbb{T}$, then by Cramer's theorem for locally compact Abelian groups ([1], see also [5, Theorem 4.6]), we get that $\mu \in \Gamma(X)$. Thus, the sufficiency when $n \geq 3$ is also proved. The necessity follows from Lemma 2. $\square$

3. The main theorem

Put $I_n(X) = I(X) \cap \Gamma_n(X)$. It follows from (5) and Lemma 1 that the inclusion

$$\Gamma(X) * I_n(X) \subset \Gamma_n(X)$$

holds. The main problem solved in this section is the following: to describe all groups X for which

$$\Gamma(X) * I_n(X) = \Gamma_n(X). \quad (17)$$

Observe that if $\mu \in \Gamma(X) * I(X)$, then μ is invariant with respect to a compact subgroup K of the group X and under the natural homomorphism $X \mapsto X/K$ induces on the factor-group X/K a Gaussian measure. We formulate now the main theorem.

Theorem 1. *Equality (17) holds for a group X , when $n = 2$, if and only if the connected component of zero c_X of X contains no more than one element of order 2. Equality (17) holds for a group X , when $n \geq 3$, if and only if c_X has the property*

$$\{x \in c_X : nx = 0\} = \{0\}. \quad (18)$$

To prove Theorem 1 we need some lemmas. First we will formulate as a lemma the following simple statement, but omit its proof.

Lemma 6. *Let n be a natural number, K be a compact subgroup of a group X . Then the following statements are equivalent:*

- (i) $K^{(n)} = K$;
- (ii) if $ny \in A(Y, K)$, then $y \in A(Y, K)$.

We note that the characteristic function of the Haar measure m_K is of the form

$$\widehat{m}_K(y) = \begin{cases} 1 & \text{if } y \in A(Y, K) \\ 0 & \text{if } y \notin A(Y, K). \end{cases} \quad (19)$$

Lemma 7. *Let K be a compact subgroup of a group X . Then the following statements are equivalent:*

- (i) $m_K \in I_n(X)$;
- (ii) $K^{(n)} = K$.

PROOF. (i) $\Rightarrow$ (ii). By Lemma 1 the characteristic function $\widehat{m}_K(y)$ satisfies equation (3). Substituting $v_1 = u$, $v_2 = \cdots = v_n = 0$ into (3), we get

$$\widehat{m}_K(nu) = \widehat{m}_K(u)\widehat{m}_K((n-1)u), \quad u \in Y. \quad (20)$$

It follows from (19) and (20) that if $ny \in A(Y, K)$, then $y \in A(Y, K)$. Hence, by Lemma 6 $K^{(n)} = K$.

(ii) $\Rightarrow$ (i). By Lemma 1 it suffices to check that the characteristic function $\widehat{m}_K(y)$ satisfies equation (3), which takes the form

$$\begin{aligned} & \prod_{j=1}^n \widehat{m}_K(u + nv_j - (v_1 + \cdots + v_n)) \\ &= \widehat{m}_K(u) \prod_{j=1}^n \widehat{m}_K(nv_j - (v_1 + \cdots + v_n)), \quad u, v_1, \dots, v_n \in Y. \end{aligned} \quad (21)$$

Both sides of equation (21) take the values either 0 or 1. If the right-hand side of (21) is equal to 1, then $u \in A(Y, K)$, $nv_j - (v_1 + \cdots + v_n) \in A(Y, K)$, $j = 1, 2, \dots, n$. Hence, $u + nv_j - (v_1 + \cdots + v_n) \in A(Y, K)$, $j = 1, 2, \dots, n$. Therefore the left-hand side of (21) is also equal to 1.

Let the left-hand side of (21) be equal to 1. This implies that

$$u + nv_j - (v_1 + \cdots + v_n) \in A(Y, K), \quad j = 1, 2, \dots, n. \quad (22)$$

It follows from (22) that

$$\sum_{j=1}^n (u + nv_j - (v_1 + \cdots + v_n)) = nu \in A(Y, K). \quad (23)$$

By Lemma 6 we get from (23) that $u \in A(Y, K)$. Then (22) implies that $nv_j - (v_1 + \cdots + v_n) \in A(Y, K)$, $j = 1, 2, \dots, n$. Hence, the right-hand side of (21) is also equal to 1. $\square$

Lemma 8 (compare with [4, Lemma 10.4]). *Let X be a discrete torsion free group. Then $\Gamma_n(X) = D(X)$.*

PROOF. If X is a discrete torsion free group, then Y is a connected compact group ([7, (24.25)]). Assume first that $Y \not\cong \mathbb{T}$. Then there exists a continuous monomorphism $p : \mathbb{R} \mapsto Y$ such that its image $p(\mathbb{R})$ is everywhere dense in Y ([7, (25.18)]). Let $\mu \in \Gamma_n(X)$. Consider the restriction of the characteristic function $\widehat{\mu}(y)$ to $p(\mathbb{R})$. Put $f(t) = \widehat{\mu}(p(t))$, $t \in \mathbb{R}$. By Lemma 1 the characteristic function $\widehat{\mu}(y)$ satisfies equation (3). Therefore $f(t)$ is a characteristic function on $\mathbb{R}$ satisfying (3) too. Taking into account that $\Gamma_n(\mathbb{R}) = \Gamma(\mathbb{R})$, we have

$$f(t) = \exp\{-\sigma t^2 + i\beta t\}, \quad \sigma \geq 0, \beta \in \mathbb{R}.$$

Since p is a monomorphism and $\overline{p(\mathbb{R})} = Y$, for any neighborhood of zero V of the group Y we can choose a sequence of real numbers $t_n \rightarrow \infty$ such that $p(t_n) \in V$ for all n . If $\sigma > 0$, then

$$|f(t_n)| = |\widehat{\mu}(p(t_n))| = \exp\{-\sigma t_n^2\} \rightarrow 0$$

when $t_n \rightarrow \infty$. Taking into account that V is an arbitrary neighborhood of zero of the group Y we come to contradiction with the continuity at zero of the function $\widehat{\mu}(y)$. Hence, $\sigma = 0$, and this implies that $|\widehat{\mu}(p(t))| = 1$ for $t \in \mathbb{R}$. Since $\overline{p(\mathbb{R})} = Y$, we have $|\widehat{\mu}(y)| = 1$ for $y \in Y$. It follows from this that $\mu \in D(X)$.

If $Y \cong \mathbb{T}$, then $X \cong \mathbb{Z}$. Since $\mathbb{Z}$ is a subgroup of $\mathbb{R}$, we have $\Gamma_n(\mathbb{Z}) \subset \Gamma_n(\mathbb{R}) = \Gamma(\mathbb{R})$. Taking into account that any Gaussian measure on $\mathbb{R}$ supported in $\mathbb{Z}$ is degenerated we completely prove the lemma. $\square$

Corollary 1. *Let Y be a connected compact group, $f(y)$ be a characteristic function on Y satisfying equation (3). Then $|f(y)| = 1$, $y \in Y$.*

Lemma 9. *Let $\mu \in \Gamma_n(X)$. Then μ is supported in a coset of a subgroup $G \cong \mathbb{R}^m \times K$, where $m \geq 0$, and K is a compact subgroup of X such that $K^{(n)} = K$.*

PROOF. Note first that if $\lambda \in \Gamma_n(X)$, then for any $x \in X$ we have $\lambda * E_x \in \Gamma_n(X)$. Therefore if it is necessary we can substitute a distribution $\lambda \in \Gamma_n(X)$ by its shift, and the shift also belongs to the class $\Gamma_n(X)$.

Let $\lambda \in M^1(X)$ be an arbitrary distribution. Consider the set

$$E = \{y \in Y : |\widehat{\lambda}(y)| = 1\}.$$

Then E is a subgroup of Y , and there exists an element $x \in X$ such that $\widehat{\lambda}(y) = (x, y)$ for $y \in E$. It follows from what has been said that we can substitute the distribution μ by its shift and assume from the beginning that

$$E = \{y \in Y : |\widehat{\mu}(y)| = 1\} = \{y \in Y : \widehat{\mu}(y) = 1\}. \quad (24)$$

It follows from (24) that $\sigma(\mu) \subset G$, where $G = A(X, E)$. Put $H = G^*$. The distribution μ considering as a distribution on G has the property

$$\{y \in H : |\widehat{\mu}(y)| = 1\} = \{0\}. \quad (25)$$

We will check that G is the desired subgroup. It follows from the structure theorem for locally compact Abelian group that $H \cong R^m \times L$, where $m \geq 0$, and L contains a compact open subgroup [7, (24.30)]. Consider the restriction of equation (3) to the connected component of zero c_L of the group L . Since the group c_L is compact, by Corollary 1 $|\widehat{\mu}(y)| = 1$ for $y \in c_L$. Taking into account (25) this implies that $c_L = \{0\}$, i.e. the group L is totally disconnected. We will prove that the group L is discrete.

By Lemma 1 the characteristic function $\widehat{\mu}(y)$ satisfies equation (3). Substitute $v_1 = u$, $v_2 = \dots = v_n = 0$ into (3). From the resulting equation we obtain

$$|\widehat{\mu}(nu)| = |\widehat{\mu}(u)|^{2n-1} |\widehat{\mu}((n-1)u)|, \quad u \in H.$$

This implies the inequality

$$|\widehat{\mu}(nu)| \leq |\widehat{\mu}(u)|^{2n-1}, \quad u \in H. \quad (26)$$

It follows from (26) that for any natural p the inequality

$$|\widehat{\mu}(n^p u)| \leq |\widehat{\mu}(u)|^{(2n-1)^p}, \quad u \in H \quad (27)$$

holds. Since $\widehat{\mu}(0) = 1$ and the function $\widehat{\mu}(y)$ is continuous, there exists a neighborhood of zero V of the group L such that $|\widehat{\mu}(y)| > 0$ for $y \in V$. Inasmuch as the group L is totally disconnected, for any neighborhood of zero of L , in particular for V , there exists a compact subgroup W such that $W \subset V$ ([7, (7.7)]). Thus, we have

$$|\widehat{\mu}(y)| > 0, \quad y \in W. \quad (28)$$

Assume that at a point $y_0 \in W$ the inequality

$$|\widehat{\mu}(y_0)| < 1 \quad (29)$$

holds. Since the group W is compact, the sequence $n^p y_0$, $p = 1, 2, \dots$ contains a converging subsequence $n^{p_j} y_0 \rightarrow \tilde{y}$, $\tilde{y} \in W$. It follows from (27) and (29) that $\hat{\mu}(\tilde{y}) = 0$, that contradicts (28). Hence, $|\hat{\mu}(y)| = 1$ for $y \in W$. Taking into account (25), we get that $W = \{0\}$, i.e. the group L is discrete. Put $K = L^*$. Since L is discrete, this implies that K is compact.

Take $y_0 \in L_{(n)}$, i.e. $ny_0 = 0$. It follows from (26) that $|\hat{\mu}(y_0)| = 1$. Taking into account (25), we obtain that $y_0 = 0$. Hence, $L_{(n)} = \{0\}$. This implies that $K^{(n)} = K$. $\square$

Lemma 10. *Let $X = R^m \times K$, where $m \geq 0$ and K is a compact group such that $K^{(n)} = K$. Then $X_{(n)} \subset c_X$.*

PROOF. We have $A(Y, Y^{(n)}) = X_{(n)}$, $A(Y, b_Y) = c_X$ ([7, (24.17)]). The lemma will be proved if we check that $Y^{(n)} \supset b_Y$. Put $L = K^*$. Obviously, b_Y is a discrete torsion group. Since $K^{(n)} = K$, we have $L_{(n)} = \{0\}$. Take $y_0 \in b_Y$. Let M be a finite cyclic subgroup generated by y_0 . It follows from $L_{(n)} = \{0\}$ that the restriction of the mapping $y \mapsto ny$, $y \in Y$, to M is a monomorphism. Taking into account that the group M is finite, this mapping is an isomorphism. Hence, $y_0 \in Y^{(n)}$. $\square$

Lemma 11. *Let X be a locally compact Abelian group. Then the following statements are equivalent:*

- (i) *for any compact subgroup K of the group X , satisfying the condition $K^{(n)} = K$, the equality $(K^*)^n = K^*$ holds;*
- (ii) *the connected component of zero c_X of the group X has the property (18);*
- (iii) *for any compact subgroup K of the group X , satisfying the condition $K^{(n)} = K$, the factor group X/K contains no subgroup topologically isomorphic to $\mathbb{T}$.*

PROOF. The equivalence of (i) and (ii) was proved in [3] (see also [4, Lemma 11.15]). To prove the equivalence of (i) and (iii) note first that $K^{(n)} = K$ if and only if $K^{(p)} = K$ for any prime divisor p of the number n . Hence it suffices to prove the equivalence of (i) and (iii) assuming that n is a prime number.

Let us prove (iii) $\Rightarrow$ (i). Let n be a prime number and K be a compact subgroup of X such that $K^{(n)} = K$. Put $L = K^*$. Then L is a discrete group satisfying the condition $L_{(n)} = \{0\}$. It follows from this that if y_0 is an element of finite order in L , then $y_0 \in L^{(n)}$. We will check that any element y_0 of infinite order in L also belongs to $L^{(n)}$. Thus, the implication (iii) $\Rightarrow$ (i) will be proved.

Let y_0 be an element of infinite order in L such that $y_0 \notin L^{(n)}$. Consider the factor-group $L/L^{(n)}$. Since n is a prime number, all nonzero elements of the factor-group $L/L^{(n)}$ have order n . It is obvious that $[y_0] \neq 0$. Therefore $[y_0]$ has

order n and hence, $ky_0 \notin L^{(n)}$, $k = 1, 2, \dots, n-1$. Denote by M the subgroup of L generated by y_0 , i.e. $M = \{y \in L : y = ly_0, l \in \mathbb{Z}\}$. Let $h \in L$ and $nh \in M$. Then $nh = ly_0$, $l \in \mathbb{Z}$. We have $l = qn + k$, where $q \in \mathbb{Z}$, $k \in \{0, 1, \dots, n-1\}$. This implies that $ky_0 = n(h - qy_0) \in L^{(n)}$. Therefore $k = 0$, and hence $nh = qny_0$. Since $L_{(n)} = \{0\}$, we get $h = qy_0 \in M$. Thus, the subgroup M has the property: if $nh \in M$, then $h \in M$. By Lemma 6 it follows from this that for the annihilator $G = A(K, M)$ the equality $G^{(n)} = G$ holds. Note now that $(K/G)^* \cong M \cong \mathbb{Z}$. This implies that $K/G \cong \mathbb{T}$. Since K/G is a subgroup of X/G , this contradicts to (iii).

The implication (i) $\Rightarrow$ (iii) for $n = 2$ was proved in [5, Lemma 7.7]. The proof in the general case is the same as in the case when $n = 2$. $\square$

PROOF OF THEOREM 1. Let $n = 2$. By Lemma 5 $\Gamma_2(X) = \Gamma_B(X)$ and the statement of the theorem in this case was proved in [2] (see also [4, Theorem 9.10]).

Assume that $n \geq 3$ and let us prove the sufficiency. Let $\mu \in \Gamma_n(X)$. By Lemma 1 the characteristic function $\hat{\mu}(y)$ satisfies equation (3). Substituting $v_1 = v$, $v_2 = -v$, $v_3 = \dots = v_n = 0$ in (3), we get

$$\hat{\mu}(u + nv)\hat{\mu}(u - nv)\hat{\mu}^{n-2}(u) = \hat{\mu}^n(u)|\hat{\mu}(nv)|^2, \quad u, v \in Y. \quad (30)$$

By Lemma 9 we can assume that the group X is of the form $X = R^m \times K$, where $m \geq 0$, and K is a compact group such that $K^{(n)} = K$. Applying Lemma 10 we obtain $X_{(n)} \subset c_X$. Then (18) implies that $X_{(n)} = \{0\}$, and hence $A(Y, X_{(n)}) = \overline{Y^{(n)}} = Y^{(n)} = Y$. Taking this into account, (30) implies that the characteristic function $\hat{\mu}(y)$ satisfies equation

$$\hat{\mu}(u + v)\hat{\mu}(u - v)\hat{\mu}^{n-2}(u) = \hat{\mu}^n(u)|\hat{\mu}(v)|^2, \quad u, v \in Y. \quad (31)$$

It follows from (31) that the set

$$B = \{y \in Y : \hat{\mu}(y) \neq 0\}$$

is an open subgroup of Y . Put $K = A(X, B)$. Since B is an open subgroup, the group K is compact. Inasmuch as the characteristic function $\hat{\mu}(y)$ satisfies equation (3), the function $\hat{\mu}(y)$ satisfies inequality (26), and it follows from (26) that the group B has the property: if $ny \in B$, then $y \in B$. Applying Lemma 6 we get $K^{(n)} = K$. By Lemma 11 the factor-group X/K contains no subgroup topologically isomorphic to $\mathbb{T}$. Note now that $(X/K)^* \cong B$ and consider the restriction of the characteristic function $\hat{\mu}(y)$ to B . By Lemma 1 this restriction

is the characteristic function of a distribution from $\Gamma_n(X/K)$. Since the factor-group X/K contains no subgroup topologically isomorphic to $\mathbb{T}$, we can apply Proposition 1 to the factor-group X/K . As a result we obtain the following representation for the characteristic function $\widehat{\mu}(y)$

$$\widehat{\mu}(y) = \begin{cases} (x, y) \exp\{-\varphi(y)\} & \text{if } y \in B \\ 0 & \text{if } y \notin B. \end{cases} \quad (32)$$

where $x \in X$, and $\varphi(y)$ is a continuous function on B , satisfying equation (2). It is well known that the function $\varphi(y)$ can be extended from the subgroup B to Y retaining its properties ([9, Lemma 5.2.5]). Denote by $\widetilde{\varphi}(y)$ the extended function. Let γ be a Gaussian measure on X with the characteristic function

$$\widehat{\gamma}(y) = (x, y) \exp\{-\widetilde{\varphi}(y)\}, \quad y \in Y. \quad (33)$$

Obviously, that $\widehat{\mu}(y) = \widehat{\gamma}(y)\widehat{m}_K(y)$. Hence, $\mu = \gamma * m_K$.

Let us prove the necessity. Assume that (18) is not fulfilled. By Lemma 11 there exists a compact subgroup K of the group X such that $K^{(n)} = K$ and the factor-group X/K contains a subgroup topologically isomorphic to $\mathbb{T}$. Therefore the distribution μ on the group $\mathbb{T}$, constructed in Lemma 2, can be considered as a distribution on the factor-group X/K . We retain the notation μ for this distribution. Since $(X/K)^* \cong A(Y, K)$, we may assume that the characteristic function $\widehat{\mu}(y)$ is defined on $A(Y, K)$. Consider on the group Y the function

$$h(y) = \begin{cases} \widehat{\mu}(y) & \text{if } y \in A(Y, K) \\ 0 & \text{if } y \notin A(Y, K). \end{cases}$$

Since the set $A(Y, K)$ is a subgroup and $\widehat{\mu}(y)$ is a positive definite function, $h(y)$ is also a positive definite function ([8, (32.43)]). Since K is a compact group, its annihilator $A(Y, K)$ is an open subgroup, and hence the function $h(y)$ is continuous. By Bochner's theorem there exists a distribution $\lambda \in M^1(X)$ such that $\widehat{\lambda}(y) = h(y)$. We will check that $\lambda \in \Gamma_n(X)$. By Lemma 1 it suffices to verify that the function $h(y)$ satisfies equation (3).

Take $u \in A(Y, K)$. If $nv_{j_0} - (v_1 + \dots + v_n) \notin A(Y, K)$ holds at least for one $j = j_0$, then $u + nv_{j_0} - (v_1 + \dots + v_n) \notin A(Y, K)$, and both sides of equation (3) are equal to zero. Assume that $nv_j - (v_1 + \dots + v_n) \in A(Y, K)$ for all $j = 1, 2, \dots, n$. This implies that $n(v_j - v_1) \in A(Y, K)$, and hence by Lemma 6 $v_j - v_1 = h_j \in A(Y, K)$, $j = 1, 2, \dots, n$. Substituting $v_j = v_1 + h_j$, $j = 1, 2, \dots, n$ in equation (3) and taking into account that the function $\widehat{\mu}(y)$ satisfies equation (3) on $A(Y, K)$, we get the equality.

Take $u \notin A(Y, K)$. Then the right-hand side of equation (3) is equal to zero. If in this case the left-hand side of equation (3) does not vanish, then the inclusions $u + nv_j - (v_1 + \dots + v_n) \in A(Y, K)$, $j = 1, 2, \dots, n$ are fulfilled. This implies that $\sum_{j=1}^n (u + nv_j - (v_1 + \dots + v_n)) = nu \in A(Y, K)$, and hence by Lemma 6 $u \in A(Y, K)$, contrary to the choice of u . So we proved that the function $h(y)$ satisfies equation (3). Since $\mu \notin \Gamma(X/K)$, obviously that $\lambda \notin \Gamma(X) * I(X)$. The theorem is completely proved. $\square$

Remark 2. Let $\mu \in \Gamma_n(\mathbb{R})$, $n \geq 2$. Put $\gamma = \mu * \bar{\mu} \in \Gamma_n(\mathbb{R})$. Then $\hat{\gamma}(y) = |\hat{\mu}(y)|^2 \geq 0$. By Lemma 1 the characteristic function $\hat{\gamma}(y)$ satisfies equation (3), and hence satisfies equation (30) too. Taking into account that $\mathbb{R}^{(n)} = \mathbb{R}$, (30) implies (31), and hence the set $B = \{y \in \mathbb{R} : \hat{\gamma}(y) \neq 0\}$ is an open subgroup of $\mathbb{R}$. So, $B = \mathbb{R}$. Since $\hat{\gamma}(y) > 0$, $y \in \mathbb{R}$, by Lemma 3 $\gamma \in \Gamma(\mathbb{R})$. This implies by Cramer's theorem that $\mu \in \Gamma(\mathbb{R})$. Thus, we proved the equality $\Gamma(\mathbb{R}) = \Gamma_n(\mathbb{R})$, $n \geq 2$, which we used in the proof of Lemma 8, and this proof is independent from Gery's theorem.

Remark 3. Comparing Proposition 1 and Theorem 1 we see that in both statements we have a particular case $n = 2$. If $n \geq 3$, then the description of the corresponding class of groups in Proposition 1 does not depend on n , and does not depend on n in Theorem 1.

Remark 4. Observe that if μ is an infinitely divisible distribution and $\mu \in \Gamma_n(X)$, then $\mu \in \Gamma(X) * I_n(X)$. Indeed, since μ is an infinitely divisible distribution, the set $B = \{y \in Y : \hat{\mu}(y) \neq 0\}$ is an open subgroup of Y ([12, p. 106]). Put $\nu = \mu * \bar{\mu}$ and consider the restriction of the characteristic function $\hat{\nu}(y)$ to B . Since $\hat{\nu}(y) = |\hat{\mu}(y)|^2 > 0$ for $y \in B$, by Lemma 3 $\hat{\nu}(y) = \exp\{-\varphi(y)\}$, $y \in B$. Note now that on an arbitrary locally compact Abelian group in the class of infinitely divisible distributions a Gaussian measure has only Gaussian factors ([5, Remark 4.8]). Thus, for the characteristic function $\hat{\mu}(y)$ we get the representation similar to (32), and the desired statement follows from this.

We would like to thank the referee for carefully reading the manuscript and useful remarks.

ACKNOWLEDGMENTS. This research is supported in part by the join project N 17-01-10 of the National Academy of Sciences of Ukraine and the Russian Foundation of the Basic Research "Investigation of asymptotic characteristic of random matrices and probability problems of classic groups and their representations".

References

- [1] G. M. FEL'DMAN, On the decomposition of Gaussian distributions on groups, *Theory Probab. Appl.* **22** (1977), 133–140.
- [2] G. M. FEL'DMAN, Bernstein Gaussian distributions on groups, *Theory Probab. Appl.* **31** (1986), 40–49.
- [3] G. M. FEL'DMAN, Groups admitting a characterization of the Gaussian distribution by the equidistribution of a monomial and linear statistics, *Ukrainian Math. J.* **42** (1990), 125–128.
- [4] G. M. FEL'DMAN, Arithmetic of Probability Distributions, and Characterization Problems on Abelian Groups, Vol. 116, Translations of Mathematical Monographs, *American Mathematical Society, Providence, RI*, 1993.
- [5] G. FELDMAN, Functional Equations and Characterization Problems on Locally Compact Abelian Groups, Vol. 5, EMS Tracts in Mathematics, *European Mathematical Society (EMS), Zurich*, 2008.
- [6] R. C. GEARY, The distribution of “Student’s” ratio for non-normal samples, *Suppl. J. R. Statist. Soc., London* **3** (1936), 178–184.
- [7] E. HEWITT and K. A. ROSS, Abstract Harmonic Analysis, **1**, *Springer-Verlag, Berlin, Göttingen, Heidelberg*, 1963.
- [8] E. HEWITT and K. A. ROSS, Abstract Harmonic Analysis, **2**, *Springer-Verlag, New York and Berlin*, 1970.
- [9] H. HEYER, Probability Measures on Locally Compact Groups, Vol. 94, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band, *Springer-Verlag, Berlin – New York*, 1977.
- [10] T. KAWATA and H. SAKAMOTO, On the characterisation of the normal population by the independence of the sample mean and the sample variance, *J. Math. Soc. Japan* **1** (1949), 111–115.
- [11] E. LUKACS, A characterization of the normal distribution, *Ann. Math. Statistics* **13** (1942), 91–93.
- [12] K. R. PARTHASARATHY, Probability measures on metric spaces, *Academic Press, New York and London*, 1967.
- [13] A. A. ZINGER, On independent samples from normal populations, *Uspehi Matem. Nauk (N.S.)* **6**, no. 5 (45) (1951), 172–175 (in Russian).

GENNADIY FELDMAN
 MATHEMATICAL DIVISION B. VERKIN INSTITUTE
 FOR LOW TEMPERATURE PHYSICS AND ENGINEERING OF
 THE NATIONAL ACADEMY OF SCIENCES OF UKRAINE
 47, LENIN AVE, KHARKOV, 61103
 UKRAINE

E-mail: feldman@ilt.kharkov.ua

MARGARYTA MYRONYUK
 MATHEMATICAL DIVISION B. VERKIN INSTITUTE
 FOR LOW TEMPERATURE PHYSICS AND ENGINEERING OF
 THE NATIONAL ACADEMY OF SCIENCES OF UKRAINE
 47, LENIN AVE, KHARKOV, 61103
 UKRAINE

E-mail: myronyuk@ilt.kharkov.ua

(Received April 15, 2009; revised October 25, 2009)