

Ricci solitons in manifolds with quasi-constant curvature

By CORNELIA LIVIA BEJAN (Iași) and MIRCEA CRASMAREANU (Iași)

This paper is dedicated to the memory of Professor Stere Ianus (1939–2010)

Abstract. The Eisenhart problem of finding parallel tensors treated already in the framework of quasi-constant curvature manifolds in [Jia] is reconsidered for the symmetric case and the result is interpreted in terms of Ricci solitons. If the generator of the manifold provides a Ricci soliton then this is i) expanding on para-Sasakian spaces with constant scalar curvature and vanishing D -concircular tensor field and ii) shrinking on a class of orientable quasi-umbilical hypersurfaces of a real projective space=elliptic space form.

1. Introduction

In 1923, EISENHART [Eisenhart] proved that if a positive definite Riemannian manifold (M, g) admits a second order parallel symmetric covariant tensor other than a constant multiple of the metric tensor, then it is reducible. In 1926, LEVY [Levy] proved that a parallel second order symmetric non-degenerated tensor α in a space form is proportional to the metric tensor. Note that this question can be considered as the dual to the the problem of finding linear connections making parallel a given tensor field; a problem which was considered by WONG in [Wong]. Also, the former question implies topological restrictions namely if the (pseudo) Riemannian manifold M admits a parallel symmetric $(0, 2)$ tensor field then M is locally the direct product of a number of (pseudo) Riemannian

Mathematics Subject Classification: 53Cxx, 53C44, 53C21, 53C20, 53C25.

Key words and phrases: parallel second order covariant tensor field, quasi-constant curvature, torse-forming vector field, Ricci soliton.

manifolds, [Wu] (cited by [Zhao]). Another situation where the parallelism of α is involved appears in the theory of totally geodesic maps, namely, as is pointed out in [Oniciuc, p. 114], $\nabla\alpha = 0$ is equivalent with the fact that $1 : (M, g) \rightarrow (M, \alpha)$ is a totally geodesic map.

While both Eisenhart and Levy work locally, RAMESH SHARMA gives in [Sharma1] a global approach based on Ricci identities. In addition to space-forms, Sharma considered this *Eisenhart problem* in contact geometry [Sharma2]–[Sharma4], for example for K -contact manifolds in [Sharma3]. Since then, several other studies appeared in various contact manifolds, see for example, the bibliography of [CalinCrasm].

Another framework was that of quasi-constant curvature in [Jia]; recall that the notion of *manifold with quasi-constant curvature* was introduced by BANG-YEN CHEN and KENTARO YANO in 1972, [ChenYano], and since then, was the subject of several and very interesting works, [Bernardini], [DeGhosh], [Wang], in both local and global approaches. Unfortunately, the paper of Jia contains some typos and we consider that a careful study deserves a new paper. There are two remarks regarding Jia result: i) it is in agreement with what happens in all previously recalled contact geometries for the symmetric case, ii) it is obtained in the same manner as in SHARMA's paper [Sharma1]. Our work improves the cited paper with a natural condition imposed to the generator of the given manifold, namely to be of torse-forming type with a regularity property.

Our main result is connected with the recent theory of Ricci solitons [Cao], a subject included in the Hamilton–Perelman approach (and proof) of Poincaré Conjecture. A connection between Ricci flow and quasi-constant curvature manifolds appears in [CaiZhao]; thus our treatment for Ricci solitons in quasi-constant curvature manifolds seems to be new.

Our work is structured as follows. The first section is a very brief review of manifolds with quasi-constant curvature and Ricci solitons. The next section is devoted to the (symmetric case of) Eisenhart problem in our framework and the relationship with the Ricci solitons is pointed out. A technical condition appears, which we call *regularity*, and is concerning with the non-vanishing of the Ricci curvature with respect to the generator of the given manifold. Let us remark that in the Jia's paper this condition is involved, but we present a characterization of these manifolds as well as some remarkable cases which are out of this condition namely: quasi-constant curvature locally symmetric and Ricci semi-symmetric metrics. A characterization of Ricci soliton is derived for dimension greater than 3.

Four concrete examples involved in possible Ricci solitons on quasi-constant manifolds are listed at the end. For the second example, we pointed out some

consequences which are yielded by the hypothesis of compactity, used in paper [DragomirGrimaldi], in connection with (classic by now) papers of T. Ivey and Perelman.

2. Quasi-constant curvature manifolds. Ricci solitons

Fix a triple (M, g, ξ) with M_n a smooth $n(> 2)$ -dimensional manifold, g a Riemannian metric on M and ξ an unitary vector field on M . Let η the 1-form dual to ξ with respect to g .

If there exist two smooth functions $a, b \in C^\infty(M)$ such that:

$$\begin{aligned} R(X, Y)Z &= a[g(Y, Z)X - g(X, Z)Y] + b[g(Y, Z)\eta(X) - g(X, Z)\eta(Y)]\xi \\ &\quad + b\eta(Z)[\eta(Y)X - \eta(X)Y] \end{aligned} \quad (2.1)$$

then (M, g, ξ) is called *manifold of quasi-constant curvature* and ξ is *the generator*, [ChenYano]. Using the notation of [DragomirTomassini, p. 325] let us denote $M_{a,b}^n(\xi)$ this manifold.

It follows:

$$R(X, Y)\xi = (a + b)[\eta(Y)X - \eta(X)Y] \quad (2.2)$$

$$R(X, \xi)Z = (a + b)[\eta(Z)X - g(X, Z)\xi] \quad (2.3)$$

while the Ricci curvature $S(X, Y) = \text{Tr}(Z \rightarrow R(Z, X)Y)$ is:

$$S(X, Y) = [a(n - 1) + b]g(X, Y) + b(n - 2)\eta(X)\eta(Y) \quad (2.4)$$

which means that (M, g, ξ) is an *eta-Einstein manifold*; in particular, if a, b are scalars, then (M, g, ξ) is an *quasi-Einstein manifold*, [GhoshDeBinh]. The scalar curvature is:

$$r = (n - 1)(na + 2b), \quad (2.5)$$

and we derive:

$$a = \frac{r - 2S(\xi, \xi)}{(n - 1)(n - 2)}, \quad b = \frac{nS(\xi, \xi) - r}{(n - 1)(n - 2)}. \quad (2.6)$$

Then $a + b = \frac{S(\xi, \xi)}{n - 1}$. Let us consider also the Ricci $(1, 1)$ tensor field Q given by: $S(X, Y) = g(QX, Y)$. From (2.4) we get:

$$Q(X) = [a(n - 1) + b]X + b(n - 2)\eta(X)\xi \quad (2.7)$$

which yields:

$$Q(\xi) = (a + b)(n - 1)\xi \quad (2.8)$$

and then ξ is an eigenvalue of Q .

In the last part of this section we recall the notion of Ricci solitons according to [Sharma5, p. 139]. On the manifold M , a *Ricci soliton* is a triple (g, V, λ) with g a Riemannian metric, V a vector field and λ a real scalar such that:

$$\mathcal{L}_V g + 2S + 2\lambda g = 0. \quad (2.9)$$

The Ricci soliton is said to be *shrinking*, *steady* or *expanding* according as λ is negative, zero or positive.

Also, we adopt the notion of η -*Ricci soliton* introduced in the paper [ChoKimura] as a data (g, V, λ, μ) :

$$\mathcal{L}_V g + 2S + 2\lambda g + 2\mu\eta \otimes \eta = 0. \quad (2.10)$$

3. Parallel symmetric second order tensors and Ricci solitons

Fix α a symmetric tensor field of $(0, 2)$ -type which we suppose to be parallel with respect to the Levi-Civita connection ∇ i.e. $\nabla\alpha = 0$. Applying the Ricci identity $\nabla^2\alpha(X, Y; Z, W) - \nabla^2\alpha(X, Y; W, Z) = 0$ we obtain the relation (1.1) of [Sharma1, p. 787]:

$$\alpha(R(X, Y)Z, W) + \alpha(Z, R(X, Y)W) = 0 \quad (3.1)$$

which is fundamental in all papers treating this subject. Replacing $Z = W = \xi$ and using (2.2) it results, by the symmetry of α :

$$(a + b)[\eta(Y)\alpha(X, \xi) - \eta(X)\alpha(Y, \xi)] = 0. \quad (3.2)$$

Definition 3.1. $M_{a,b}^n(\xi)$ is called *regular* if $a + b \neq 0$.

In order to obtain a characterization of such manifolds we consider:

Definition 3.2 ([RachunekMikes]). ξ is called *semi-torse forming vector field* for (M, g) if, for all vector fields X :

$$R(X, \xi)\xi = 0. \quad (3.3)$$

From (2.2) we get: $R(X, \xi)\xi = (a + b)(X - \eta(X)\xi)$ and therefore, if $X \in \ker \eta = \xi^\perp$, then $R(X, \xi)\xi = (a + b)X$ and we obtain:

Proposition 3.3. For $M_{a,b}^n(\xi)$ the following are equivalent:

- i) ξ is regular,
- ii) ξ is not semi-torse forming,
- iii) $S(\xi, \xi) \neq 0$ i.e. ξ is non-degenerate with respect to S ,
- iv) $Q(\xi) \neq 0$ i.e. ξ does not belong to the kernel of Q .

In particular, if ξ is parallel ($\nabla \xi = 0$) then M is not regular.

Remarks 3.4. i) From Theorems 2 and 3 of [Wang, p. 175] a regular $M_{a,b}^n(\xi)$ is neither recurrent nor locally symmetric.

ii) From Theorem 3 of [DragomirGrimaldi, p. 228] a regular $M_{a,b}^n(\xi)$ with a and b constants is not Ricci semi-symmetric.

In the following we restrict to the regular case. Returning to (3.2), with $X = \xi$ in:

$$\eta(Y)\alpha(X, \xi) = \eta(X)\alpha(Y, \xi) \quad (3.4)$$

we derive:

$$\alpha(Y, \xi) = \eta(Y)\alpha(\xi, \xi) = \alpha(\xi, \xi)g(Y, \xi). \quad (3.5)$$

The parallelism of α implies also that $\alpha(\xi, \xi)$ is a constant:

$$X(\alpha(\xi, \xi)) = 2\alpha(\nabla_X \xi, \xi) = 2\alpha(\xi, \xi)g(\nabla_X \xi, \xi) = 2\alpha(\xi, \xi) \cdot 0 = 0. \quad (3.6)$$

Making $Y = \xi$ in (3.1) and using (2.3) we get:

$$\eta(Z)\alpha(X, W) - g(X, Z)\alpha(\xi, W) + \eta(W)\alpha(X, Z) - g(X, W)\alpha(\xi, Z) = 0$$

which yield, via (3.5) and $W = \xi$:

$$\alpha(X, Z) = \alpha(\xi, \xi)g(X, Z). \quad (3.7)$$

In conclusion:

Theorem 3.5. *A parallel second order symmetric covariant tensor in a regular $M_{a,b}^n(\xi)$ is a constant multiple of the metric tensor.*

At the end of this section we include some applications of the above Theorem to Ricci solitons:

Naturally, two remarkable situations appear regarding the vector field V : $V \in \text{span } \xi$ or $V \perp \xi$ but the second class seems far too complex to analyse in practice. For this reason it is appropriate to investigate only the case $V = \xi$. So, we can apply the previous result for $\alpha := \mathcal{L}_\xi g + 2S$ which yields $\lambda = -S(\xi, \xi)$.

Theorem 3.6. *Fix a regular $M_{a,b}^n(\xi)$.*

- i) *A Ricci soliton $(g, \xi, -S(\xi, \xi) \neq 0)$ can not be steady but is shrinking if the constant $S(\xi, \xi)$ is positive or expanding if $S(\xi, \xi) < 0$.*

- ii) An η -Ricci soliton (g, ξ, λ, μ) provided by the parallelism of $\alpha + 2\mu\eta \otimes \eta$ is given by:

$$\lambda + \mu = -S(\xi, \xi) \neq 0. \quad (3.8)$$

- iii) If $n \geq 4$ and $b \neq 0$ then $(g, \xi, -S(\xi, \xi))$ is a Ricci soliton if and only if ξ is geodesic i.e. $\nabla_\xi \xi = 0$ and:

$$\frac{\xi(a+b)}{4b} + a(n-1) + b = \frac{a+b}{n-1}. \quad (3.9)$$

PROOF. iii) We have three cases:

- I) $\alpha + 2\lambda g = 0$ on $\text{span } \xi$ yields the above expression of λ .
 II) $\alpha + 2\lambda g = 0$ on $\ker \eta = \xi^\perp$ gives:

$$\frac{\xi(a+b)}{4b} + \lambda + a(n-1) + b = 0 \quad (3.10)$$

where we use the formula (3.5) of [GanchevMihova, p. 123].

- III) $\alpha + 2\lambda g = 0$ on $(U, \xi) \in \ker \eta \oplus \text{span } \xi$ gives:

$$g(\nabla_U \xi, \xi) + g(U, \nabla_\xi \xi) = 0.$$

But the first term is zero since ξ is unitary while the second implies that $\nabla_\xi \xi \in \text{span } \xi$. But again, ξ being unitary we have that $\nabla_\xi \xi$ is orthogonal to ξ . $\square$

Example 3.7. A para-Sasakian manifold with constant scalar curvature and vanishing D -concircular tensor is an $M_{a,b}^n(\xi)$ with [DragomirGrimaldi, p. 186]:

$$a = \frac{r + 2(n-1)}{(n-1)(n-2)}, \quad b = \frac{-r - n(n-1)}{(n-1)(n-2)}$$

and then, a Ricci soliton (g, ξ) on it is expanding. This result can be considered as a version in para-contact geometry of the Corollary of [Sharma5, p. 140] which states that a Ricci soliton g of a compact K -contact manifold is Einstein, Sasakian and shrinking.

From (3.9) we get $r = -n$ and returning to formulae above it results:

$$a = \frac{1}{n-1}, \quad b = \frac{-n}{n-1}.$$

Example 3.8. Let $N_{n+1}(c)$ be a space form with the metric g and M a *quasi-umbilical hypersurface* in N , [ChenYano], [Wang, p. 175], i.e. there exist two smooth functions α, β on M and a 1-form η of norm 1 such that the second fundamental form is:

$$h_{ij} = \alpha g_{ij} + \beta \eta_i \eta_j.$$

According to the cited papers M is an $M_{a,b}^n(\xi)$ with:

$$a = c + \alpha^2, \quad b = \alpha\beta$$

and ξ the g -dual of η . This $M_{a,b}^n(\xi)$ is regular if and only if $c + \alpha^2 + \alpha\beta \neq 0$. Therefore, a Ricci soliton (g, ξ) on this $M_{a,b}^n(\xi)$ is shrinking if $c + \alpha^2 + \alpha\beta > 0$ and expanding if $c + \alpha^2 + \alpha\beta < 0$.

Inspired by Theorem 3 of [DragomirGrimaldi, p. 185] let $N = \mathbb{R}P^{n+1}(c)$, $c > 0$ and M an orientable quasi-umbilical hypersurface with $b = \alpha\beta > 0$. Then:

- i) a Ricci soliton (g, ξ) on it is shrinking and M is a real homology sphere (all Betti numbers vanish) if it is also compact,
- ii) using the result of [Ivey], for $n = 3$ the manifold is of constant curvature being compact; so the case $n = 4$ is the first important in any conditions or the case $n = 3$ without compactness when we (possibly) give up at the topology of real homology sphere,
- iii) using again a classic result, now due to Perelman [Perelman], the compactness implies that the Ricci soliton is gradient i.e. η is exact.

Example 3.9. Let (M_0^{2n}, ω_0, B) be a generalized Hopf manifold [DragomirOrnea], and M^n an n -dimensional anti-invariant and totally geodesic submanifold. We set $\|\omega_0\| = 2c$ and suppose that B is unitary. Then, formula (12.40) of [DragomirOrnea, p. 162] gives that if $R^\perp = 0$ then M^n is of quasi-constant curvature with $a = c^2$ and $b = -\frac{1}{4}$. Therefore, M^n is regular for $\|\omega_0\| \neq 1$ and a Ricci soliton is shrinking if $\|\omega_0\| > 1$ and expanding if $\|\omega_0\| < 1$.

Example 3.10. Suppose that ξ is a *torse-forming vector field* i.e. there exist a smooth function f and a 1-form ω such that:

$$\nabla_X \xi = fX + \omega(X)\xi. \quad (3.11)$$

From the fact that ξ has unitary length it results $f + \omega(\xi) = 0$ which means that ξ is exactly a geodesic vector field.

Particular cases:

- i) ([RachunekMikes]) If ω is exact then ξ is called *conccircular*; let $\omega = -du$ with u a smooth function on M . Then $f = -\omega(\xi) = \xi(u)$.
- ii) If $\omega = -f\eta$ then we call ξ of *Kenmotsu type* since (3.11) becomes similar to a expression well-known in Kenmotsu manifolds, [CalinCrasm].

Let us restrict to ii). From (3.11) a straightforward computation gives:

$$R(X, Y)\xi = X(f)[Y - \eta(Y)\xi] - Y(f)[X - \eta(X)\xi] + f^2[\eta(X)Y - \eta(Y)X] \quad (3.12)$$

and a comparison with (2.2) yields $a + b = -f^2$ and f must be a constant, different from zero from regularity of the manifold. So, a possible Ricci soliton in a Kenmotsu type case must be expanding and with $S(\xi, \xi)$ and the scalar curvature constants, a result similar to Propositions 3 and 4 of [CalinCrasm].

References

- [Bernardini] ADELE BERNARDINI, Riemannian manifolds with quasicontant curvature, *Atti Sem. Mat. Fis. Univ. Modena* **30** (1981), 16–31.
- [CaiZhao] Q. H. CAI and P. B. ZHAO, Stability of quasi-DeTurck flows in Riemannian manifolds of quasi-constant sectional curvatures, *Chin. Ann. Math. Ser. A* **29** (2008), 97–106.
- [Cao] H.-D. CAO, Geometry of Ricci solitons, *Chinese Ann. Math. Ser. B* **27** (2006), 121–142.
- [CalinCrasm] C. CALIN and M. CRASMAREANU, From the Eisenhart problem to Ricci solitons in f -Kenmotsu manifolds, *Bull. Malays. Math. Sci. Soc.* **33** (2010), 361–368.
- [ChenYano] B.-Y. CHEN and K. YANO, Hypersurfaces of a conformally flat space, *Tensor* **26** (1972), 318–322.
- [ChoKimura] J. T. CHO and M. KIMURA, Ricci solitons and real hypersurfaces in a complex space form, *Tohoku Math. J.* **61** (2009), 205–212.
- [DeGhosh] U. C. DE and G. C. GHOSH, On quasi Einstein manifolds, *Period. Math. Hungar.* **48** (2004), 223–231.
- [DragomirGrimaldi] S. DRAGOMIR and RENATA GRIMALDI, On the topology of Riemann spaces of quasi-constant curvature, *Publ. Inst. Math. (Beograd)* **46**(60) (1989), 183–187.
- [DragomirOrnea] S. DRAGOMIR and L. ORNEA, Locally conformal Kähler geometry, Progress in Mathematics, 155, *Birkhäuser Boston, Inc., Boston, MA*, 1998.
- [DragomirTomassini] S. DRAGOMIR and G. TOMASSINI, Differential geometry and analysis on CR manifolds, Progress in Mathematics, 246, *Birkhäuser Boston, Inc., Boston, MA*, 2006.
- [Eisenhart] L. P. EISENHART, Symmetric tensors of the second order whose first covariant derivatives are zero, *Trans. Amer. Math. Soc.* **25** (1923), 297–306.
- [GanchevMihova] G. GANCHEV and V. MIHOVA, Riemannian manifolds of quasi-constant sectional curvatures, *J. Reine Angew. Math.* **522** (2000), 119–141.
- [GhoshDeBinh] G. C. GHOSH, U. C. DE and T. Q. BINH, Certain curvature restrictions on a quasi Einstein manifold, *Publ. Math. Debrecen* **69** (2006), 209–217.

- [Ivey] T. IVEY, Ricci solitons on compact three-manifolds, *Differential Geom. Appl.* **3** (1993), 301–307.
- [Jia] X.-Q. JIA, Second order parallel tensors on quasi-constant curvature manifolds, *Chinese Quart. J. Math.* **17** (2002), 101–105.
- [Levy] H. LEVY, Symmetric tensors of the second order whose covariant derivatives vanish, *Ann. of Math.* (2) **27** (1925), 91–98.
- [Oniciuc] C. ONICIUC, Nonlinear connections on tangent bundle and harmonicity, *Ital. J. Pure Appl. Math.* **6** (1999), 109–122.
- [Perelman] G. PERELMAN, The entropy formula for the Ricci flow and its geometric applications, 2002, [arXiv: math.DG/0211159v1](https://arxiv.org/abs/math.DG/0211159v1).
- [RachunekMikes] L. RACHUNEK and J. MIKES, On tensor fields semiconjugated with torsion-forming vector fields, *Acta Univ. Palacki. Olomuc. Fac. Rerum Nat. Math.* **44** (2005), 151–160.
- [Sharma1] R. SHARMA, Second order parallel tensor in real and complex space forms, *Internat. J. Math. Math. Sci.* **12** (1989), 787–790.
- [Sharma2] R. SHARMA, Second order parallel tensors on contact manifolds. I, *Algebras Groups Geom.* **7** (1990), 145–152.
- [Sharma3] R. SHARMA, Second order parallel tensors on contact manifolds. II, *C. R. Math. Rep. Acad. Sci. Canada* **13** (1991), 259–264.
- [Sharma4] R. SHARMA, On the curvature of contact metric manifolds, *J. Geom.* **53** (1995), 179–190.
- [Sharma5] R. SHARMA, Certain results on K -contact and (k, μ) -contact manifolds, *J. Geom.* **89** (2008), 138–147.
- [Wang] Y. D. WANG, On some properties of Riemannian spaces of quasiconstant curvature, *Tensor* **35** (1981), 173–176.
- [Wong] Y.-C. WONG, Existence of linear connections with respect to which given tensor fields are parallel or recurrent, *Nagoya Math. J.* **24** (1964), 67–108.
- [Wu] H. WU, Holonomy groups of indefinite metrics, *Pacific J. Math.* **20** (1967), 351–392.
- [Zhao] G. ZHAO, Symmetric covariant tensor fields of order 2 on pseudo-Riemannian manifolds, *Vienna Preprint ESI 479* (1997), <http://www.esi.ac.at/preprints/esi479.ps>.

CORNELIA LIVIA BEJAN
SEMINARUL MATEMATIC "AL. MYLLER",
UNIVERSITY "AL. I. CUZA"
IASI, 700506
ROMANIA

E-mail: bejanliv@yahoo.com

MIRCEA CRASMAREANU
FACULTY OF MATHEMATICS
UNIVERSITY "AL. I. CUZA"
IASI, 700506
ROMANIA

E-mail: mcrasm@uaic.ro

URL: <http://www.math.uaic.ro/~mcrasm>

(Received January 4, 2010)