

The residual nilpotency of the augmentation ideal

By B. KIRÁLY (Eger)

1. Introduction

Let R be a commutative ring with unity, G a group and RG its group ring and let $A(RG)$ denote the *augmentation ideal* of RG , that is the kernel of the ring homomorphism $RG \rightarrow R$ which maps the group elements to 1. It is easy to see that an R -module $A(RG)$ is a free module with the elements $g - 1$ ($1 \neq g \in G$) as a basis.

The ideal $A(RG)$ of the group ring RG is said to be *residually nilpotent* if $\bigcap_{n=1}^{\infty} A^n(RG) = 0$. For convenience, we adopt the following notation:

$$A^\omega(RG) = \bigcap_{n=1}^{\infty} A^n(RG).$$

We are interested in the residual nilpotence of the augmentation ideals of group rings.

In the case when R is a field, the question about the residual nilpotence of the augmentation ideal is completely solved (see in particular [6], VI, Theorem 2.26).

If R is the ring of integers and if the finitely generated group G has torsion elements then the question about the residual nilpotency of the augmentation ideal is solved in [1].

In [5] the author gives a complete characterization of the residual nilpotence of the augmentation ideal for a group ring over the integers.

For a group ring RG HARTLEY (see [3], Theorem E) gives a sufficient condition for the residual nilpotence of the augmentation ideal in the case when $\bigcap_{n=1}^{\infty} p^n R = 0$ and the group G has a finite N series.

In this paper we give sufficient conditions for the residual nilpotence of the augmentation ideal for all arbitrary group ring RG (Theorem 3.1). These conditions are also necessary if the group G has a generalized torsion element (Theorem 3.2). In the case when G is without a generalized torsion element and the torsion group of the additive group of a ring R for some prime p has no elements of infinite p -height, the question about the residual nilpotence of the augmentation ideal is solved in Theorem 3.3.

2. Notations and some known facts

If H is a normal subgroup of G , then $I(RH)$ (or $I(H)$ for short when it is obvious from the context what ring R we are working with) denotes the ideal of RG generated by all elements of the form $h - 1$, $h \in H$. It is well known that $I(RH)$ is the kernel of the natural epimorphism $\psi^* : RG \rightarrow RG/H$ induced by the group homomorphism ψ of G onto G/H .

If K, L are two subgroups of G , then $[K, L]$ denotes the subgroup generated by all commutators $[g, h] = g^{-1}h^{-1}gh$, $g \in K$, $h \in L$.

Let p be a prime and n a natural number. Then G^{p^n} is the subgroup generated by all elements of the form g^{p^n} , $g \in G$.

The subgroup $W_p(G)$ defined by $W_p(G) = \bigcap_{n=1}^{\infty} G^{p^n} \gamma_n(G)$, where $\gamma_n(G)$ is the n -th term of the lower central series of G , i.e. $\gamma_1(G) = G$, $\gamma_2(G) = G'$ is the commutator subgroup $[G, G]$ of G , and $\gamma_n(G) = [\gamma_{n-1}(G), G]$.

The ideal $J_p(R)$ of a ring R is defined by $J_p(R) = \bigcap_{n=1}^{\infty} p^n R$.

If $\mathfrak{C}$ denotes a class of groups (by which we understand that $\mathfrak{C}$ contains all groups of order 1 and, with each $H \in \mathfrak{C}$, all isomorphic copies of H), we define the class $R\mathfrak{C}$ of *residually- $\mathfrak{C}$* groups by letting $G \in R\mathfrak{C}$ if and only if: whenever $1 \neq g \in G$, there exists a normal subgroup H_g of the group G such that $G/H_g \in \mathfrak{C}$ and $g \notin H_g$.

We use the following notations for standard group classes: $\mathfrak{N}_0$ – torsion-free nilpotent groups, $\mathfrak{N}_p$ – nilpotent p -groups of finite exponent, that is, nilpotent groups in which for some $n = n(G)$ every element g satisfies the equation $g^{p^n} = 1$.

Let $\mathfrak{C}$ be a class of groups. A group G is said to be *discriminated* by $\mathfrak{C}$ if for every finite subset $g_1, g_2, \dots, g_n$ of distinct elements of G , there exists a group $H \in \mathfrak{C}$ and a homomorphism ϕ of G into H , such that $\phi(g_i) \neq \phi(g_j)$ for $g_i \neq g_j$, ($1 \leq i, j \leq n$).

A series $G = H_1 \supseteq H_2 \supseteq \dots \supseteq H_n \supseteq \dots$ of normal subgroups of a group G is called an *N -series* if $[H_i, H_j] \subseteq H_{i+j}$ for all $i, j \geq 1$ and also each of the Abelian groups H_i/H_j is a direct product of (possibly infinitely

many) cyclic group which are either infinite or of order p^k , where p is a fixed prime and k bounded by some integer depending only on G .

It is easy to see that the lower central series of a nilpotent p -group of finite exponent is an N -series.

The n -th dimension subgroup $D_n(RG)$ of G over R is the set of group elements $g \in G$ such that $g - 1$ lies in the n -th power of $A(RG)$. It is well known that for every natural number n the inclusion

$$\gamma_n(G) \subseteq D_n(RG)$$

holds.

An element g of a group G is called a *generalized torsion element* if for all natural numbers n the order of the element $g\gamma_n(G)$ of the factor group $G/\gamma_n(G)$ is finite.

It is clear that the torsion elements of the group G are generalized torsion elements of G .

If $g \in G$ is a generalized torsion element then Ω_g denotes the set of prime divisors of the orders of the elements $g\gamma_n(G) \in G/\gamma_n(G)$ for all $n = 2, 3, \dots$.

Lemma 2.1. *For every natural number n the inclusions*

$$I(\gamma_n(G)) \subseteq I(D_n(RG)) \subseteq A^n(RG)$$

hold.

The statement is well known.

Lemma 2.2. *Let G be discriminated by a class of groups $\mathfrak{C}$ and let x be a nonzero element of RG . Then there exists a group $H \in \mathfrak{C}$ and a homomorphism ϕ of RG into RH such that $\phi(x) \neq 0$.*

The proof is evident.

Lemma 2.3. *If G is discriminated by a class of groups $\mathfrak{C}$ and for each $H \in \mathfrak{C}$ the equality $A^\omega(RH) = 0$ holds, then $A^\omega(RG) = 0$.*

The lemma follows immediately from Lemma 2.2.

Lemma 2.4. *Let a class $\mathfrak{C}$ of groups be closed for the taking of subgroups (that is all subgroups of any member of the class $\mathfrak{C}$ are again in the class $\mathfrak{C}$) and also for finite direct products and let G be a residually- $\mathfrak{C}$ group. Then G is discriminated by $\mathfrak{C}$.*

The proof can be obtained immediately.

In this paper we shall use the following theorems:

Theorem 2.1 ([3], Theorem E). *Let G be a group with a finite N -series and R be a commutative ring with unity satisfying $J_p(R) = 0$. Then $A^\omega(RG) = 0$.*

Theorem 2.2. ([6], VI, Theorem 2.15). *If G is a residually torsion-free nilpotent group and R is a commutative ring with unity such that its additive group is torsion-free, then $A^\omega(RG) = 0$.*

3. The residual nilpotency of the augmentation ideal

In this section R is a commutative ring with unity.

Lemma 3.1. *Let $g \in G$ and $g^{p^n} \in D_t(RG)$ for a prime p and a natural number n . Then there exists a natural number m such that*

$$p^m(g-1) \in A^t(RG).$$

PROOF. We prove this by induction on t . For $t = 1$ the statement is obvious. Let $p^s(g-1) \in A^{t-1}(RG)$ for some s . From the decomposition g^{p^n} as $(g-1+1)^{p^n}$ we have that

$$g^{p^m} - 1 = p^m(g-1) + \sum_{i=2}^{t-1} \binom{p^m}{i} (g-1)^i + \sum_{i=t}^{p^m} \binom{p^m}{i} (g-1)^i$$

for every m . If $m \geq n(s+t)$, then p^s divides $\binom{p^m}{i}$ ($\binom{p^m}{i}$ is the binomial coefficient p^m over i) for $i = 1, 2, \dots, t-1$ and $g^{p^m} \in D_t(RG)$. Therefore we have

$$g^{p^m} - 1 = p^m(g-1) + p^s(g-1)^2 \sum_{i=2}^{t-1} d_i (g-1)^{i-2} + \sum_{i=t}^{p^m} \binom{p^m}{i} (g-1)^i$$

where $d_i p^s = \binom{p^m}{i}$ for $i=2, 3, \dots, t-1$. By Lemma 2.1 $g^{p^m} - 1 \in A^t(RG)$. Then from the induction hypothesis and from the preceding identity $p^m(g-1) \in A^t(RG)$ follows. $\square$

Lemma 3.2. *Let $h \in G^{p^n} \gamma_n(G)$ for a natural number n . Then for all natural numbers t and s for which $n \geq t+s$*

$$h - 1 \equiv p^s F_t(h) \pmod{A^t(RG)}$$

holds, where $F_t(h) \in A(RG)$.

PROOF. Writing the element h as $h = h_1^{p^n} h_2^{p^n} \dots h_m^{p^n} y_n$ ($h_i \in G$, $y_n \in \gamma_n(G)$) and using the identity

$$(1) \quad xy - 1 = (x-1)(y-1) + (x-1) + (y-1)$$

we have

$$h - 1 = \left(h_1^{p^n} h_2^{p^n} \cdots h_m^{p^n} - 1 \right) (y_n - 1) + \left(h_1^{p^n} h_2^{p^n} \cdots h_m^{p^n} - 1 \right) + (y_n - 1).$$

Since $t \leq n$, by Lemma 2.1, $(y_n - 1) \in A^t(RG)$. It is clear that p^s divides $\binom{p^n}{j}$ for $j = 1, 2, \dots, t-1$. Then from the preceding identity

$$\begin{aligned} h - 1 &\equiv \sum_{i=1}^m \left(h_i^{p^n} - 1 \right) b_i \equiv p^s \sum_{i=1}^m \sum_{j=1}^{t-1} d_j (h_i - 1)^j b_i \equiv \\ &\equiv p^s F_t(h) \pmod{A^t(RG)} \end{aligned}$$

follows, where $F_t(h) = \sum_{i=1}^m \sum_{j=1}^{t-1} d_j (h_i - 1)^j b_i$, $b_i \in RG$ and $p^s d_j = \binom{p^n}{j}$. $\square$

We recall that if g is a generalized torsion element of a group G then Ω_g is the set of the prime divisors of the orders of the elements $g\gamma_k G \in G/\gamma_k(G)$ for all $k = 2, 3, \dots$ and also that $J_p(R) = \bigcap_{n=1}^{\infty} p^n R$.

Lemma 3.3. *Let g be a generalized torsion element of a group G , Λ an arbitrary subset of Ω_g , $r \in \bigcap_{p \in \Lambda} J_p(R)$ and let $x \in \bigcap_{p \in \Omega_g \setminus \Lambda} \bigcap_{n=1}^{\infty} I(G^{p^n} \gamma_n(G))$.*

Then one of the following statement holds:

- 1) *if Λ is the proper subset of Ω_g , then $r(g-1)x \in A^\omega(RG)$;*
- 2) *if $\Lambda = \Omega_g$, then $r(g-1) \in A^\omega(RG)$;*
- 3) *if $\Lambda = \emptyset$, then $(g-1)x \in A^\omega(RG)$.*

PROOF. Clearly, it is enough to show that for an arbitrary natural number t the elements $r(g-1)$, $(g-1)x$, $r(g-1)x$ all lie in the ideal $A^t(RG)$.

If $g \in \gamma_t(G)$ then, by Lemma 2.1, $(g-1) \in A^t(RG)$ and the statements follow.

Now let $g \notin \gamma_t(G)$ and let $n_t = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_s^{\alpha_s}$ be the order of the element $g\gamma_t(G)$ of $G/\gamma_t(G)$. We can renumber the primes so that $p_1, p_2, \dots, p_k \in \Lambda$ and $p_i \notin \Lambda$ for all $i > k$.

Let $g\gamma_t(G) = g_1 g_2 \cdots g_s \gamma_t(G)$ be a decomposition of the element $g\gamma_t(G)$ of the nilpotent group $G/\gamma_t(G)$ into a product of p_i -elements $g_i \gamma_t(G)$ such that $g_i^{q_i} \in \gamma_t(G)$, where $q_i = p_i^{\alpha_i}$, $i = 1, 2, \dots, s$. Then $g = g_1 g_2 \cdots g_s y_t$ for suitable $y_t \in \gamma_t(G)$. From (1) we conclude that

$$g - 1 = v + w + (y_t - 1)$$

where $v = \sum_{i=1}^k (g_i - 1)a_i$, $w = \sum_{i=k+1}^s (g_i - 1)a_i$ and $a_i \in RG$.

(If $\Lambda \cap \{p_1, p_2, \dots, p_s\} = \emptyset$ we assume that $v = 0$, $k = 0$, and in the case $\Lambda \cap \{p_1, p_2, \dots, p_s\} = \{p_1, p_2, \dots, p_s\}$ we assume that $w = 0$ and $k = s$.) Since $y_t - 1 \in A^t(RG)$,

$$(2) \quad g - 1 \equiv v + w \pmod{A^t(RG)}.$$

We know that $g_i^{q_i} \in \gamma_t(G)$ holds for all $i = 1, 2, \dots, s$. Therefore $g_i^{q_i} \in D_t(RG)$ and, by Lemma 3.1, there exist natural numbers m_i for $i = 1, 2, \dots, s$ such that

$$(3) \quad p_i^{m_i}(g_i - 1) \in A^t(RG).$$

We notice that if $i \leq k$ then $p_i \in \Lambda$. So, for all $i = 1, 2, \dots, k$, we can decompose the element r from the ideal $\bigcap_{p \in \Lambda} J_p(R)$ as $r = p_i^{m_i} r_i$ ($r_i \in R$).

Therefore $r(g - 1) \equiv \sum_{i=1}^k r_i p_i^{m_i} (g_i - 1)a_i + rw \pmod{A^t(RG)}$. Then from (2) and (3) we obtain that

$$(4) \quad r(g - 1) \equiv rw \pmod{A^t(RG)}.$$

Now we prove that every component $(g_i - 1)a_i x$ of the sum $wx = \sum_{i=k+1}^s (g_i - 1)a_i x$ lies in the ideal $A^t(RG)$. Let p_j be a fixed prime, where $k + 1 \leq j \leq s$. Then $p_j \in \Omega_g \setminus \Lambda$ and from the conditions

$x \in \bigcap_{p \in \Omega_g \setminus \Lambda} \bigcap_{n=1}^{\infty} I(G^{p^n} \gamma_n(G))$ of our lemma it follows that

$x \in \bigcap_{n=1}^{\infty} I(G^{q^n} \gamma_n(G))$, where $q = p_j$. For every natural number n we

can decompose the element x as $x = \sum_{m=1}^{\ell} \beta_m z_m (f_m - 1)$, where $\beta_m \in R$,

$f_m \in G^{q^n} \gamma_n(G)$, and z_m are the elements from a left transversal of the cosets of $G^{q^n} \gamma_n(G)$ in G . If $n \geq m_j + t$ then, by Lemma 3.2, we have that

$$f_m - 1 \equiv p_j^{m_j} F_t(f_m) \pmod{A^t(RG)}$$

where $F_t(f_m) \in A(RG)$, and so

$$(5) \quad x \equiv p_j^{m_j} \sum_{m=1}^{\ell} \beta_m z_m F_t(f_m) \pmod{A^t(RG)}$$

holds. Then by (3) we obtain that

$$(g_j - 1)a_jx \equiv p_j^{m_j}(g_j - 1)a_j \sum_{m=1}^{\ell} \beta_m z_m F_t(f_m) \equiv 0 \pmod{A^t(RG)}$$

i.e. $(g_j - 1)a_jx \in A^t(RG)$ for all $j = k + 1, k + 2, \dots, s$. Thus we have

$$(6) \quad wx = 0 \pmod{A^t(RG)}$$

and

$$(7) \quad rwx = 0 \pmod{A^t(RG)}.$$

If Λ is a proper subset of Ω_g (case 1) then by (4) and (7) we get $r(g - 1)x \in A^t(RG)$.

If $\Lambda = \Omega_g$ (case 2) then $w = 0$ in (2) and by (4) we obtain that $r(g - 1)x \in A^t(RG)$.

If $\Lambda = \emptyset$ (case 3) then $v = 0$ in (2) and by (6) $(g - 1)x \in A^t(RG)$ follows. Because t is arbitrary, from the above facts it follows that the elements $r(g - 1)x$, $r(g - 1)$ and $(g - 1)x$ lie in $A^\omega(RG)$, which proves the lemma. $\square$

Let G be a nilpotent p -group of finite exponent. It is clear that its lower central series is an N -series.

We now prove the following

Lemma 3.4. *For a nilpotent p -group G of finite exponent*

$$A^\omega(RG) \subseteq J_p(R) \cdot A(RG).$$

PROOF. Let $x = \sum_{i=1}^n \alpha_i g_i$ be an element of RG and let $R_p = R/J_p(R)$. Then $J_p(R_p) = 0$ and, by Theorem 2.1, $A^\omega(R_p G) = 0$. Let $\bar{\phi}(x) = \sum_{i=1}^n \phi(\alpha_i) g_i$ where ϕ is the natural homomorphism of R onto R_p . Then $\bar{\phi}$ is a homomorphism of RG onto $R_p G$. If $x \in A^\omega(RG)$ then $\bar{\phi}(x)$ lies in $A^\omega(R_p G)$. Consequently, $\bar{\phi}(x) = 0$ and $\alpha_i \in J_p(R)$ for $i = 1, 2, \dots, n$. $\square$

Let Ω be a nonempty subset of the set of primes and let $\mathfrak{N}_p$ be the class of nilpotent p -groups of finite exponent. Define $\mathfrak{N}_\Omega$ by $\mathfrak{N}_\Omega = \bigcup_{p \in \Omega} \mathfrak{N}_p$.

We have the following

Theorem 3.1. *Let Ω be a nonempty subset of the set of primes such that $\bigcap_{p \in \Omega} J_p(R) = 0$ and a group G is discriminated by the class of groups $\mathfrak{N}_\Omega$. If for every proper subset Λ of the set Ω at least one of the conditions*

- 1) $\bigcap_{p \in \Lambda} J_p(R) = 0$;
- 2) G is discriminated by the class of groups $\mathfrak{N}_{\Omega \setminus \Lambda}$;

holds, then $A^\omega(RG) = 0$.

PROOF. Let $x = \sum_{i=1}^n \alpha_i g_i \in A^\omega(RG)$ and let Λ be the set of those primes of Ω for which there exists a homomorphism ϕ_p of G into the nilpotent p -group H_p of finite exponent with the property $\phi_p(g_i) \neq \phi_p(g_j)$ for all $i \neq j$. Since the group G is discriminated by the class of groups $\mathfrak{N}_\Omega$, Λ is nonempty.

Let p be an arbitrary element of Λ . If $\phi_p^* : RG \rightarrow RH_p$ is the ring homomorphism, induced by the homomorphism ϕ_p , then $\phi_p^*(x) = \sum_{i=1}^n \alpha_i \phi_p(g_i)$ lies in $A^\omega(RH_p)$. Because p is an arbitrary element of the set Λ , by Lemma 3.4 we obtain that

$$\alpha_i \in \bigcap_{p \in \Lambda} J_p(R)$$

for all $i = 1, 2, \dots, n$. If $\bigcap_{p \in \Lambda} J_p(R) \neq 0$ then from the conditions of this theorem it follows that G is discriminated by the class of groups $\mathfrak{N}_{\Omega \setminus \Lambda}$. Then there exists an element $q \in \Omega \setminus \Lambda$ and a homomorphism ϕ_q of the group G into the nilpotent q -group of finite exponent H_q for which $\phi_q(g_i) \neq \phi_q(g_j)$ for all $i \neq j$, $i, j = 1, 2, \dots, n$. Then by construction of the set Λ it follows that $q \in \Lambda$. This is a contradiction. Consequently, $\bigcap_{p \in \Lambda} J_p(R) = 0$ and $\alpha_i = 0$ for $i = 1, 2, \dots, n$. Therefore $x = 0$ and $A^\omega(RG) = 0$. $\square$

Lemma 3.5. *Let $\mathfrak{G}$ be a class of groups and $\{G_\alpha\}$ ($\alpha \in J$) a family of the normal subgroups of G such that the conditions*

- 1) $G/G_\alpha \in \mathfrak{G}$ for each $\alpha \in J$;
- 2) G_α is torsion-free for all $\alpha \in J$;

hold. If G is not discriminated by a class of groups $\mathfrak{G}$ then there exists a finite subset of distinct elements $g_1, g_2, \dots, g_s$ from G such that the nonzero element $y = (g_1 - 1)(g_2 - 1) \cdots (g_s - 1)$ lies in the ideal $\bigcap_{\alpha \in J} I(G_\alpha)$.

PROOF. If the group G is not discriminated by the class of groups $\mathfrak{G}$ then there exists a finite subset of distinct elements $h_1, h_2, \dots, h_m$ from G

such that for every $\alpha \in J$, $h_i G_\alpha = h_j G_\alpha$ for some $i \neq j$. Then $h_i h_j^{-1} \in G_\alpha$ and the element $h_i h_j^{-1}$ has infinite order.

Let $M = \{g_1, g_2, \dots, g_s\}$ be the set of all those elements of the form $h_i h_j^{-1}$ which have infinite order ($1 \leq i, j \leq m$). It is well known that if g is an element of infinite order then from the equation $(g-1)x = 0$ it follows that $x = 0$ (see for example [7], III. Proposition 4.18). Since g_i ($i=1, 2, \dots, s$) have infinite order, the product $y = (g_1-1)(g_2-1) \cdots (g_s-1)$ is nonzero. From the construction of the set M it follows that for every $\alpha \in J$ there exists at least one element g_i of M such that $g_i \in G_\alpha$. Consequently, $y \in \bigcap_{\alpha \in J} I(G_\alpha)$. $\square$

Theorem 3.2. *Let a group G contain a generalized torsion element. Then $A(RG)$ is residually nilpotent if and only if there exists a nonempty subset Ω of the set of primes such that $\bigcap_{p \in \Omega} J_p(R) = 0$, the group G is discriminated by the class of groups $\mathfrak{N}_\Omega$, and for every proper subset Λ of the set Ω at least one of the conditions*

- 1) $\bigcap_{p \in \Lambda} J_p(R) = 0$;
- 2) G is discriminated by the class of groups $\mathfrak{N}_{\Omega \setminus \Lambda}$;

holds.

PROOF. Let $A^\omega(RG) = 0$ and let the group G contain a torsion element. Then in G there exists a p -element g . Let p^n be the order of g . We show that in this case the set $\Omega = \{p\}$ satisfies the conditions of this theorem.

Then element g^{p^n} belongs to $\gamma_t(G)$ and $\gamma_t(G) \subseteq D_t(RG)$ for every t . Therefore, by Lemma 3.1,

$$(8) \quad p^m(g-1) \in A^t(RG)$$

for some m depending only on t . If $r \in J_p(R)$ then for every m the element r can be decomposed as $r = p^m r_m$, ($r_m \in R$). Therefore from (8) we obtain that $r(g-1) \in A^t(RG)$ for every t . Hence $r(g-1) \in A^\omega(RG)$ and $r(g-1) = 0$. This equation is possible only if $r = 0$. Consequently, $J_p(R) = 0$.

Now we show that the group G is discriminated by the class of groups $\mathfrak{N}_p$. Let $1 \neq h \in \bigcap_{n=1}^{\infty} G^{p^n} \gamma_n(G)$. Hence for every t and m there exists i such that $h \in G^{p^i} \gamma_i(G)$ and $i \geq t + m$. Then, by Lemma 3.2,

$$(9) \quad h - 1 \equiv p^m F_t(h) \pmod{A^t(RG)}$$

follows, where $F_t(h) \in A(RG)$. Therefore from (8) we have that $(g-1) \cdot (h-1) \in A^t(RG)$ for all t . Then $(g-1)(h-1) \in A^\omega(RG)$ and $(g-1)(h-1) = 0$. This is possible only if $g = h$, $g^2 = 1$, $p = 2$ (because g is a p -element of G) and the characteristic of the ring R equals 2. Then (9) implies that $h-1 \in A^t(RG)$ for all t . Therefore $h-1 \in A^\omega(RG)$ and $h = 1$ which is a contradiction. Consequently, $\bigcap_{n=1}^{\infty} G^{p^n} \gamma_n(G) = 1$. From this equation it follows that G is a residually- $\mathfrak{N}_p$ group. Since the class of the groups $\mathfrak{N}_p$ is closed for taking subgroups and forming finite direct products, by Lemma 2.4 we have that G is discriminated by the class of groups $\mathfrak{N}_p$. Consequently, we can choose the set Ω such that $\Omega = \{p\}$.

Now let G be a torsion-free group with the generalized torsion element g of infinite order. Because $A^\omega(RG) = 0$, from Lemma 2.1 $g \notin \bigcap_{n=1}^{\infty} \gamma_n(G)$ follows. Therefore Ω is nonempty.

Now we show that the set Ω can be chosen as $\Omega = \Omega_g$. By Lemma 3.3 (case 2) for every $r \in \bigcap_{p \in \Omega} J_p(R)$ the element $r(g-1)$ lies in $A^\omega(RG)$. Then $r(g-1) = 0$. Therefore $r = 0$ and $\bigcap_{p \in \Omega} J_p(R) = 0$.

Let Λ be a subset of Ω . If G is not discriminated by $\mathfrak{N}_{\Omega \setminus \Lambda}$ then, according to Lemma 3.5 (here we suppose that the family $\{G_\alpha\}$ of the normal subgroups of G coincides with $\{G^{p^n} \gamma_n(G), p \in \Omega \setminus \Lambda, n = 2, 3, \dots\}$, the class $\mathfrak{G}$ is $\mathfrak{N}_{\Omega \setminus \Lambda}$) there exists a nonzero element x in the ideal

$$\bigcap_{p \in \Omega_g \setminus \Lambda} \bigcap_{n=1}^{\infty} I(G^{p^n} \gamma_n(G)) \text{ of the form } x = (g_1 - 1)(g_2 - 1) \cdots (g_s - 1).$$

If Λ is empty then, by case 3 of Lemma 3.3, we have that $(g-1)x \in A^\omega(RG)$. Therefore $(g-1)x = 0$. In the group ring of torsion-free groups such an equation is impossible. Hence G is discriminated by the class of groups $\mathfrak{N}_\Omega$.

Now let Λ be a proper subset of the set Ω and r be an arbitrary element from $\bigcap_{p \in \Lambda} J_p(R)$. If G is not discriminated by $\mathfrak{N}_{\Omega \setminus \Lambda}$ then, by Lemma 3.3 (case 1), $r(g-1)x \in A^\omega(RG)$. Therefore $r(g-1)x = 0$. This equation is possible only if $r = 0$. Hence if G is not discriminated by the class of groups $\mathfrak{N}_{\Omega \setminus \Lambda}$ then $\bigcap_{p \in \Lambda} J_p(R) = 0$.

Sufficiency is proved in Theorem 3.1. $\square$

If the additive group of a ring R is torsion-free and if a group G has no generalized torsion elements then the question about the residual nilpotency of the augmentation ideal is solved (see in particular [2] Theorem 15.5).

The torsion subgroup $T(R^+)$ of the additive group R^+ of a ring R is the direct sum of its primary components $S_p(R^+)$ which are the ideals of the ring R .

Let Π be the set of those primes for which the p -primary components $S_p(R^+)$ of $T(R^+)$ are nonzero.

Lemma 3.6. *Let $A^\omega(RG) = 0$ and $T(R^+) \neq 0$. Then G is a residually- $\mathfrak{N}_p$ group for all $p \in \Pi$.*

PROOF. Let $p \in \Pi$ and $0 \neq r \in S_p(R^+)$. Then $p^s r = 0$ for some s . If $g \in \bigcap_{n=1}^{\infty} G^{p^n} \gamma_n(G)$ then, by Lemma 3.2, we obtain that for all t $g - 1 \equiv p^s F_t(g) \pmod{A^t(RG)}$ ($F_t(g) \in A(RG)$, $n \geq t + s$, $s \geq 1$). Hence $r(g - 1) \equiv p^s r F_t(g) \equiv 0 \pmod{A^t(RG)}$ for all t and $r(g - 1) \in A^\omega(RG)$. Consequently, $g = 1$ and $\bigcap_{n=1}^{\infty} G^{p^n} \gamma_n(G) = 1$ i.e. G is a residually- $\mathfrak{N}_p$ group. $\square$

The element g of the additive Abelian group G is called an element of *infinite p -height* if the equation $p^n x = g$ has a solution in G for all natural numbers n .

Theorem 3.3. *Let the torsion group $T(R^+)$ of the additive group R^+ of a ring R be nonzero, and suppose that for some $p \in \Pi$ the group $T(R^+)$ has no elements of infinite p -height. Further let G be a group with no generalized torsion elements. Then $A^\omega(RG) = 0$ if and only if G is a residually- $\mathfrak{N}_p$ group for all $p \in \Pi$.*

PROOF. Let G be a residually- $\mathfrak{N}_p$ group for all $p \in \Pi$ and let $R_p = R/J_p(R)$. Then $J_p(R_p) = 0$ and, by Theorem 2.1, we obtain that $A^\omega(R_p H) = 0$ for all $H \in \mathfrak{N}_p$ and every prime $p \in \Pi$. Therefore, by Lemmas 2.3 and 2.4, we have that $A^\omega(R_p G) = 0$. If the element $x = \sum_{i=1}^n \alpha_i g_i$ lies in the ideal $A^\omega(RG)$ then from the last equation we conclude that

$$(10) \quad \alpha_i \in J_p(R)$$

for all $p \in \Pi$ and every $i = 1, 2, \dots, n$.

Let $\sqrt{\gamma_n(G)} = \{g \in G \mid g^m \in \gamma_n(G) \text{ for some integer } m \geq 1\}$ be the isolator of the subgroup $\gamma_n(G)$ in G . Obviously, the $G/\sqrt{\gamma_n(G)}$ are torsion-free nilpotent groups and $\bigcap_{n=1}^{\infty} \sqrt{\gamma_n(G)} = 1$ because the group G has no generalized torsion elements. Consequently, $G \in \mathbf{RN}_0$ i.e. G is a residually torsion-free nilpotent group. Since the additive group of the ring $R/T(R^+)$ is without torsion, by Theorem 2.2. $A^\omega(R/T(R^+)G) = 0$

follows. From this we infer that $\alpha_i \in T(R^+)$ and by (10) we obtain that $\alpha_i \in \bigcap_{p \in \Pi} (J_p(R) \cap T(R^+))$ for $i = 1, 2, \dots, n$. It is known that $T(R^+)$ is the serving subgroup of R^+ . Therefore

$$\begin{aligned} J_p(R) \cap T(R^+) &= \left(\bigcap_{n=1}^{\infty} p^n R \right) \cap T(R^+) = \bigcap_{n=1}^{\infty} ((p^n R) \cap T(R^+)) = \\ &= \bigcap_{n=1}^{\infty} p^n T(R^+) = J_p(T(R^+)). \end{aligned}$$

Hence $\alpha_i \in \bigcap_{p \in \Pi} J_p(T(R^+))$ for $i = 1, 2, \dots, n$. Because for some $p = p_0 \in \Pi$, the torsion group $T(R^+)$ has no element of infinite p_0 -height, $J_{p_0}(T(R^+)) = 0$. Therefore $\bigcap_{p \in \Pi} J_p(T(R^+)) = 0$ and $\alpha_i = 0$ for all $i = 1, 2, \dots, n$. Consequently, $x = 0$ and $A^\omega(RG) = 0$.

The necessity is proved in Lemma 3.5. $\square$

Remark 1. It is obvious that $T(R^+) \subseteq pT(R^+)$ for all $p \notin \Pi$ and consequently the question about the residual nilpotency of the augmentation ideals remains open in the case when the group G has no generalized torsion elements and the torsion group of the additive group of the ring R has an element of infinite p -height for all primes p .

Remark 2. Theorems 3.1, 3.2 and 3.3 were announced in [4].

References

- [1] A. A. BOVDI, On the intersection of powers of the augmentation ideal, *Math. Notes* **2,2** (1967), 129–132, (In Russian).
- [2] A. A. BOVDI, Group rings, *UMK VO, Kiev*, 1988, (In Russian).
- [3] B. HARTLEY, The residual nilpotence of wreath products, *Proc. London Math. Soc.* **20,3** (1970), 365–392.
- [4] B. L. KIRÁLY, The residual nilpotency of the augmentation ideal of group rings, Preprint, *UkrNIINTI, No 1498-Uk87, Uzhorod* (1987), 1–22, (In Russian).
- [5] A. I. LICHTMAN, The residual nilpotence of the augmentation ideal and the residual nilpotence of some classes of groups, *Israel J. Math.* **26** (1974), 276–293.
- [6] I. B. PASSI, Group rings and their augmentation ideals, *Lecture Notes in Mathematics*, vol. 715, *Springer-Verlag, Berlin-Heidelberg-New York*, 1979.
- [7] S. K. SEHGAL, Topics in group rings, *Marcell-Dekker, New York-Basel*, 1978.

BERTALAN KIRÁLY
 ESZTERHÁZY KÁROLY TEACHERS' TRAINING COLLEGE
 DEPARTMENT OF MATHEMATICS
 LEÁNYKA U. 4.
 3301 EGER, PF. 43.
 HUNGARY

(Received May 3, 1993)