

Some convergence theorems for the q -integral

By MINGJIN WANG (Changzhou)

Abstract. First, we use Ramanujan's ${}_1\psi_1$ summation formula to obtain an inequality for the bilateral basic hypergeometric series ${}_r\psi_r$. Then, we give some convergence theorems for the q -integral.

1. Introduction

Convergence is an important problem in the study of q -series. There are some papers in the literature [7], [8], [9], [10]. For example, ITO used inequality technique to give a sufficient condition for convergence of a special q -series called Jackson integral [7]. In this paper, we gave some convergence theorems for the q -integral. We first recall some definitions, notation and known results in [1] which will be used in this paper. Throughout the whole paper, it is supposed that $0 < q < 1$. The q -shifted factorials are defined as

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k), \quad (a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k). \quad (1.1)$$

We also adopt the following compact notation for multiple q -shifted factorials:

$$(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \dots (a_m; q)_n, \quad (1.2)$$

where n is an integer or ∞ . We may extend the definition (1.1) of $(a; q)_n$ to

$$(a; q)_\alpha = \frac{(a; q)_\infty}{(aq^\alpha; q)_\infty}, \quad (1.3)$$

Mathematics Subject Classification: Primary: 26D15; Secondary: 33D15.

Key words and phrases: inequality, the bilateral basic hypergeometric function ${}_r\psi_r$, Ramanujan's ${}_1\psi_1$ summation formula, convergence, q -integral.

Supported by the National Natural Science Foundation (grant 11271057) of China.

for any complex number α . In particular,

$$(a; q)_{-n} = \frac{(a; q)_{\infty}}{(aq^{-n}; q)_{\infty}} = \frac{1}{(aq^{-n}; q)_n} = \frac{(-q/a)^n}{(q/a; q)_n} q^{\binom{n}{2}}. \quad (1.4)$$

The bilateral basic hypergeometric series ${}_r\psi_s$ is defined by

$$\begin{aligned} {}_r\psi_s \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix}; q, z \right) \\ = \sum_{n=-\infty}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_s; q)_n} (-1)^{(s-r)n} q^{\binom{s-r}{2}n} z^n. \end{aligned} \quad (1.5)$$

The following is the well known Ramanujan's ${}_1\psi_1$ summation formula

$$\sum_{n=-\infty}^{\infty} \frac{(a; q)_n}{(b; q)_n} z^n = \frac{(q, b/a, az, q/az; q)_{\infty}}{(b, q/a, z, b/az; q)_{\infty}}, \quad |b/a| < |z| < 1. \quad (1.6)$$

2. An inequality for the bilateral basic hypergeometric series

Inequality technique is one of the useful tools in the study of special functions [2], [3], [4], [8], [9], [10]. In this section, we use Ramanujan's ${}_1\psi_1$ summation formula to derive an inequality for the bilateral basic hypergeometric series ${}_{r+1}\psi_{r+1}$, which can be used to discuss the convergence of the q -series. The main result of this section is the following inequality.

Theorem 2.1. *Let a, b be any real numbers such that $q < b < a < 1$ or $a < b < 0$, and let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for any $b/a < |z| < 1$, we have*

$$\left| {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, z \right) \right| \leq M \frac{(q, b/a, a|z|, q/a|z|; q)_{\infty}}{(b, q/a, |z|, b/a|z|; q)_{\infty}}, \quad (2.1)$$

where

$$M = \max \left\{ \prod_{i=1}^r \frac{(-|a_i|; q)_{\infty}}{(|b_i|; q)_{\infty}}, \prod_{i=1}^r \frac{(-q/|b_i|; q)_{\infty}}{(q/|a_i|; q)_{\infty}} \right\}.$$

PROOF. Let $n \geq 0$. Since,

$$|(a_j; q)_n| = \prod_{i=0}^{n-1} |1 - a_j q^i| \leq \prod_{i=0}^{n-1} (1 + |a_j| q^i) = (-|a_j|; q)_n \leq (-|a_j|; q)_{\infty}$$

and

$$|(b_j; q)_n| = \prod_{i=0}^{n-1} |1 - b_j q^i| \geq \prod_{i=0}^{n-1} (1 - |b_j| q^i) = (|b_j|; q)_n \geq (|b_j|; q)_\infty > 0,$$

we have

$$\left| \frac{(a_j; q)_n}{(b_j; q)_n} \right| \leq \frac{(-|a_j|; q)_\infty}{(|b_j|; q)_\infty}. \quad (2.2)$$

Hence,

$$\left| \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_r; q)_n} \right| \leq \prod_{i=1}^r \frac{(-|a_i|; q)_\infty}{(|b_i|; q)_\infty}. \quad (2.3)$$

On the other hand, using (1.4) and (2.2) gives

$$\left| \frac{(a_i; q)_{-n}}{(b_i; q)_{-n}} \right| = \left| \left(\frac{b_i}{a_i} \right)^n \frac{(q/b_i; q)_n}{(q/a_i; q)_n} \right| \leq \left| \frac{b_i}{a_i} \right|^n \frac{(-q/|b_i|; q)_\infty}{(q/|a_i|; q)_\infty}. \quad (2.4)$$

Consequently,

$$\begin{aligned} \left| \frac{(a_1, a_2, \dots, a_r; q)_{-n}}{(b_1, b_2, \dots, b_r; q)_{-n}} \right| &\leq \left| \frac{b_1 b_2 \dots b_r}{a_1 a_2 \dots a_r} \right|^n \prod_{i=1}^r \frac{(-q/|b_i|; q)_\infty}{(q/|a_i|; q)_\infty} \\ &\leq \prod_{i=1}^r \frac{(-q/|b_i|; q)_\infty}{(q/|a_i|; q)_\infty}. \end{aligned} \quad (2.5)$$

Combining (2.3) and (2.5) gives

$$\left| \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_r; q)_n} \right| \leq M = \max \left\{ \prod_{i=1}^r \frac{(-|a_i|; q)_\infty}{(|b_i|; q)_\infty}, \prod_{i=1}^r \frac{(-q/|b_i|; q)_\infty}{(q/|a_i|; q)_\infty} \right\}, \quad (2.6)$$

where $n = \dots - 2, -1, 0, 1, 2, \dots$

Under the condition $q < b < a < 1$ or $a < b < 0$, it is easy to know

$$\frac{(a; q)_n}{(b; q)_n} > 0, \quad n = \dots - 2, -1, 0, 1, 2, \dots, \quad (2.7)$$

and

$$0 < b/a < 1. \quad (2.8)$$

Using (2.6), (2.7), (2.8), and Ramanujan's ${}_1\psi_1$ summation formula (1.6), we have

$$\left| {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, z \right) \right| = \left| \sum_{n=-\infty}^{\infty} \frac{(a, a_1, \dots, a_r; q)_n}{(b, b_1, \dots, b_r; q)_n} z^n \right|$$

$$\begin{aligned}
&\leq \sum_{n=-\infty}^{\infty} \left| \frac{(a, a_1, \dots, a_r; q)_n}{(b, b_1, \dots, b_r; q)_n} z^n \right| = \sum_{n=-\infty}^{\infty} \left\{ \frac{(a; q)_n |z|^n}{(b; q)_n} \cdot \left| \frac{(a_1, \dots, a_r; q)_n}{(b_1, \dots, b_r; q)_n} \right| \right\} \\
&\leq M \sum_{n=-\infty}^{\infty} \frac{(a; q)_n}{(b; q)_n} |z|^n = M \frac{(q, b/a, a|z|, q/a|z|; q)_{\infty}}{(b, q/a, |z|, b/a|z|; q)_{\infty}}, \quad (2.9)
\end{aligned}$$

where $b/a < |z| < 1$. Therefor inequality (2.1) holds. Thus, we complete the proof. $\square$

3. Some convergence theorems for the q -integral

F. H. JACKSON defined the q -integral by [6]

$$\int_0^d f(t) d_q t = d(1-q) \sum_{n=0}^{\infty} f(dq^n) q^n, \quad (3.1)$$

and

$$\int_c^d f(t) d_q t = \int_0^d f(t) d_q t - \int_0^c f(t) d_q t. \quad (3.2)$$

In this section, we use the inequality (2.1) to give some sufficient conditions for convergence of the q -integral. First, we give the following lemma:

Lemma 3.1. *Let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$, let $\{u_n\}$ and $\{v_n\}$ be any convergence real number series such that $q < v_n < u_n < 1$ or $u_n < v_n < 0$ and let $\{c_n\}$ and $\{d_n\}$ be any convergence real number series such that $v_n/u_n < |d_n| < 1$. If*

$$\lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = p < 1,$$

then the q -series

$$\sum_{n=1}^{\infty} c_n {}_{r+1}\psi_{r+1} \left(\begin{matrix} u_n, a_1, \dots, a_r \\ v_n, b_1, \dots, b_r \end{matrix}; q, d_n \right) \quad (3.3)$$

converges absolutely.

PROOF. Letting $a = u_n, b = v_n$ and $z = d_n$ in (2.1) gives

$$\left| {}_{r+1}\psi_{r+1} \left(\begin{matrix} u_n, a_1, \dots, a_r \\ v_n, b_1, \dots, b_r \end{matrix}; q, d_n \right) \right| \leq M \frac{(q, v_n/u_n, u_n|d_n|, q/u_n|d_n|; q)_{\infty}}{(v_n, q/u_n, |d_n|, v_n/u_n|d_n|; q)_{\infty}}. \quad (3.4)$$

Multiplying both sides of (3.4) by $|c_n|$ gets

$$\left| c_n {}_{r+1}\psi_{r+1} \left(\begin{matrix} u_n, a_1, \dots, a_r \\ v_n, b_1, \dots, b_r \end{matrix}; q, d_n \right) \right| \leq |c_n| M \frac{(q, v_n/u_n, u_n|d_n|, q/u_n|d_n|; q)_\infty}{(v_n, q/u_n, |d_n|, v_n/u_n|d_n|; q)_\infty}. \quad (3.5)$$

Let

$$e_n = |c_n| M \frac{(q, v_n/u_n, u_n|d_n|, q/u_n|d_n|; q)_\infty}{(v_n, q/u_n, |d_n|, v_n/u_n|d_n|; q)_\infty}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{e_{n+1}}{e_n} = \lim_{n \rightarrow \infty} \left| \frac{c_{n+1}}{c_n} \right| = p < 1,$$

the ratio test shows that the series

$$\sum_{n=0}^{\infty} e_n$$

is convergent. From (3.5), it is sufficient to establish that the q -series (3.3) is absolutely convergent. $\square$

Using Lemma 3.1, we can easily get some convergence theorems for the q -integral.

Theorem 3.2. *Let a, b be any real numbers such that $q < b < a < 1$ or $a < b < 0$, let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for any $0 < d < 1$ and $\alpha > -1$, the q -integral*

$$\int_0^d t^\alpha {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, z \right) d_q t \quad (3.6)$$

converges absolutely. Where $z = b/a + (1 - b/a)t$.

PROOF. It is easy to see that

$$b/a < b/a + (1 - b/a)dq^n < 1, \quad n = 0, 1, \dots$$

By the definition of q -integral (3.1), we get

$$\begin{aligned} & \int_0^d t^\alpha {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, z \right) d_q t \\ &= d^{1+\alpha} (1 - q) \sum_{n=0}^{\infty} q^{n(1+\alpha)} {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, b/a + (1 - b/a)dq^n \right). \end{aligned} \quad (3.7)$$

Using Lemma 3.1 with $u_n = a$, $v_n = b$, $c_n = q^{n(1+\alpha)}$, $d_n = b/a + (1 - b/a)dq^n$, and noticing

$$\lim_{n \rightarrow \infty} \frac{q^{(n+1)(1+\alpha)}}{q^{n(1+\alpha)}} = q^{(1+\alpha)} < 1,$$

we know the q -integral (3.6) converges absolutely. $\square$

By the definition of q -integral (3.2), we immediately get

Corollary 3.3. *Let a, b be any real numbers such that $q < b < a < 1$ or $a < b < 0$, let a_i, b_i be any real numbers such that $|a_i| > q$, $|b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for any $0 < c, d < 1$ and $\alpha > -1$, the q -integral*

$$\int_c^d t^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ b, b_1, \dots, b_r \end{matrix}; q, z \right) d_q t \quad (3.8)$$

converges absolutely. Where $z = b/a + (1 - b/a)t$.

Theorem 3.4. *Let a, b and c be any real numbers such that $a < 0, b < 0$, $b/a < c < 1$, let a_i, b_i be any real numbers such that $|a_i| > q$, $|b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for any $0 < d < 1$ and $\alpha > -1$, the q -integral*

$$\int_0^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \quad (3.9)$$

converges absolutely.

PROOF. By the definition of q -integral (3.1), we get

$$\begin{aligned} & \int_0^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \\ &= d^{1+\alpha} (1-q) \sum_{n=0}^{\infty} q^{n(1+\alpha)} {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ bdq^n, b_1, \dots, b_r \end{matrix}; q, cdq^n \right). \end{aligned} \quad (3.10)$$

Using Lemma 3.1 with $u_n = a$, $v_n = bdq^n$, $c_n = q^n$ and $d_n = cdq^n$ and noticing

$$a < bdq^n < 0, \quad bdq^n/a < cdq^n < 1,$$

we get the q -integral (3.9) converges absolutely. $\square$

By the definition of q -integral (3.2), we immediately get

Corollary 3.5. Let a, b and c be any real numbers such that $a < 0, b < 0, b/a < c < 1$, let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for any $0 < d, e < 1$ and $\alpha > -1$, the q -integral

$$\int_e^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} a, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \quad (3.11)$$

converges absolutely.

Theorem 3.6. Let a, b and c be any real numbers such that $a < 0, b < 0, b/a < c < 1$, let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for $\alpha > -1, \beta \leq 0$ and $1 \leq d < 1/c$, the q -integral

$$\int_0^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} az^\beta, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \quad (3.12)$$

converges absolutely.

PROOF. By the definition of q -integral (3.1), we get

$$\begin{aligned} & \int_0^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} az^\beta, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \\ &= d^{1+\alpha} (1-q) \sum_{n=0}^{\infty} q^{n(1+\alpha)} {}_{r+1}\psi_{r+1} \left(\begin{matrix} adq^{n\beta}, a_1, \dots, a_r \\ bdq^n, b_1, \dots, b_r \end{matrix}; q, cdq^n \right). \end{aligned} \quad (3.13)$$

Using Lemma 3.1 with $u_n = adq^{n\beta}, v_n = bdq^n, c_n = q^{n(1+\alpha)}$ and $d_n = cdq^n$ and noticing

$$aq^{n\beta} < bq^n < 0, \quad bq^{n(1-\beta)}/a < cdq^n < 1,$$

we have the q -integral (3.12) converges absolutely. $\square$

Corollary 3.7. Let a, b and c be any real numbers such that $a < 0, b < 0, b/a < c < 1$, let a_i, b_i be any real numbers such that $|a_i| > q, |b_i| < 1$ for $i = 1, 2, \dots, r$ with $r \geq 1$ and $|b_1 b_2 \dots b_r| \leq |a_1 a_2 \dots a_r|$. Then for $\alpha > -1, \beta \leq 0$ and $1 \leq d, e < 1/c$, the q -integral

$$\int_e^d z^\alpha \cdot {}_{r+1}\psi_{r+1} \left(\begin{matrix} az^\beta, a_1, \dots, a_r \\ bz, b_1, \dots, b_r \end{matrix}; q, cz \right) d_q z \quad (3.14)$$

converges absolutely.

References

- [1] G. GASPER and M. RAHMAN, Basic Hypergeometric Series, *Cambridge Univ. Press, Cambridge, MA*, 1990.
- [2] C. GIORDANO, A. LAFORGIA and JOSIP PEČARIĆ, Supplements to known inequalities for some special functions, *J. Math. Anal. Appl.* **200** (1996), 34–41.
- [3] ÁRPÁD ELBERT and A. LAFORGIA, An inequality for the product of two integrals relating to the incomplete gamma function, *J. Inequal. Appl.* **5**, no. 1 (2000), 39–51, doi:10.1155/S1025583400000035.
- [4] M. E. H. ISMAIL and A. LAFORGIA, Functional inequalities for incomplete gamma and related functions, *Math. Inequal. Appl.* **9** (2006), 233–247.
- [5] A. LAFORGIA and P. NATALINI, On some Turn-type inequalities, *J. Inequal. Appl.* **2006**, Article ID 29828 (2006), 6, doi:10.1155/JIA/2006/29828.
- [6] F. H. JACKSON, On q -definite integrals, *Quart. J.* **41** (1910), 193–203.
- [7] M. ITO, Convergence and asymptotic behavior of Jackson integrals associated with irreducible reduced root systems, *J. Approximation Theory* **124** (2003), 154–180.
- [8] M. WANG, Two inequalities for ${}_r\phi_r$ and applications, *J. Inequal. Appl.* **2008**, Article ID 471527 (2008), 6, doi:10.1155/2008/471527.
- [9] M. WANG and X. ZHAO, On convergence of q -series involving ${}_{r+1}\phi_r$ basic hypergeometric series, *J. Inequal. Appl.* **2009**, Article ID 170526 (2009), 9, doi:10.1155/2009/170526.
- [10] M. WANG, Inequalities for terminating ${}_{r+1}\phi_r$ and applications, *J. Math. Inequal.* **4** (2010), 121–128.

MINGJIN WANG
 DEPARTMENT OF APPLIED MATHEMATICS
 CHANGZHOU UNIVERSITY
 CHANGZHOU, JIANGSU 213164
 P.R. CHINA
E-mail: wang197913@126.com

(Received September, 13, 2011)