

Title: On a conjecture about repdigits in k -generalized Fibonacci sequences

Author(s): Jhon J. Bravo and Florian Luca

The k -generalized Fibonacci sequence $(F_n^{(k)})_n$ resembles the Fibonacci sequence in that it starts with $0, \dots, 0, 1$ (a total of k terms) and each term afterwards is the sum of the k preceding terms. F. LUCA [4] in 2000 and recently D. MARQUES [5] proved that 55 and 44 are the largest numbers with only one distinct digit (so called *repdigits*) in the sequences $(F_n^{(2)})_n$ and $(F_n^{(3)})_n$, respectively. Further, Marques conjectured that there are no repdigits having at least 2 digits in a k -generalized Fibonacci sequence for any $k > 3$. In the present paper, we confirm this conjecture.

Address:

Jhon J. Bravo
Departamento de Matemáticas
Universidad del Cauca
Calle 5 No. 4-70
Popayán
Colombia
Current address:
Instituto de Matemáticas
Universidad Nacional Autónoma de México
Circuito Exterior
Ciudad Universitaria, C. P. 04510
México D.F.
Mexico

Address:

Florian Luca
Fundación Marcos Moshinsky
Instituto de Ciencias Nucleares UNAM
Circuito Exterior, C.U., Apdo. Postal 70-543
Mexico D.F. 04510
Mexico