

## A condition for a Finsler space to be a Riemannian space

By SHÔJI WATANABE (Tokyo)

### 0. Introduction

In a Finsler space, there are known canonical Finsler connections, that is, the Cartan connection  $CT$ , the Berwald connection  $B\Gamma$ , the Rund connection  $R\Gamma$  and the Hashiguchi connection  $H\Gamma$ .

The concept of the generalized Berwald  $P^1$ -connections was introduced by AIKOU and HASHIGUCHI [1] and MATSUMOTO [2]. In the previous paper [4], we considered a generalized Berwald  $P^1$ -connection  $\Gamma$ . Using this Finsler connection, we obtained a condition for a Finsler space to be a Riemannian space in each of the above-mentioned Finsler connections  $CT$ ,  $B\Gamma$ ,  $R\Gamma$ ,  $H\Gamma$  and  $\Gamma$ , respectively [4].

The purpose of the present paper is to improve these conditions for a Finsler space to be a Riemannian space and to make up these conditions into a condition valid for any of the Finsler connections mentioned above.

Throughout the present paper, the terminology and notations are those of MATSUMOTO's monograph [3].

### 1. Preliminaries

Let  $F^n = (M, L)$  be a Finsler space with a Finsler metric  $L(x, y)$ , where  $x$  denotes a point of the underlying manifold  $M$  and  $y$  denotes a supporting element. The fundamental tensor  $g_{ij}$  is given by  $g_{ij} = \frac{1}{2} \partial^2 L^2 / \partial y^i \partial y^j$ . We shall express a Finsler connection  $F\Gamma(F_h^i{}_j, N_j^i, C_h^i{}_j)$  in terms of its coefficients. The  $(v)hv$ -torsion tensor  $P_{jk}^i$  and the  $hv$ -curvature tensor  $P_h^i{}_{jk}$  of a Finsler connection  $F\Gamma$  are given by

$$(1.1) \quad P_{jk}^i = \dot{\partial}_k N_j^i - F_k^i{}_j,$$

and

$$(1.2) \quad P_h^i{}_{jk} = \dot{\partial}_k F_h^i{}_{\cdot j} - C_h^i{}_{k|j} + C_h^i{}_{\cdot r} P^r{}_{jk},$$

respectively, where “ $|$ ” denotes the  $h$ -covariant differentiation with respect to  $F\Gamma$  and  $\dot{\partial}_k = \partial/\partial y^k$ .

As well-known, the Berwald connection  $B\Gamma$  is given by  $B\Gamma = (G_h^i{}_{\cdot j}, G^i{}_{\cdot j}, 0)$ , where

$$\begin{aligned} G_h^i{}_{\cdot j} &= \frac{1}{2} \dot{\partial}_h \dot{\partial}_j (\gamma_m^i{}_{\cdot r} y^m y^r), \\ \gamma_m^i{}_{\cdot r} &= \frac{1}{2} g^{ij} (\partial g_{jm} / \partial x^r + \partial g_{jr} / \partial x^m - \partial g_{mr} / \partial x^j), \\ G^i{}_{\cdot j} &= G_h^i{}_{\cdot j} y^h = \frac{1}{2} \dot{\partial}_j (\gamma_m^i{}_{\cdot r} y^m y^r). \end{aligned}$$

Then the known Finsler connections  $B\Gamma$ ,  $H\Gamma$ ,  $R\Gamma$ ,  $C\Gamma$  and our Finsler connections  $\Gamma$  and  $\Gamma'$  are denoted by the following table:

|                    |           |                                           |                   |                     |
|--------------------|-----------|-------------------------------------------|-------------------|---------------------|
| Finsler connection | $F\Gamma$ | $F_h^i{}_{\cdot j}$                       | $N^i{}_{\cdot j}$ | $C_h^i{}_{\cdot j}$ |
| Berwald            | $B\Gamma$ | $G_h^i{}_{\cdot j}$                       | $G^i{}_{\cdot j}$ | 0                   |
| Hashiguchi         | $H\Gamma$ | $G_h^i{}_{\cdot j}$                       | $G^i{}_{\cdot j}$ | $g_h^i{}_{\cdot j}$ |
| Rund               | $R\Gamma$ | $G_h^i{}_{\cdot j} - g_h^i{}_{\cdot j 0}$ | $G^i{}_{\cdot j}$ | 0                   |
| Cartan             | $C\Gamma$ | $G_h^i{}_{\cdot j} - g_h^i{}_{\cdot j 0}$ | $G^i{}_{\cdot j}$ | $G_h^i{}_{\cdot j}$ |
|                    | $\Gamma$  | $G_h^i{}_{\cdot j} + L g_h^i{}_{\cdot j}$ | $G^i{}_{\cdot j}$ | 0                   |
|                    | $\Gamma'$ | $G_h^i{}_{\cdot j} + L g_h^i{}_{\cdot j}$ | $G^i{}_{\cdot j}$ | $g_h^i{}_{\cdot j}$ |

where  $g_h^i{}_{\cdot j} = \frac{1}{2} g^{ir} \dot{\partial}_r g_{hj}$  and  $g_h^i{}_{\cdot j|0} = g_h^i{}_{\cdot j|k} y^k$ .

For the above-mentioned Finsler connections, the following equations are satisfied:

$$(1.3) \quad F_h^i{}_{\cdot j} = G_h^i{}_{\cdot j} - P^i{}_{jh},$$

$$(1.4) \quad P_0^i{}_{jk} := P_h^i{}_{jk} y^h = P^i{}_{jk}.$$

Using (1.3), (1.2) is rewritten as follows:

$$(1.5) \quad P_h^i{}_{jk} = G_h^i{}_{\cdot j \cdot k} - P^i{}_{hj \cdot k} - C_h^i{}_{k|j} + C_h^i{}_{\cdot r} P^r{}_{jk},$$

where  $\cdot k = \dot{\partial}_k$ .

## 2. The main result

In this section, we shall assume that the Finsler connection  $F\Gamma$  is one of the Finsler connections  $B\Gamma$ ,  $H\Gamma$ ,  $R\Gamma$ ,  $C\Gamma$ ,  $\Gamma$  and  $\Gamma'$ .

**Theorem.** *Let  $A_h^i{}_j = Lg_h^i{}_j$ . Then a Finsler space  $F^n$  is Riemannian, if and only if*

$$(2.1) \quad P_h^i{}_{jk} = -A_h^i{}_{j \cdot k}.$$

To prove the theorem, we shall prove the following

**Lemma.** *Let  $F^n$  be a Finsler space with the  $hv$ -curvature tensor  $P_h^i{}_{jk}$  satisfying (2.1). Then  $F^n$  is a Landsberg space satisfying*

$$(2.2) \quad P_j^i{}_{k} = A_j^i{}_{\cdot k}.$$

PROOF of the Lemma. Contracting (2.1) by  $y^h$  and paying attention to (1.4), we have (2.2). Thus, from (1.5), (2.1) and (2.2), we get

$$-A_h^i{}_{j \cdot k} = G_h^i{}_{j \cdot k} - A_h^i{}_{j \cdot k} - C_h^i{}_{k|j} + C_h^i{}_{\cdot r} A_j^r{}_{\cdot k}.$$

Hence we have

$$(2.3) \quad G_h^i{}_{j \cdot k} - C_h^i{}_{k|j} + C_h^i{}_{\cdot r} A_j^r{}_{\cdot k} = 0.$$

*Case 1.* If  $F\Gamma$  is  $H\Gamma$  or  $C\Gamma$  or  $\Gamma'$ , then we have  $C_h^i{}_{\cdot j} = g_h^i{}_{\cdot j}$ . In this case, contracting (2.3) by  $y^j$ , we have  $g_h^i{}_{k|0} = 0$ . This means that  $F^n$  is a Landsberg space.

*Case 2.* If  $F\Gamma$  is  $B\Gamma$  or  $R\Gamma$  or  $\Gamma$ , then we have  $C_h^i{}_{\cdot j} = 0$ . Hence (2.3) reduces to  $G_h^i{}_{j \cdot k} = 0$ . This means that  $F^n$  is a Berwald space and hence a Landsberg space. Thus the lemma is proved.

PROOF of the Theorem. The necessity of (2.1) is evident. Assume (2.1), then by the Lemma, we have that  $F^n$  is a Landsberg space satisfying (2.2). First, we shall consider cases of  $H\Gamma$  and  $B\Gamma$ . For  $H\Gamma$  and  $B\Gamma$ , the  $(v)hv$ -torsion tensor  $P^i{}_{jk}$  vanishes. So from (2.2) we have  $A_j^i{}_{\cdot k} = P^i{}_{jk} = 0$ . Accordingly,  $F^n$  is a Riemannian space. Next, we shall consider cases of  $\Gamma$  and  $\Gamma'$ . For these Finsler connections we have  $P^i{}_{jk} = -A_j^i{}_{\cdot k}$ . So we have  $A_j^i{}_{\cdot k} = P^i{}_{jk} = -A_j^i{}_{\cdot k}$ . Hence  $A_j^i{}_{\cdot k} = 0$ . This means that  $F^n$  is Riemannian. Third, we shall consider cases of  $C\Gamma$  and  $R\Gamma$ . For these Finsler connections, we have  $P^i{}_{jk} = g_j^i{}_{k|0}$ . So, noting that  $F^n$  is a Landsberg space, we obtain  $A_j^i{}_{\cdot k} = P^i{}_{jk} = g_j^i{}_{k|0} = 0$ . This means that  $F^n$  is Riemannian. Thus the proof of the Theorem is complete.

### References

- [1] T. AIKOU and M. HASHIGUCHI, On generalized Berwald connections, *Rep. Fac. Sci. Kagoshima Univ. (Math. Phys. Chem.)* **17** (1984), 9–13.
- [2] M. MATSUMOTO, The induced and intrinsic Finsler connections of a hypersurface and Finsler projective geometry, *J. Math. Kyoto Univ.* **25** (1985), 107–144.
- [3] M. MATSUMOTO, Foundations of Finsler geometry and special Finsler spaces, *Kaiseisha Press, Otsu, Japan*, 1986.
- [4] S. WATANABE, The generalized Berwald  $P^1$ -connection with  $P^i_{jk} = -A_j^i{}_{,k}$  and its applications, *Publ. Math. Debrecen* **42** (1993), 39–44.

SHÔJI WATANABE  
DEPARTMENT OF MATHEMATICS  
SCIENCE UNIVERSITY OF TOKYO  
WAKAMIYA-CHO 26, SHINJUKU-KU  
TOKYO 162, JAPAN

*(Received August 2, 1993)*