

Groups whose proper subgroups are (locally finite)-by-(locally nilpotent)

By AMEL DILMI (Setif) and NADIR TRABELSI (Setif)

Abstract. If $\mathfrak{X}$ is a class of groups, then a group G is called a *minimal non- $\mathfrak{X}$ -group* if it is not an $\mathfrak{X}$ -group but all its proper subgroups belong to $\mathfrak{X}$. Let π be a set of primes and let $\mathfrak{X}$ be a quotient and subgroup closed class of locally nilpotent groups such that every infinite locally graded minimal non- $\mathfrak{X}$ -group is a countable p -group for some prime p . Our main result in the present paper states that G is an infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{X}$ -group if and only if there exists a prime $p \notin \pi$ such that G is an infinitely generated minimal non- $\mathfrak{X}$ p -group; where $L\mathfrak{F}_\pi$ denotes the class of locally finite π -groups.

1. Introduction

If $\mathfrak{X}$ is a class of groups, then a group G is called a *minimal non- $\mathfrak{X}$ -group* if it is not an $\mathfrak{X}$ -group but all its proper subgroups belong to $\mathfrak{X}$. Many results have been obtained on minimal non- $\mathfrak{X}$ -groups for several choices of $\mathfrak{X}$. In particular, in [10] a complete description of infinitely generated minimal non-nilpotent groups having a maximal subgroup is given. These groups are metabelian Chernikov p -groups, where p is a prime. Later in [14], infinitely generated minimal non-nilpotent groups without maximal subgroups have been studied and it was proved, among many results, that they are countable p -groups. In [15] it is proved that if G is a minimal non- $(L\mathfrak{F})\mathfrak{N}$ (respectively, non- $(L\mathfrak{F})\mathfrak{N}_c$) group, then G is a finitely generated perfect group which has no proper subgroups of finite index and $G/\text{Frat}(G)$ is simple, where $L\mathfrak{F}$ (respectively, $\mathfrak{N}$, $\mathfrak{N}_c$) denotes the class of locally finite (respectively, nilpotent, nilpotent of class at most c) groups. Therefore there are

Mathematics Subject Classification: 20F50, 20F19.

Key words and phrases: Locally nilpotent, Locally finite, π -groups, Minimal non-nilpotent.

no minimal non- $(L\mathfrak{F})\mathfrak{N}$ -groups (respectively, non- $(L\mathfrak{F})\mathfrak{N}_c$ -groups) which are infinitely generated (or equivalently locally graded). In the present paper, we generalize these last results by considering the classes $(L\mathfrak{F}_\pi)\mathfrak{N}$ and $(L\mathfrak{F}_\pi)\mathfrak{N}_c$, where π is a given set of primes and $L\mathfrak{F}_\pi$ denotes the class of locally finite π -groups. It turns out that infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{N}$ -groups exist. For if G is an infinitely generated minimal non-nilpotent group, then it is a p -group for some prime p and hence it is an infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{N}$ -group for every set π not containing p . We will prove that the converse is also true. In fact our results on $(L\mathfrak{F}_\pi)\mathfrak{N}$ and $(L\mathfrak{F}_\pi)\mathfrak{N}_c$ will be consequences of more general results on $(L\mathfrak{F}_\pi)\mathfrak{X}$ (respectively, $(L\mathfrak{F}_\pi)\mathfrak{V}$), where $\mathfrak{X}$ (respectively, $\mathfrak{V}$) denotes a quotient and subgroup closed class (respectively, a variety) of locally nilpotent groups such that infinite locally graded minimal non- $\mathfrak{X}$ -groups are countable p -groups. Our main result (Theorem 4.1) states that a group G is an infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{X}$ -group if and only if there exists a prime $p \notin \pi$ such that G is an infinitely generated minimal non- $\mathfrak{X}$ p -group.

Although our main result concerns the class $(L\mathfrak{F}_\pi)\mathfrak{X}$, we will introduce two more general classes, $\Omega_\pi\mathfrak{Z}$ and $\Omega_\pi\mathfrak{V}$, in order to make our preliminary results as general as we can do, for further investigations.

Recall that a group is locally graded if every non-trivial finitely generated subgroup has a proper non-trivial finite homomorphic image.

Our notation and terminology are standard and follow [13]. In the sequel, π denotes a given set of primes. If H is a subgroup of a group G then we denote by $T_\pi(H)$ the subgroup generated by all π -elements of H , that is elements of finite order whose prime divisors belong to π . We will write simply T_π for $T_\pi(G)$. A (not necessary periodic) group G is said to be π -free if it has no π -elements. Finally we denote by $\mathfrak{T}_\pi$ the class of π -groups, that is groups with only π -elements.

2. Minimal non- $\Omega_\pi\mathfrak{Z}$ -groups

Let $\mathfrak{Z}$ be a quotient and subgroup closed class of locally nilpotent groups and let Ω_π be either the class $L\mathfrak{F}_\pi$ or $\mathfrak{T}_\pi$. In this section we will prove a result on infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -groups which will be used many times in this note.

First we prove two useful lemmas.

Lemma 2.1. *Let G be a group generated by π -elements. If G belongs to $L(\Omega_\pi\mathfrak{N})$, then it belongs to Ω_π .*

PROOF. Since $L(\Omega_\pi) = \Omega_\pi$, there is no loss of generality if we assume that G is finitely generated and hence it belongs to $\Omega_\pi\mathfrak{N}$. Let N be a normal subgroup of G such that $N \in \Omega_\pi$ and G/N is nilpotent. We have that $(G/N)/(G/N)'$ is an abelian group generated by finitely many π -elements and hence it is a finite π -group. We deduce by [13, Corollary of Theorem 2.26] that G/N is also a finite π -group as it is nilpotent. Therefore G belongs to Ω_π . $\square$

Lemma 2.2. *Let G be an infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -group. Then T_π belongs to Ω_π and if H is a proper subgroup of G , then so is HT_π .*

PROOF. As G is not finitely generated, $G \in L(\Omega_\pi\mathfrak{Z}) \subseteq L(\Omega_\pi\mathfrak{N})$ and hence $T_\pi \in \Omega_\pi$ by Lemma 2.1. Now let H be a proper subgroup of G . If $G = HT_\pi$, then $G/T_\pi \simeq H/H \cap T_\pi$ is in $\Omega_\pi\mathfrak{Z}$ and hence it belongs to $\mathfrak{Z}$ since it is a π -free group. It follows that $G \in \Omega_\pi\mathfrak{Z}$, which is a contradiction. Therefore HT_π is a proper subgroup of G . $\square$

Proposition 2.3. *An infinitely generated group G is a minimal non- $\Omega_\pi\mathfrak{Z}$ -group if and only if the following conditions are satisfied:*

- (i) T_π belongs to Ω_π , and if H is a proper subgroup of G then so is HT_π ;
- (ii) G/T_π is a π -free infinitely generated minimal non- $\mathfrak{Z}$ -group.

PROOF. Assume that G is an infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -group. Then (i) is satisfied by Lemma 2.2. As $T_\pi \in \Omega_\pi$ by (i), we have that $G/T_\pi \notin \mathfrak{Z}$. Since G is in $L(\Omega_\pi\mathfrak{Z})$ and G/T_π is π -free, we deduce that G/T_π is a minimal non- $\mathfrak{Z}$ -group which is locally in $\mathfrak{Z}$. Hence G/T_π is an infinitely generated π -free-group and (ii) is proved.

Conversely, assume that conditions (i) and (ii) are satisfied. So G/T_π is an infinitely generated π -free group and a minimal non- $\mathfrak{Z}$ -group. Hence $G \notin \Omega_\pi\mathfrak{Z}$ and is infinitely generated. Let H be a proper subgroup of G . Then $H \cap T_\pi \in \Omega_\pi$ and $H/H \cap T_\pi \simeq HT_\pi/T_\pi \neq G/T_\pi$ by (ii). It follows that $H/H \cap T_\pi \in \mathfrak{Z}$ and hence $H \in \Omega_\pi\mathfrak{Z}$. Therefore G is an infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -group. $\square$

Corollary 2.4. *Let G be an infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -group. If G splits over T_π , that is if there exists a subgroup H of G such that $T_\pi \cap H = 1$ and $G = T_\pi H$, then G is a π -free infinitely generated minimal non- $\mathfrak{Z}$ -group.*

PROOF. Assume that G is an infinitely generated minimal non- $\Omega_\pi\mathfrak{Z}$ -group such that $G = T_\pi H$. It follows by condition (i) of Proposition 2.3 that $G = H$. Since $T_\pi \cap H = 1$ we deduce that G is π -free and hence $T_\pi = 1$. It follows by condition (ii) of Proposition 2.3 that G is a π -free infinitely generated minimal non- $\mathfrak{Z}$ -group. $\square$

By Proposition 2.3, if there exists an infinitely generated minimal non- $\Omega_\pi \mathfrak{Z}$ -group, then there exists an infinitely generated minimal non- $\mathfrak{Z}$ -group. Since minimal non- $L\mathfrak{N}$ -groups are obviously finitely generated, $L\mathfrak{N}$ being the class of locally nilpotent groups, we have the following application of Proposition 2.3.

Corollary 2.5. *If G is an infinitely generated group whose proper subgroups are in $\Omega_\pi(L\mathfrak{N})$, then so is G .*

3. Minimal non- $\Omega_\pi \mathfrak{Y}$ -groups

Let $\mathfrak{Y}$ be a quotient and subgroup closed class of locally nilpotent groups such that locally graded minimal non- $\mathfrak{Y}$ -groups are periodic. Our main result in this section is about finitely generated non- $(L\mathfrak{F}_\pi)\mathfrak{Y}$ -groups.

Clearly infinitely generated minimal non- $\Omega_\pi \mathfrak{Y}$ -groups are characterized by Proposition 2.3 which implies that they are periodic. If we take π to be the set of all primes, then $\Omega_\pi = \Omega$ is either the class of locally finite or periodic groups. In this case a non-trivial group cannot be periodic and π -free, so condition (ii) of Proposition 2.3 is not satisfied and hence Proposition 2.3 has the following consequence.

Corollary 3.1. *If G is an infinitely generated group all of whose proper subgroups are in $\Omega \mathfrak{Y}$, then so is G .*

Now we will prove a result on finitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{Y}$ -groups. To this end we need the following lemma which implies together with the remark above that infinite locally graded minimal non- $\Omega_\pi \mathfrak{Y}$ -groups are periodic.

Lemma 3.2. *Let G be a non-periodic group all of whose proper subgroups are in $\Omega_\pi \mathfrak{Y}$. If G is finitely generated and has a proper subgroup of finite index, then it belongs to $\Omega_\pi \mathfrak{Y}$.*

PROOF. Let G as stated and let N be a normal proper subgroup of finite index. Then N is finitely generated and hence $N \in \Omega_\pi \mathfrak{N}$. It follows that if $H := T_\pi(N)$, then $H \in \Omega_\pi$ and $G/H \in \mathfrak{N}\mathfrak{F}$. Therefore G/H satisfies the maximal condition on subgroups and hence every proper subgroup of G/H belongs to $\mathfrak{F}_\pi \mathfrak{N}$. Consequently every proper subgroup of G is in $\Omega_\pi \mathfrak{N}$. We deduce by [15, Corollary 2.2] that G is in $\Omega \mathfrak{N}$ and hence it has a torsion part. Therefore T_π is periodic and hence it is proper in G . So that $T_\pi \in \Omega_\pi$ by Lemma 2.1. Since G/T_π is a π -free group, all its proper subgroups belong to $\mathfrak{Y}$. As G is not periodic, G/T_π is a non-periodic nilpotent-by-finite group and hence it cannot be a minimal non- $\mathfrak{Y}$ -group. Consequently $G/T_\pi \in \mathfrak{Y}$ and so $G \in \Omega_\pi \mathfrak{Y}$, as claimed. $\square$

Note that if a minimal non- $(L\mathfrak{F}_\pi)\mathfrak{Y}$ -group is infinitely generated then it is an infinite locally graded group; the next result implies that the converse is true, that is an infinite locally graded minimal non- $(L\mathfrak{F}_\pi)\mathfrak{Y}$ -group is infinitely generated.

Proposition 3.3. *Let G be an infinite finitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{Y}$ -group. Then G is a perfect group which has no proper subgroups of finite index and $G/\text{Frat}(G)$ is simple.*

PROOF. Let G as stated. Assume that G has a proper normal subgroup of finite index, say N . Then N is finitely generated and hence $N \in (L\mathfrak{F}_\pi)\mathfrak{N}$. Since G is infinite, so is N and hence N is not periodic. It follows by Lemma 3.2 that G is in $(L\mathfrak{F}_\pi)\mathfrak{Y}$, which is a contradiction. So G has no proper subgroups of finite index and hence it is perfect. Since G is finitely generated, $G/\text{Frat}(G)$ is non-trivial. Let $N/\text{Frat}(G)$ be a non-trivial normal subgroup of $G/\text{Frat}(G)$. Therefore there exists a maximal subgroup M of G such that $N \not\leq M$ and hence $G = MN$. Let $F := T_\pi(M)$; since $M \in (L\mathfrak{F}_\pi)\mathfrak{Y}$, $F \in L\mathfrak{F}_\pi$ by Lemma 2.1. If $g \in G$, then it is easy to see that $(NF)^g \leq NF$ and hence NF is normal in G . Moreover

$$G/NF = NM/NF \simeq M/M \cap NF \simeq (M/F)/(M \cap NF/F)$$

which is a finitely generated $\mathfrak{Y}$ -group and hence it is nilpotent. But G is perfect, so we deduce that $G = NF$. It follows that G/N is a finitely generated group in $L\mathfrak{F}_\pi$ and hence it is finite. Therefore G/N is trivial, a contradiction that gives that $G/\text{Frat}(G)$ is simple. $\square$

4. Minimal non- $(L\mathfrak{F}_\pi)\mathfrak{X}$ -groups

Let $\mathfrak{X}$ be a quotient and subgroup closed class of locally nilpotent groups such that infinite locally graded minimal non- $\mathfrak{X}$ -groups are countable p -groups, where p is a prime. Now we are ready to prove our main result, which is an easy consequence of Proposition 2.3.

Theorem 4.1. *A group G is an infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{X}$ -group if and only if there exists a prime $p \notin \pi$ such that G is an infinitely generated minimal non- $\mathfrak{X}$ p -group.*

PROOF. Assume that G is an infinitely generated minimal non- $(L\mathfrak{F}_\pi)\mathfrak{X}$ -group. Then by Proposition 2.3, $T_\pi \in L\mathfrak{F}_\pi$ and G/T_π is an infinite π -free locally nilpotent minimal non- $\mathfrak{X}$ -group. Therefore there exists a prime $p \notin \pi$ such that G/T_π is a countable p -group. It follows that G is locally finite, so that we can

deduce by [8, Lemma 1.D.4] that there exists a p -subgroup P of G such that $G = PT_\pi$. Therefore G is an infinitely generated minimal non- $\mathfrak{X}$ p -group by Corollary 2.4.

The converse is clear. $\square$

One can deduce the following consequence of Theorem 4.1.

Corollary 4.2. *An infinitely generated group G having non-trivial π -elements and whose proper subgroups are in $(L\mathfrak{F}_\pi)\mathfrak{X}$ is itself in $(L\mathfrak{F}_\pi)\mathfrak{X}$.*

Since by [10] and [14] (respectively, by [2, Theorem 1.2]) an infinitely generated minimal non-nilpotent (respectively, non-Baer) group is a countable p -group for some prime p , then we can take $\mathfrak{X}$ to be the class $\mathfrak{N}$ (respectively, of Baer groups).

Other possibility for the class $\mathfrak{X}$ is the class ZA of hypercentral groups as by [6, Lemma 2.2] an infinitely generated minimal non- ZA -group is periodic and since ZA is a $\mathbf{N}_0$ -closed class (i.e. the product of two normal hypercentral subgroups is hypercentral) [13, p. 51 of Part 1] and a countably recognizable class (i.e. a group is hypercentral whenever its countable subgroups are hypercentral) [13, Corollary 2 of Theorem 8.34].

If $G = A\langle x \rangle$ is a locally dihedral 2-group, that is the semi-direct product of a quasicyclic 2-group A by a cyclic group $\langle x \rangle$ of order 2 which acts by inversion on A , then it is easy to see that G is an infinitely generated minimal non-nilpotent group. So that if π is a set of odd primes then G is a minimal non- $(L\mathfrak{F}_\pi)\mathfrak{N}$ -group.

5. Minimal non- $(L\mathfrak{F}_\pi)\mathfrak{V}$ -groups

Now let $\mathfrak{V}$ be a variety of locally nilpotent groups. In this section we will prove that there are no infinite locally graded minimal non- $(L\mathfrak{F}_\pi)\mathfrak{V}$ -groups.

To prove next lemma we adapt the proof of its analog on the variety $\mathfrak{N}_c$ [5, Corollary 2].

Lemma 5.1. *Let G be a locally graded minimal non- $\mathfrak{V}$ -group. Then G is finite.*

PROOF. Let G as stated and assume for a contradiction that G is infinite. Then G is obviously finitely generated and hence it is nilpotent-by-finite so that G satisfies the maximal condition on subgroups. It follows that every proper subgroup of G is nilpotent. If G is not nilpotent, then it is a minimal non-nilpotent group and hence $G/\text{Frat}(G)$ is simple by [10, Theorem 3.3]. Therefore

$G/\text{Frat}(G)$ is finite and hence so is G by [9], a contradiction. It follows that G is an infinite nilpotent group. It is known that therefore G is (torsion-free)-by-finite. Let N be a normal subgroup of G such that N is torsion-free and G/N is finite. Hence N is a residually finite p -group for every prime p . Let p and q be distinct primes and let H and K be proper normal subgroups of N such that N/H and N/K are respectively of p -power and q -power order. Then $N/H \cap K$ is a finite nilpotent group with two primary components. So $G/H_G \cap K_G$ is a finite nilpotent group with at least two primary components and hence it is a direct product of two proper subgroups, so that it belongs to $\mathfrak{V}$. It follows that $G/H_G \in \mathfrak{V}$ and hence $G \in \mathfrak{V}$ as N is a residually finite p -group, a contradiction which implies that G is finite. $\square$

Proposition 5.2. *Let G be an infinite minimal non- $(L\mathfrak{F}_\pi)\mathfrak{V}$ -group. Then G is a finitely generated perfect group which has no proper subgroups of finite index and $G/\text{Frat}(G)$ is simple.*

PROOF. Let G as stated and assume for a contradiction that G is not finitely generated. Then by Proposition 2.3, $T_\pi \in L\mathfrak{F}_\pi$ and G/T_π is a locally nilpotent minimal non- $\mathfrak{V}$ -group and hence it is finite by Lemma 5.1. It follows that there exists a (finitely generated) proper subgroup F of G such that $G = T_\pi F$. We deduce by condition (i) of Proposition 2.3 that $G = F$, a contradiction. Therefore G is finitely generated and hence by Proposition 3.3 G is perfect, has no proper subgroups of finite index and $G/\text{Frat}(G)$ is simple. $\square$

Proposition 5.2 has the following consequence.

Corollary 5.3. *Let G be an infinite locally graded group whose proper subgroups are in $(L\mathfrak{F}_\pi)\mathfrak{V}$. Then G is a $(L\mathfrak{F}_\pi)\mathfrak{V}$ -group*

Clearly $\mathfrak{V}$ can stand for the variety $\mathfrak{N}_c$. According to the solution of the Restricted Burnside Problem [17], [18] the class of locally nilpotent groups of exponent a given positive integer n is a variety and hence it is another possibility for the variety $\mathfrak{V}$.

In [11] it is constructed an infinite simple torsion-free finitely generated group whose proper subgroups are cyclic. This group is both a minimal non- $(L\mathfrak{F}_\pi)\mathfrak{N}$ -group and a minimal non- $(L\mathfrak{F}_\pi)\mathfrak{N}_1$ -group for every set of primes π .

6. Some consequences

In this section we deduce some results on infinitely generated minimal non- $\mathfrak{C}_\pi\mathfrak{N}$ and non- $\mathfrak{F}_\pi\mathfrak{N}$ -groups (respectively, non- $\mathfrak{C}_\pi\mathfrak{N}_c$ and non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -groups), where

$\mathfrak{C}_\pi = L\mathfrak{F}_\pi \cap \mathfrak{C}$, $\mathfrak{F}_\pi = L\mathfrak{F}_\pi \cap \mathfrak{F}$, $\mathfrak{C}$ is the class of Chernikov groups and c is a positive integer.

In [1, Theorem 2.1], it is proved that a locally graded group is a minimal non- $\mathfrak{C}\mathfrak{N}$ -group if and only if it is a minimal non- $\mathfrak{N}$ -group without maximal subgroups. Using Theorem 4.1 we generalise this result.

Corollary 6.1. *Let G be an infinitely generated group. Then G is a minimal non- $\mathfrak{C}_\pi\mathfrak{N}$ -group if and only if G is a minimal non- $\mathfrak{N}$ -group such that either G has no maximal subgroups or there exists a prime $p \notin \pi$ such that G is a p -group having maximal subgroups.*

PROOF. Assume that G is a minimal non- $\mathfrak{C}_\pi\mathfrak{N}$ -group. First we prove that the product of two normal $\mathfrak{C}_\pi\mathfrak{N}$ -subgroups of G is likewise a $\mathfrak{C}_\pi\mathfrak{N}$ -group. For if H and K are two normal $\mathfrak{C}_\pi\mathfrak{N}$ -subgroups of G , then for some positive integers c and d , $\gamma_c(H)$ and $\gamma_d(K)$ are two normal $\mathfrak{C}_\pi$ -subgroups of G and hence so is $\gamma_c(H)\gamma_d(K)$. Since

$$HK/\gamma_c(H)\gamma_d(K) = (H\gamma_d(K)/\gamma_c(H)\gamma_d(K))(K\gamma_c(H)/\gamma_c(H)\gamma_d(K))$$

is nilpotent, we deduce that HK is a $\mathfrak{C}_\pi\mathfrak{N}$ -group. It follows by [10, Theorem 2.12] that every nilpotent image of G is abelian. Clearly G is either a $\mathfrak{C}\mathfrak{N}$ -group or a minimal non- $\mathfrak{C}\mathfrak{N}$ -group. Therefore G is either a $\mathfrak{C}\mathfrak{A}$ -group, $\mathfrak{A}$ being the class of abelian groups, or a minimal non- $\mathfrak{N}$ -group without maximal subgroups by [1, Theorem 2.1]. By Theorem 4.1 we have also that G is either a $(L\mathfrak{F}_\pi)\mathfrak{A}$ -group or there exists a prime $p \notin \pi$ such that G is a minimal non- $\mathfrak{N}$ - p -group. Consequently, if we assume that G is not a minimal non- $\mathfrak{N}$ -group, then $G \in \mathfrak{C}\mathfrak{A} \cap (L\mathfrak{F}_\pi)\mathfrak{A}$, so that $G' \in (L\mathfrak{F}_\pi) \cap \mathfrak{C} = \mathfrak{C}_\pi$, which is a contradiction. Hence G is a minimal non- $\mathfrak{N}$ -group and either G has no maximal subgroups or for some prime $p \notin \pi$ G is a p -group having maximal subgroups.

Conversely, if G is a minimal non- $\mathfrak{N}$ -group having maximal subgroups, then G is π -free and hence $G \notin \mathfrak{C}_\pi\mathfrak{N}$ which gives that G is a minimal non- $\mathfrak{C}_\pi\mathfrak{N}$ -group. Now if G is a minimal non- $\mathfrak{N}$ -group without maximal subgroups, then it is a minimal non- $\mathfrak{C}\mathfrak{N}$ -group by [1, Theorem 2.1] and hence $G \notin \mathfrak{C}_\pi\mathfrak{N}$ which again gives that G is a minimal non- $\mathfrak{C}_\pi\mathfrak{N}$ -group. $\square$

In [12, Theorem 1], it is proved that a locally graded group whose proper subgroups are in $\mathfrak{C}\mathfrak{N}_c$ is itself a $\mathfrak{C}\mathfrak{N}_c$ -group. We generalise this result using Corollary 5.3.

Corollary 6.2. *Let G be an infinite locally graded group whose proper subgroups are in $\mathfrak{C}_\pi\mathfrak{N}_c$. Then G is a $\mathfrak{C}_\pi\mathfrak{N}_c$ -group.*

PROOF. Let G as stated; then G belongs to $\mathfrak{C}\mathfrak{N}_c$ by [12, Theorem 1]. Since G is a $(L\mathfrak{F}_\pi)\mathfrak{N}_c$ -group by Corollary 5.3, we deduce that G belongs to $\mathfrak{C}_\pi\mathfrak{N}_c$. $\square$

Corollary 6.2 has the following consequence which will be used in next results.

Lemma 6.3. *If G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -group, then G is a minimal non- $\mathfrak{F}\mathfrak{N}_c$ -group.*

PROOF. Let G be a minimal non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -group. Then by Corollary 6.2, G belongs to $\mathfrak{C}_\pi\mathfrak{N}_c$ and hence $G \notin \mathfrak{F}\mathfrak{N}_c$. Therefore G is a minimal non- $\mathfrak{F}\mathfrak{N}_c$ -group. $\square$

Combining [16, Theorem 3.5] and the fact that infinitely generated minimal non- $\mathfrak{N}$ -groups are soluble by [3, Theorem 1.3] we have that a group is an infinitely generated minimal non- $\mathfrak{F}\mathfrak{N}$ -group if and only if it is an infinitely generated minimal either non- $\mathfrak{N}$ -group or non- $\mathfrak{F}\mathfrak{A}$ -group. We generalise this result using Theorem 4.1.

Corollary 6.4. *Let G be an infinitely generated group. Then G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ -group if and only if G is either a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$ -group, or a minimal non- $\mathfrak{N}$ -group which is either without maximal subgroups, or there exists a prime $p \notin \pi$ such that G is a p -group having maximal subgroups.*

PROOF. Assume that G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ -group. As in the proof of Corollary 6.1, one can prove that every nilpotent image of G is abelian. Clearly G is either a $\mathfrak{F}\mathfrak{N}$ -group or a minimal non- $\mathfrak{F}\mathfrak{N}$ -group. So G is either a $\mathfrak{F}\mathfrak{A}$ -group, or a minimal non- $\mathfrak{F}\mathfrak{A}$ -group or a minimal non- $\mathfrak{N}$ -group without maximal subgroups. On the other hand G is either a $L(\mathfrak{F}_\pi)\mathfrak{N}$ -group or a minimal non- $L(\mathfrak{F}_\pi)\mathfrak{N}$ -group, which implies that G is either a $L(\mathfrak{F}_\pi)\mathfrak{A}$ -group or a minimal non- $\mathfrak{N}$ - p -group for some prime $p \notin \pi$. Therefore if G is not a minimal non- $\mathfrak{N}$, then G is in $L(\mathfrak{F}_\pi)\mathfrak{A}$ and is either a $\mathfrak{F}\mathfrak{A}$ -group or a minimal non- $\mathfrak{F}\mathfrak{A}$ -group. Since clearly $G \notin \mathfrak{F}\mathfrak{A}$, we have that G is a minimal non- $\mathfrak{F}\mathfrak{A}$ -group which is a $(L\mathfrak{F}_\pi)\mathfrak{A}$ and hence it is a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$.

Conversely, if G is a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$ -group, then G is a minimal non- $\mathfrak{F}\mathfrak{A}$ -group by Lemma 6.3 and hence G is a minimal non- $\mathfrak{F}\mathfrak{N}$ -group by [16, Theorem 3.5]. We deduce that $G \notin \mathfrak{F}_\pi\mathfrak{N}$ and hence G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ -group. If G is a minimal non- $\mathfrak{N}$ - p -group for some prime $p \notin \pi$, then G is π -free and hence $G \notin \mathfrak{F}_\pi\mathfrak{N}$, so that G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ -group. Now if G is a minimal non- $\mathfrak{N}$ - p -group without maximal subgroups, for some prime $p \in \pi$, then G is a minimal non- $\mathfrak{F}\mathfrak{N}$ by [16, Theorem 3.5] and a $L(\mathfrak{F}_\pi)\mathfrak{A}$ -group by Theorem 4.1 and hence G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ -group. $\square$

In [4, Theorem 2], it is proved that a locally graded group is a minimal non- $\mathfrak{F}\mathfrak{N}_c$ -group if and only if it is a minimal non- $\mathfrak{F}\mathfrak{A}$ -group. We generalise this result using Corollary 5.3.

Corollary 6.5. *An infinitely generated group G is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -group if and only if G is a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$ -group.*

PROOF. Let G be a minimal non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -group. Then G is a minimal non- $\mathfrak{F}\mathfrak{N}_c$ -group by Lemma 6.3. It follows that G is a minimal non- $\mathfrak{F}\mathfrak{A}$ by [4, Theorem 2] and hence G is a minimal non- $\mathfrak{F}\mathfrak{N}$ -group by [16, Theorem 3.5]. Since the product of two normal $\mathfrak{F}\mathfrak{N}$ -subgroups of G is a $\mathfrak{F}\mathfrak{N}$ -group and as G is a $(L\mathfrak{F}_\pi)\mathfrak{N}_c$ -group by Corollary 5.3, we deduce that G belongs to $(L\mathfrak{F}_\pi)\mathfrak{A}$. Consequently G is a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$ -group.

Conversely assume that G is a minimal non- $\mathfrak{F}_\pi\mathfrak{A}$ -group. Then clearly every proper subgroup of G is in $\mathfrak{F}_\pi\mathfrak{N}_c$. Since G cannot be a $\mathfrak{F}_\pi\mathfrak{N}_c$ -group by Lemma 6.3, it is a minimal non- $\mathfrak{F}_\pi\mathfrak{N}_c$ -group. $\square$

It is easy to see that the well known example of Heineken-Mohamed [7], which is a minimal non- $\mathfrak{N}$ -group without maximal subgroups, is both a minimal non- $\mathfrak{F}_\pi\mathfrak{N}$ and non- $\mathfrak{C}_\pi\mathfrak{N}$ -group for all sets π of primes.

ACKNOWLEDGEMENT. Both authors are grateful to the referees for careful reading and for their invaluable comments.

References

- [1] A. ARIKAN and N. TRABELSI, On groups whose proper subgroups are Chernikov-by-Baer or (periodic divisible abelian)-by-Baer, *J. Algebra Appl.* **12** (2013), 1–15.
- [2] A. ARIKAN and N. TRABELSI, On minimal non-Baer-groups, *Comm. Algebra* **39** (2011), 2489–2497.
- [3] A. O. ASAR, Locally nilpotent p -groups whose proper subgroups are hypercentral or nilpotent-by-Chernikov, *J. London Math. Soc.* **61** (2000), 412–422.
- [4] B. BRUNO and R. E. PHILLIPS, On minimal conditions related to Miller-Moreno type groups, *Rend. Sem. Mat. Univ. Padova* **69** (1983), 153–168.
- [5] B. BRUNO and R. E. PHILLIPS, Groups with restricted non-normal subgroups, *Math. Z* **176** (1981), 199–221.
- [6] S. FRANCIOSI, F. DE GIOVANNI and Y. P. SYSAK, Groups with many polycyclic-by-nilpotent subgroups, *Ric. Mat.* **48** (1999), 361–378.
- [7] H. HEINEKEN and I. J. MOHAMED, A group with trivial centre satisfying the normalizer condition, *J. Algebra* **10** (1968), 368–376.
- [8] O. H. KEGEL and B. A. F. WEHRFRITZ, Locally Finite Groups, *North-Holland, Amsterdam*, 1973.

- [9] J. C. LENNOX, Polycyclic Frattini factors in certain finitely generated groups, *Arch. Math.* **24** (1973), 571–578.
- [10] M. F. NEWMAN and J. WIEGOLD, Groups with many nilpotent subgroups, *Arch. Math.* **15** (1964), 241–250.
- [11] A. Y. OL'SHANSKII, An infinite simple torsion-free Noetherian group, *Izv. Akad. Nauk. SSSR Ser. Mat.* **43** (1979), 1328–1393.
- [12] J. OTAL and J. M. PENA, Groups in which every proper subgroup is Chernikov-by-nilpotent or nilpotent-by-Chernikov, *Arch. Math.* **51** (1988), 193–197.
- [13] D. J. S. ROBINSON, Finiteness Conditions and Generalized Soluble Groups, *Springer Verlag, Berlin*, 1972.
- [14] H. SMITH, Groups with few non-nilpotent subgroups, *Glasgow Math. J.* **39** (1997), 141–151.
- [15] N. TRABELSI, On minimal non-(torsion-by-nilpotent) and non-((locally finite)-by-nilpotent) groups, *C. R. Acad. Sci. Paris Ser. I* **344** (2007), 353–356.
- [16] M. XU, Groups whose proper subgroups are finite-by-nilpotent, *Arch. Math.* **66** (1996), 353–359.
- [17] E. I. ZEL'MANOV, Solution of the restricted Burnside problem for 2-groups, *USSR-Sb.* **72** (1992), 543–565.
- [18] E. I. ZEL'MANOV, Solution of the restricted Burnside problem for groups of odd exponent, *Math. USSR-Izv.* **36** (1991), 41–60.

AMEL DILMI
LABORATORY OF FUNDAMENTAL
AND NUMERICAL MATHEMATICS
DEPARTMENT OF MATHEMATICS
UNIVERSITY SETIF 1
SETIF 19000
ALGERIA

E-mail: di.amel@yahoo.fr

NADIR TRABELSI
LABORATORY OF FUNDAMENTAL
AND NUMERICAL MATHEMATICS
DEPARTMENT OF MATHEMATICS
UNIVERSITY SETIF 1
SETIF 19000
ALGERIA

E-mail: nadir.trabelsi@yahoo.fr

(Received September 10, 2014; revised February 3, 2015)