

## A minimax theorem not involving convexities of the function

By I. JOÓ (Budapest)

*Dedicated to Professor Ákos Császár for his 70<sup>th</sup> birthday*

In this paper we prove the following

**Theorem.** *Let  $X$  be a compact topological space,  $Y$  be an arbitrary topological space and  $f : X \times Y \rightarrow \mathbb{R}$  a function continuous in  $x$  for every fixed  $y \in Y$  and lower semicontinuous in  $y$  for every fixed  $x \in X$ . This means that the level sets  $H_c^y = \{x : f(x, y) \geq c\}$  and  $H_x^c = \{y : f(x, y) \leq c\}$  are closed in  $X$  resp in  $Y$  and  $\hat{H}_c^y = \{x : f(x, y) \succ c\}$  are open in  $X$ . Suppose further that*

a) *The intersection of any (not necessarily finite numbers of  $H_x^c$  sets),  $x \in X$ ,  $c \in \mathbb{R}$  is connected (may be empty).*

b) *The intersection of any finitely many  $\hat{H}_c^y$  is connected (may be empty).*

Then

$$\sup_x \inf_y f(x, y) = \inf_y \sup_x f(x, y).$$

PROOF of the Theorem. Define an interval structure on  $Y$  as follows: let  $[y_1, y_2] = \bigcap_{x,c} \{H_x^c : y_1, y_2 \in H_x^c\}$ .

If we choose the constant  $c$  sufficiently large, we obtain  $y_1, y_2 \in H_x^c$ . Hence  $[y_1, y_2]$  is well-defined, closed connected and  $y_1, y_2 \in [y_1, y_2]$ . The sets  $H_x^c$  become convex in the sense that  $y_1, y_2 \in H_x^c$  implies  $[y_1, y_2] \in H_x^c$ . Consequently we can apply Theorem B of JOÓ [1] stating that if

- (1)  $H_x^c \cap [y_1, y_2]$  is closed in  $[y_1, y_2]$
- (2)  $H_x^c$  is convex

(3)  $\bigcap_{i=1}^n \hat{H}_c^{y_i}$  is connected for every finite intersection  
 then  $\bigcap_{i=1}^n \hat{H}_c^{y_i} \neq \emptyset$  for every  $c < c^y = \inf_y \sup_x f$  hence the sets  $H_c^y$  have also the finite intersection property. In our case (1) holds since  $H_x^c$  and  $[y_1, y_2]$  are closed in  $Y$ , (2) was remarked above and (3) is identical to b), hence Theorem B of [1] applies. Since the sets  $H_c^y$  are compact, this implies  $\bigcap_{y \in Y} H_c^y \neq \emptyset$  i.e. there exist  $x_0 \in X$  with  $f(x_0, y) \geq c$  for all  $y \in Y$ . Hence  $c < c^*$  implies  $c \leq \sup_x \inf_y f$  i.e.  $c^* \leq \sup_x \inf_y f$  which proves the desired minimax equality.  $\square$

### References

- [1] I. Joó, On a fixed point theorem, *Publicationes Mathematicae, Debrecen* **36** (1989), 127–128.

I. JOÓ  
 DEPARTMENT OF ANALYSIS  
 L. EÖTVÖS UNIVERSITY  
 M-1088-BUDAPEST  
 MÚZEUM KRT. 6-8.

(Received November 15, 1993; revised May 13, 1994)