

Title: GK-dimension of 2×2 generic Lie matrices

Author(s): Vesselin Drensky, Plamen Koshlukov and Gustavo Grings Machado

Recently Machado and Koshlukov have computed the Gelfand–Kirillov dimension of the relatively free algebra $F_m = F_m(\text{var}(sl_2(K)))$ of rank m in the variety of algebras generated by the three-dimensional simple Lie algebra $sl_2(K)$ over an infinite field K of characteristic different from 2. They have shown that $\text{GKdim}(F_m) = 3(m - 1)$. The algebra F_m is isomorphic to the Lie algebra generated by m generic 2×2 matrices. Now we give a new proof for $\text{GKdim}(F_m)$ using classical results of Procesi and Razmyslov combined with the observation that the commutator ideal of F_m is a module of the center of the associative algebra generated by m generic traceless 2×2 matrices.

Address:

Vesselin Drensky
Institute of Mathematics and Informatics
Bulgarian Academy of Sciences
Acad. G. Bonchev Str., Block 8
1113 Sofia
Bulgaria

Address:

Plamen Koshlukov
Department of Mathematics
IMECC, UNICAMP
Sérgio Buarque de Holanda 651
Campinas, SP 13083-859
Brazil

Address:

Gustavo Grings Machado
Department of Mathematics
CCNE, UFSM
Faixa de Camobi, Campus UFSM
Santa Maria, RS 97105-900
Brazil