

Conformal vector fields on submanifolds of a Euclidean space

By HANAN ALOHALI (Riyadh), HAILA ALODAN (Riyadh)
and SHARIEF DESHMUKH (Riyadh)

Abstract. In this paper, we investigate n -dimensional immersed compact submanifold M of a Euclidean space R^{n+p} , with the immersion $\psi : M \rightarrow R^{n+p}$, where the tangential component ψ^T of ψ is a conformal vector field. A characterization of n -sphere in the Euclidean space R^{n+p} is obtained. Also conditions under which ψ^T is a conformal vector field in the general case and those in the special case where the submanifold has flat normal connection and $p = 2$ are obtained as well.

1. Introduction

Given an immersed n -dimensional submanifold M of a Euclidean space $(R^{n+p}, \langle, \rangle)$, where $\langle, \rangle$ is the Euclidean metric, one of the important questions is to find conditions under which the submanifold M lies on the hypersphere $S^{n+p-1}(c)$ of the Euclidean space R^{n+p} . This question has been studied in [ALO07], [ALO02], [ALOD02]. Recall that a smooth vector field ξ on a Riemannian manifold (M, g) is said to be a conformal vector field if its flow consists of conformal transformations of the Riemannian manifold (M, g) and it is equivalent to the requirement that the vector field ξ satisfies

$$\mathcal{L}_\xi g = 2\rho g,$$

where $\mathcal{L}_\xi$ is the Lie derivative with respect to the vector field ξ , and ρ is a smooth function on M , called the potential function of the conformal vector

Mathematics Subject Classification: 53C20, 53A50.

Key words and phrases: Ricci curvature, conformal gradient vector field, flat normal vector field, submanifolds.

This work is supported by the Deanship of Scientific Research of King Saud University, College of Science, Research Center.

field ξ . Conformal vector fields have been used to characterize spheres among compact Riemannian manifolds (cf. [DES12], [DES08], [DES10]). If M is an n -dimensional immersed submanifold of the Euclidean space R^{n+p} with the immersion $\psi : M \rightarrow R^{n+p}$, then treating ψ as the position vector field of points of M , we can express it as

$$\psi = \psi^T + \psi^\perp,$$

where ψ^T is the tangential component of ψ to M , and $\psi^\perp$ is the normal component of ψ . Thus, we get a globally defined vector field ψ^T on the submanifold M , which might be either a Killing vector field or a conformal vector field. However, the covariant derivative of ψ^T being symmetric (see Section 2), asking ψ^T be a Killing vector field, will not yield interesting geometry. Therefore, it is a natural question to find conditions under which the vector field ψ^T is a conformal vector field on M , as well as to study the geometry of the submanifold for which the vector field ψ^T is a conformal vector field. In this paper, we address these questions. It is interesting to note that in the case when ψ^T is a nonzero conformal vector field on the compact submanifold M , under suitable restrictions on the Ricci curvatures, the submanifold is shown to be isometric to the sphere $S^n(c)$ of constant curvature c (cf. Theorem 3.1). We also find conditions under which the vector field ψ^T is a conformal vector field on the submanifold M (cf. Theorems 3.2 and 4.1). Finally, we use the conformal vector field associated to the normal component $\psi^\perp$ on the submanifold M to find a necessary and sufficient condition for the submanifold to lie on the hypersphere $S^{n+p-1}(c)$ (cf. Theorem 3.3).

2. Preliminaries

Let M be an n -dimensional submanifold of the Euclidean space R^{n+p} with immersion $\psi : M \rightarrow R^{n+p}$. We denote by $\langle, \rangle$ and $\bar{\nabla}$ the Euclidean metric and the Euclidean connection, respectively, on R^{n+p} , we also denote by g and ∇ the induced metric and the Riemannian connection on the submanifold M . Then, we have the following equations for the submanifold M (cf. [CHE83]):

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \bar{\nabla}_X N = -A_N X + \nabla_X^\perp N \quad (2.1)$$

$X, Y \in \mathfrak{X}(M)$, $N \in \Gamma(\Lambda)$, where $\mathfrak{X}(M)$ is the Lie algebra of smooth vector fields on M , $\Gamma(\Lambda)$ is the space of smooth sections of the normal bundle Λ of M , h is the second fundamental form, A_N is the Weingarten map with respect to the normal $N \in \Gamma(\Lambda)$ which is related to the second fundamental form h by

$$g(A_N X, Y) = g(h(X, Y), N), \quad X, Y \in \mathfrak{X}(M),$$

and $\nabla^\perp$ is the connection in the normal bundle Λ . We also have the Gauss equation

$$R(X, Y)Z = A_{h(Y, Z)}X - A_{h(X, Z)}Y, \quad X, Y, Z \in \mathfrak{X}(M), \quad (2.2)$$

where $R(X, Y)Z$, $X, Y, Z \in \mathfrak{X}(M)$ is the curvature tensor field of the submanifold M . The Ricci tensor field of M is given by

$$\text{Ric}(X, Y) = ng(h(X, Y), H) - \sum_{i=1}^n g(h(X, e_i), h(Y, e_i)), \quad (2.3)$$

where $\{e_1, \dots, e_n\}$ is a local orthonormal frame on M , and

$$H = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i)$$

is the mean curvature vector field.

The Ricci operator Q is the symmetric operator defined by

$$\text{Ric}(X, Y) = g(Q(X), Y), \quad X, Y \in \mathfrak{X}(M).$$

If we express $\psi = \psi^T + \psi^\perp$, where $\psi^T \in \mathfrak{X}(M)$ is the tangential component and $\psi^\perp \in \Gamma(\Lambda)$ is the normal component of ψ , and if we denote by $B = A_{\psi^\perp}$ the Weingarten map with respect to the normal vector field $\psi^\perp$, then using equation (2.1), we get

$$\nabla_x \psi^T = X + BX, \quad \nabla_x^\perp \psi^\perp = -h(X, \psi^T), \quad X, Y \in \mathfrak{X}(M). \quad (2.4)$$

We use the mean curvature vector field H to define a smooth function $F : M \rightarrow R$ on the submanifold M by $F = \langle H, \psi^\perp \rangle$. Now, for an n -dimensional submanifold $\psi : M \rightarrow R^{n+p}$, and a local orthonormal frame $\{e_1, \dots, e_n\}$ on M , we have

$$\begin{aligned} \text{div } \psi^T &= \sum_{i=1}^n \langle \nabla_{e_i} \psi^T, e_i \rangle = \sum_{i=1}^n \langle e_i + A_{\psi^\perp} e_i, e_i \rangle \\ &= n + \sum_{i=1}^n \langle h(e_i, e_i), \psi^\perp \rangle = n + n \langle H, \psi^\perp \rangle = n(1 + F), \end{aligned}$$

that is,

$$\text{div } \psi^T = n(1 + F). \quad (2.5)$$

We have the following Lemmas:

Lemma 2.1 (Hsiung–Minkowski formula). *Let M be an n -dimensional compact submanifold of the Euclidean space R^{n+p} . Then*

$$\int_M (1 + F) dv = 0.$$

Lemma 2.2 ([ALO07]). *Let M be an n -dimensional submanifold of R^{n+p} . Then the tensor field B satisfies*

- (i) $\text{Tr } B = nF$;
- (ii) $(\nabla B)(X, Y) - (\nabla B)(Y, X) = R(X, Y)\psi^T$;
- (iii) $\sum_{i=1}^n (\nabla B)(e_i, e_i) = n\nabla F + Q(\psi^T)$;

where $(\nabla B)(X, Y) = \nabla_X BY - B\nabla_X Y$, $X, Y \in \mathfrak{X}(M)$.

Lemma 2.3 ([ALO07]). *Let $\psi : M \rightarrow R^{n+p}$ be an n -dimensional compact submanifold. Then a necessary and sufficient condition for $\psi(M) \subset S^{n+p-1}(c)$ is that $\psi^T = 0$ and $F = -1$.*

Definition 2.1. A smooth vector field ξ on a Riemannian manifold (M, g) is said to be a conformal vector field if there exists a smooth function ρ on M that satisfies $\mathcal{L}_\xi g = 2\rho g$, ρ called a potential function, where $\mathcal{L}_\xi g$ is the Lie derivative of g with respect to ξ . We say that ξ is a non-trivial conformal vector field if the potential function ρ is not a constant. A conformal vector field ξ is said to be a gradient conformal vector field if $\xi = \nabla f$, for a smooth function f on M .

Using Koszul's formula, we immediately obtain the following for a vector field ξ on M :

$$2g(\nabla_X \xi, Y) = (\mathcal{L}_\xi g)(X, Y) + d\eta(X, Y), \quad X, Y \in \mathfrak{X}(M),$$

where η is the 1-form dual to ξ , that is, $\eta(X) = g(X, \xi)$, $X \in \mathfrak{X}(M)$. Define a skew-symmetric tensor field φ of type $(1, 1)$ on M by $d\eta(X, Y) = 2g(\varphi X, Y)$, and a symmetric tensor field C of type $(1, 1)$ by

$$\mathcal{L}_\xi g(X, Y) = 2g(CX, Y), \quad X, Y \in \mathfrak{X}(M),$$

then, for a smooth vector field ξ on M , we have

$$\nabla_X \xi = CX + \varphi X, \quad X, Y \in \mathfrak{X}(M). \quad (2.6)$$

Using the definition of a conformal vector field and equation (2.6), we have

Lemma 2.4 ([DES12]). *Let ξ be a conformal vector field on an n -dimensional Riemannian manifold (M, g) , with potential function ρ . Then*

$$\nabla_X \xi = \rho X + \varphi X, \quad X \in \mathfrak{X}(M) \quad \text{and} \quad \operatorname{div} \xi = n\rho.$$

Remark 2.1 ([DES08]). Let ξ be a conformal gradient vector field on a compact Riemannian manifold (M, g) . Then, for $\rho = n^{-1} \operatorname{div} \xi$,

$$\int_M \rho dv = 0.$$

Let λ_1 be the nonzero eigenvalue of the Laplacian operator Δ acting on the smooth functions of a compact Riemannian manifold (M, g) , where we adopt the sign convention of the Laplacian operator as $\Delta f = \operatorname{div} \nabla f$. Then, for a smooth function f on M satisfying

$$\int_M f dv = 0,$$

by minimum principle we have

$$\int_M \|\nabla f\|^2 dv \geq \lambda_1 \int_M f^2 dv, \quad (2.7)$$

and the equality holds if and only if $\Delta f = -\lambda_1 f$. Moreover, for a smooth function f , the Hessian operator H_f is given by

$$H_f X = \nabla_X \nabla f, \quad X \in \mathfrak{X}(M),$$

and on a compact Riemannian manifold, we have the following Bochner formula:

$$\int_M \left\{ \operatorname{Ric}(\nabla f, \nabla f) + \|H_f\|^2 - (\Delta f)^2 \right\} dv = 0. \quad (2.8)$$

3. Submanifolds with ψ^T as conformal vector field

Let M be an n -dimensional submanifold of the Euclidean space R^{n+p} , with immersion $\psi : M \rightarrow R^{n+p}$. In this section, we study the geometry of the submanifold M for which the vector field ψ^T is a conformal vector field. First, we prove the following Lemmas.

Lemma 3.1. *Let M be an n -dimensional submanifold of the Euclidean space R^{n+p} , with immersion $\psi : M \rightarrow R^{n+p}$ and $f = \frac{1}{2} \|\psi^\perp\|^2$. If the gradient ∇f of the smooth function f is a conformal vector field, then*

$$\text{Ric}(\psi^T, \psi^T) + n\psi^T(F) + n\rho + nF + \|B\|^2 = 0,$$

where ρ is the potential function of ∇f .

PROOF. As ∇f is a conformal vector field with potential function say ρ , we have

$$\mathcal{L}_{\nabla f} g = 2\rho g.$$

Since the 1-form dual to the conformal vector field ∇f is closed, we have $\varphi = 0$, and Lemma 2.4 takes the form

$$\nabla_X(\nabla f) = \rho X \quad \text{and} \quad \Delta f = n\rho, \quad (3.1)$$

where Δ is the Laplacian operator. Now, for $X \in \mathfrak{X}(M)$, we have

$$\begin{aligned} g(\nabla f, X) &= X(f) = X\left(\frac{1}{2} \|\psi^\perp\|^2\right) = g(\bar{\nabla}_X \psi^\perp, \psi^\perp) = g(-A_{\psi^\perp} X + \nabla_X^\perp \psi^\perp, \psi^\perp) \\ &= g(\nabla_X^\perp \psi^\perp, \psi^\perp) = -g(h(X, \psi^T), \psi^\perp) = -g(A_{\psi^\perp} \psi^T, X), \end{aligned}$$

which gives $\nabla f = -A_{\psi^\perp} \psi^T = -B\psi^T$. Putting $\xi = \psi^T$, we get $\nabla f = -B\xi$, and consequently,

$$\nabla_X(\nabla f) = -\nabla_X B\xi = -[(\nabla B)(X, \xi) + B\nabla_X \xi],$$

which, using equation (2.4), gives

$$\begin{aligned} \nabla_X(\nabla f) &= -(\nabla B)(X, \xi) - B(X + BX) \\ &= -(\nabla B)(X, \xi) - BX - B^2 X. \end{aligned} \quad (3.2)$$

Now, using Lemma 2.2 and the fact that B is a symmetric operator, we have

$$\begin{aligned} \sum_{i=1}^n g((\nabla B)(e_i, \xi), e_i) &= g\left(\sum_{i=1}^n (\nabla B)(e_i, e_i), \xi\right) \\ &= g(n\nabla F + Q(\xi), \xi) = n\xi(F) + \text{Ric}(\xi, \xi). \end{aligned} \quad (3.3)$$

Also, using equations (3.1) and (3.2), we get

$$\begin{aligned} \sum_{i=1}^n g((\nabla B)(e_i, \xi), e_i) &= \sum_{i=1}^n g(-\rho e_i - Be_i - B^2 e_i, e_i) \\ &= -n\rho - \text{Tr } B - \|B\|^2. \end{aligned} \quad (3.4)$$

Then, using $\text{Tr } B = nF$ and equations (3.3) and (3.4), we arrive at

$$\text{Ric}(\xi, \xi) + n\xi(F) + n\rho + nF + \|B\|^2 = 0,$$

which proves the Lemma. $\square$

Lemma 3.2. *Let $\psi : M \rightarrow R^{n+p}$ be an n -dimensional compact submanifold. Then*

$$\int_M \left\{ \text{Ric}(\psi^T, \psi^T) - n^2(1+F)^2 + \|B\|^2 - n \right\} dv = 0.$$

PROOF. Taking $\xi = \psi^T$, we have

$$\text{div}(F\xi) = g(\nabla F, \xi) + F \text{div } \xi = g(\nabla F, \xi) + nF(1+F).$$

Consider a local orthonormal frame $\{e_1, \dots, e_n\}$, then using Lemma 2.2 and equation (2.5) to compute $\text{div}(B\xi)$, we get

$$\begin{aligned} \text{div}(B\xi) &= \sum_{i=1}^n g(\nabla_{e_i} B\xi, e_i) = \sum_{i=1}^n g((\nabla B)(e_i, \xi) + B\nabla_{e_i} \xi, e_i) \\ &= \sum_{i=1}^n [g((\nabla B)(e_i, e_i), \xi) + g(\nabla_{e_i} \xi, Be_i)] \\ &= g(n\nabla F + Q(\xi), \xi) + \sum_{i=1}^n [g(e_i, Be_i) + g(Be_i, Be_i)] \\ &= ng(\nabla F, \xi) + \text{Ric}(\xi, \xi) + \text{Tr } B + \|B\|^2 \\ &= ng(\nabla F, \xi) + \text{Ric}(\xi, \xi) + nF + \|B\|^2, \end{aligned}$$

and

$$g(\nabla F, \xi) = \text{div}(F\xi) - nF^2 - nF,$$

which gives

$$ng(\nabla F, \xi) = n \text{div}(F\xi) - n^2 F^2 - n^2 F.$$

Consequently,

$$\text{div}(B\xi) = n \text{div}(F\xi) - n^2 F^2 - n^2 F + \text{Ric}(\xi, \xi) + nF + \|B\|^2,$$

and we have

$$\text{div}(B\xi - nF\xi) = \text{Ric}(\xi, \xi) - n^2 F^2 - n^2 F + nF + \|B\|^2,$$

which after integration gives

$$\int_M \left\{ \text{Ric}(\xi, \xi) - n^2 (F^2 - 1) + \|B\|^2 - n \right\} dv = 0. \quad (3.5)$$

Also using Lemma 2.1, we have

$$\int_M (1 + F)^2 dv = \int_M (F^2 - 1) dv,$$

which, together with equation (3.5), gives

$$\int_M \left\{ \text{Ric}(\xi, \xi) - n^2 (1 + F)^2 + \|B\|^2 - n \right\} dv = 0. \quad \square$$

Theorem 3.1. *Let $\psi : M \rightarrow R^{n+p}$ be an n -dimensional compact submanifold with the tangential component ψ^T , a nonzero conformal vector field with potential function ρ , and λ_1 be the first nonzero eigenvalue of the Laplacian operator on the submanifold M . If $c = n^{-1}\lambda_1$ and the Ricci tensor on M satisfies*

- (i) $\text{Ric}(\nabla\rho + c\psi^T, \nabla\rho + c\psi^T) \geq 0$,
 - (ii) $\text{Ric}(\nabla\rho, \nabla\rho) \leq (n-1)c\|\nabla\rho\|^2$,
- then M is isometric to a sphere $S^n(c)$.*

PROOF. Let $\xi = \psi^T$ be a conformal vector field with potential function ρ . If we define $f = \frac{1}{2}\|\psi\|^2$, then it is easy to show that $\xi = \nabla f$. Thus ξ is a gradient conformal vector field, and consequently, as the 1-form η dual to ξ being $\eta = df$ is closed, we get that $\varphi = 0$. Then, by Lemma 2.4, we have

$$\nabla_X \xi = \rho X,$$

and using equation (2.4) in the above equation, we have

$$BX + X = \rho X,$$

which gives $B = (\rho - 1)I$ and $\text{div } \xi = n\rho$. However, as $\xi = \nabla f$, we have $\Delta f = n\rho$.

Now,

$$(\nabla B)(X, Y) = \nabla_X BY - B\nabla_X Y = \nabla_X (\rho - 1)Y - (\rho - 1)\nabla_X Y = X(\rho)Y,$$

which, together with Lemma 2.2, gives

$$X(\rho)Y - Y(\rho)X = R(X, Y)\xi. \quad (3.6)$$

The above equation immediately gives

$$\operatorname{Ric}(\xi, X) = \sum_{i=1}^n R(e_i, X; \xi, e_i) = g(X, \nabla \rho) - nX(\rho),$$

and consequently, we have

$$Q(\xi) = -(n-1)\nabla \rho. \quad (3.7)$$

The above equation gives

$$\operatorname{Ric}(\xi, \xi) = -(n-1)\xi(\rho) = -(n-1)[\operatorname{div}(\rho\xi) - \rho \operatorname{div} \xi],$$

that is,

$$\operatorname{Ric}(\xi, \xi) = -(n-1)\operatorname{div}(\rho\xi) + n(n-1)\rho^2. \quad (3.8)$$

Also, equation (3.7) gives

$$\operatorname{Ric}(\xi, \nabla \rho) = g(-(n-1)\nabla \rho, \nabla \rho) = -(n-1)\|\nabla \rho\|^2. \quad (3.9)$$

Let λ_1 be the first nonzero eigenvalue of the Laplacian operator on M . Then Remark 2.1, together with equation (2.7), gives

$$\int_M \|\nabla \rho\|^2 dv \geq \lambda_1 \int_M \rho^2 dv, \quad (3.10)$$

with equality holding if and only if $\Delta \rho = -\lambda_1 \rho$.

Using $c = n^{-1}\lambda_1$ and equations (3.8), (3.9) and (3.10), we arrive at

$$\begin{aligned} & \int_M \operatorname{Ric}(\nabla \rho + c\xi, \nabla \rho + c\xi) dv \\ &= \int_M \left\{ \operatorname{Ric}(\nabla \rho, \nabla \rho) + n(n-1)c^2\rho^2 - 2(n-1)c\|\nabla \rho\|^2 \right\} dv \\ &\leq \int_M \left\{ \operatorname{Ric}(\nabla \rho, \nabla \rho) - (n-1)c\|\nabla \rho\|^2 \right\} dv, \end{aligned}$$

Using the conditions in the statement, and the above inequality, we conclude that

$$\operatorname{Ric}(\nabla \rho + c\xi, \nabla \rho + c\xi) = 0 \quad \text{and} \quad \operatorname{Ric}(\nabla \rho, \nabla \rho) - (n-1)c\|\nabla \rho\|^2 = 0. \quad (3.11)$$

Thus we have

$$\text{Ric}(\nabla\rho, \nabla\rho) + 2c \text{Ric}(\nabla\rho, \xi) + c^2 \text{Ric}(\xi, \xi) = 0,$$

which, together with equation (3.9) and the second equation in (3.11), gives

$$\text{Ric}(\xi, \xi) = (n-1)c^{-1} \|\nabla\rho\|^2. \quad (3.12)$$

Now, using $\nabla f = \xi$, that is, $H_f(X) = \rho X$ and $\Delta f = n\rho$ in the Bochner Formula (2.8), we arrive at

$$\int_M \{ \text{Ric}(\xi, \xi) + n\rho^2 - n^2\rho^2 \} dv = 0,$$

which, together with equation (3.12), gives

$$\int_M \|\nabla\rho\|^2 dv = nc \int_M \rho^2 dv = \lambda_1 \int_M \rho^2 dv.$$

This equality in (3.10) gives $\Delta\rho = -\lambda_1\rho$, which, together with $\Delta f = n\rho$, gives $\Delta(\rho + \lambda_1 n^{-1}f) = 0$, and on compact M , we have $\rho + \lambda_1 n^{-1}f = \text{constant}$. This last equation, together with $H_f(X) = \rho X$, gives $\nabla\rho = -c\nabla f$, that is,

$$\nabla_X \nabla\rho = -c\rho X. \quad (3.13)$$

If ρ is a constant, then we have $-c\nabla f = 0$, that is, $\xi = 0$, which is a contradiction, as ξ is a nonzero conformal vector field. Hence the nonconstant function ρ satisfies the OBATA's differential equation (3.13) (cf. [OBA62]), and therefore is isometric to the sphere $S^n(c)$. $\square$

In the following result, we consider the tangential component ψ^T and find conditions under which it becomes a conformal vector field on the submanifold M .

Theorem 3.2. *Let $\psi : M \rightarrow R^{n+p}$ be an n -dimensional compact submanifold, with $\lambda = \inf \frac{1}{n-1} \text{Ric} > 0$. If $\|\psi^T\|^2 \geq n\lambda^{-1}(1+F)^2$, then ψ^T is a conformal vector field on M .*

PROOF. Taking $\xi = \psi^T$ in Lemma 3.2, we get

$$\int_M \left\{ \text{Ric}(\xi, \xi) - n^2(1+F)^2 + \|B\|^2 - n \right\} dv = 0,$$

which gives

$$\int_M \left(\text{Ric}(\xi, \xi) - \lambda(n-1)\|\xi\|^2 \right) + (\|B\|^2 - nF^2) + \left((n-1) \left(\lambda\|\xi\|^2 - n(1+F)^2 \right) \right) = 0.$$

Using $\text{Ric}(\xi, \xi) \geq (n-1)\lambda\|\xi\|^2$, the Schwarz inequality $\|B\|^2 \geq nF^2$ and the condition in the statement $\lambda\|\xi\|^2 \geq n(1+F)^2$ in the above equation, we get the equality $\|B\|^2 = nF^2$, which holds if and only if $B = FI$. Thus

$$\nabla_X \xi = BX + X = FX + X = (1+F)X = \rho X,$$

where $\rho = (1+F)$, that is,

$$\mathcal{L}_\xi g = 2\rho g,$$

which proves that $\xi = \psi^T$ is a conformal vector field. $\square$

In the next result, we consider a conformal vector field on the submanifold M associated with the normal component $\psi^\perp$, and it is interesting to note that in this case we get the criterion for the submanifold to lie on the hypersphere in the Euclidean space, that is, we get a criterion for a spherical submanifold.

Theorem 3.3. *Let $\psi : M \rightarrow R^{n+p}$ be an n -dimensional compact submanifold with mean curvature H . Suppose that the smooth function $f = \frac{1}{2}\|\psi^\perp\|^2$ gives the conformal vector field ∇f on M , and that $\nabla_{\psi^T}^\perp H = 0$. Then $h(\psi^T, \psi^T) = 0$ if and only if $\psi(M) \subset S^{n+p-1}(c)$ for some constant $c > 0$.*

PROOF. Suppose that $h(\psi^T, \psi^T) = 0$. Then, for $\xi = \psi^T$, we have

$$\xi(F) = g(\nabla_\xi^\perp H, \psi^\perp) + g(H, \nabla_\xi^\perp \psi^\perp) = -g(H, h(\xi, \xi)) = 0,$$

that is, $\xi(F) = 0$, which, together with Lemma 3.1, gives

$$\text{Ric}(\xi, \xi) + n\xi(F) + n\rho + nF + \|B\|^2 = 0.$$

Integrating the above equation, we get

$$\int_M \left\{ \text{Ric}(\xi, \xi) + nF + \|B\|^2 \right\} dv = \int_M \left\{ \text{Ric}(\xi, \xi) + \|B\|^2 - n \right\} dv = 0,$$

where we used Lemma 2.1.

Now, using Lemma 3.2 in the above equation, we get

$$\int_M -n^2(1+F)^2 dv = 0,$$

that is, $F = -1$, which, by virtue of Lemma 2.3, gives $\psi(M) \subset S^{n+p-1}(c)$ for some constant $c > 0$.

Conversely, if $\psi(M) \subset S^{n+p-1}(c)$, $c > 0$, then by Lemma 2.3 $F = -1$ and $\psi^T = 0$, and this proves $h(\xi, \xi) = 0$. $\square$

4. Submanifolds with flat normal connection

In this section, we study codimension-two submanifolds in the Euclidean space R^{n+2} with flat normal connection, and find conditions under which the tangential component of the position vector field is a conformal vector field. Let $\psi : M \rightarrow R^{n+2}$ be an immersion of a compact manifold with a flat normal connection and a mean curvature vector field H . We assume that the mean curvature vector field H is nowhere zero, and choose a local orthonormal frame $\{N_1, N_2\}$ of normals such that $H = \alpha N_1$, where $\alpha = \|H\|$. Then, using the definition of the smooth function $F = \langle \psi^\perp, H \rangle$, in this case we have

$$\psi^\perp = \frac{F}{\alpha} N_1 + \mu N_2, \quad \mu = \langle N_2, \psi^\perp \rangle. \quad (4.1)$$

Define a smooth 1-form ω by $\omega(X) = g(\nabla_X^\perp N_1, N_2)$, $X \in \mathfrak{X}(M)$, and let v be the smooth vector field on M dual to ω .

Lemma 4.1. *Let $\psi : M \rightarrow R^{n+2}$ be an immersion of a smooth manifold with a local orthonormal frame $\{N_1, N_2\}$ of normals such that $H = \alpha N_1$. Then, the normal connection on M is flat if and only if ω is closed.*

PROOF. Using $\omega(X) = g(\nabla_X^\perp N_1, N_2)$, we have $\nabla_X^\perp N_1 = \omega(X) N_2$ and that $\nabla_X^\perp N_2 = -\omega(X) N_1$. We compute $R^\perp(X, Y) N_1$ to get

$$R^\perp(X, Y) N_1 = X(\omega(Y)) N_2 - Y(\omega(X)) N_2 - \omega([X, Y]) N_2 = d\omega(X, Y) N_2,$$

and similarly we have

$$R^\perp(X, Y) N_2 = -d\omega(X, Y) N_1, \quad X, Y \in \mathfrak{X}(M),$$

which proves the normal connection is flat if and only if $d\omega = 0$, that is, ω is closed. $\square$

Let M be a submanifold of R^{n+2} with flat normal connection. Then as the smooth 1-form ω , which is dual to smooth vector field v , is closed using equation (2.6), we have a symmetric tensor field C that is given by $\nabla_X v = CX$, for $X \in \mathfrak{X}(M)$.

Lemma 4.2. *Let $\psi : M \rightarrow R^{n+2}$ be an immersion of a smooth manifold with a local orthonormal frame $\{N_1, N_2\}$ of normals such that $H = \alpha N_1$ and shape operators $A_1 = A_{N_1}$ and $A_2 = A_{N_2}$. Then*

$$(i) \quad \sum_{i=1}^n (\nabla A_1)(e_i, e_i) = n\nabla\alpha + A_2v,$$

$$(ii) \sum_{i=1}^n (\nabla A_2)(e_i, e_i) = n\alpha v - A_1 v,$$

where $\{e_1, \dots, e_n\}$ is a local orthonormal frame on M .

PROOF. Using the expression

$$(Dh)(X, Y, Z) = \nabla_X^\perp h(Y, Z) = h(\nabla_X Y, Z) - h(\nabla_X Z, Y),$$

and the Codazzi equation of the submanifold

$$(Dh)(X, Y, Z) = (Dh)(Y, Z, X), \quad X, Y, Z \in \mathfrak{X}(M),$$

we get

$$(\nabla A_1)(X, Y) - (\nabla A_1)(Y, X) = A_{\nabla_X^\perp N_1} Y - A_{\nabla_Y^\perp N_1} X, \quad (4.2)$$

and that

$$(\nabla A_2)(X, Y) - (\nabla A_2)(Y, X) = A_{\nabla_X^\perp N_2} Y - A_{\nabla_Y^\perp N_2} X. \quad (4.3)$$

Also we have

$$\text{Tr } A_1 = n\alpha \quad \text{and} \quad \text{Tr } A_2 = 0, \quad (4.4)$$

and consequently, we get

$$\sum_{i=1}^n g((\nabla A_1)(X, e_i), e_i) = \sum_{i=1}^n g(\nabla_X A_1 e_i, e_i) - g(A_1 \nabla_X e_i, e_i) = ng(X, \nabla \alpha).$$

Using equations (4.2) and (4.3) in the above equation, we arrive at the desired result in (i).

Similarly, using equations (4.3) and (4.4), we get

$$\sum_{i=1}^n g((\nabla A_2)(X, e_i), e_i) = \sum_{i=1}^n g(\nabla_X A_2 e_i, e_i) - g(A_2 \nabla_X e_i, e_i) = X(\text{Tr } A_2) = 0,$$

and arrive at the desired result in (ii). $\square$

In the following main result of this section, we find necessary conditions for the vector field $\xi = \psi^T$ on the submanifold M of the Euclidean space R^{n+2} with flat normal connection to be a conformal vector field. Let $\psi : M \rightarrow R^{n+2}$ be a compact submanifold with flat normal connection, and v be the vector field dual to the closed 1-form ω given in Lemma 4.1, and $h = \text{Tr } C$, C being the symmetric tensor field given by $CX = \nabla_X v$.

Theorem 4.1. *Let $\psi : M \rightarrow R^{n+2}$ be an immersion of a compact manifold with a flat normal connection, and $\{N_1, N_2\}$ a local orthonormal frame of normals such that $H = \alpha N_1$, $H(p) \neq 0$, $p \in M$. If there is a constant c and the following conditions hold:*

- (i) $\text{Ric}(v, v) \geq \frac{n-1}{n}h^2$,
- (ii) $\text{Ric}(\xi - cv, \xi - cv) \geq 0$,
- (iii) $|ch - nF| \leq n$,

where $\xi = \psi^T$, then ξ is a conformal vector field.

PROOF. Using the definition of the curvature tensor field and

$$\nabla_X v = CX, \quad (4.5)$$

we get

$$R(X, Y)v = (\nabla C)(X, Y) - (\nabla C)(Y, X). \quad (4.6)$$

Since $h = \text{Tr } C$, the above equation gives

$$\text{Ric}(X.v) = g\left(\sum_{i=1}^n (\nabla C)(e_i, e_i) - \nabla h, X\right),$$

that is,

$$Q(v) = \sum_{i=1}^n (\nabla C)(e_i, e_i) - \nabla h. \quad (4.7)$$

Now, using equation (4.7) in computing $\text{div } Cv$, we get

$$\text{div } Cv = \text{Ric}(v, v) + v(h) + \|C\|^2. \quad (4.8)$$

Also, equation (4.5) gives $\text{div } v = h$, and thus we have

$$\text{div } hv = v(h) + h^2,$$

which on integration gives

$$\int_M v(h) dv = - \int_M h^2 dv.$$

Now, integrating equation (4.8) and using the above equation, we get

$$\int_M \left\{ \text{Ric}(v, v) + \|C\|^2 - h^2 \right\} dv = 0,$$

that is,

$$\int_M \left\{ \left(\text{Ric}(v, v) - \frac{n-1}{n} h^2 \right) + \left(\|C\|^2 - \frac{1}{n} h^2 \right) \right\} dv = 0.$$

Thus the condition (i) in the statement, together with Schwarz inequality $\|C\|^2 \geq \frac{1}{n} h^2$, gives

$$\text{Ric}(v, v) = \frac{n-1}{n} h^2 \quad \text{and} \quad \|C\|^2 = \frac{1}{n} h^2. \quad (4.9)$$

The second equation in (4.9) gives

$$C = \frac{h}{n} I \quad \text{and} \quad \nabla_X v = \frac{h}{n} X. \quad (4.10)$$

Now, using equation (4.7), we get

$$\text{Ric}(v, v) = - \left(\frac{n-1}{n} \right) v(h),$$

which, together with equation (4.9), gives $v(h) = -h^2$. Also, the first equation in (4.10) and $\text{Tr } B = F$ give $\text{Tr } CB = hF$.

Using equation (4.1) in (2.4), we get

$$X \left(\frac{F}{\alpha} \right) N_1 + \frac{F}{\alpha} \nabla_X^\perp N_1 + X(\mu) N_2 + \mu \nabla_X^\perp N_2 = -h(X, \xi), \quad (4.11)$$

which, taking inner product with N_1 , gives

$$\nabla \left(\frac{F}{\alpha} \right) = \mu v - A_1 \xi, \quad (4.12)$$

similarly, taking inner product with N_2 gives

$$\nabla \mu = -A_2 \xi - \frac{F}{\alpha} v. \quad (4.13)$$

Now, we compute the divergence of the vector field Bv ,

$$\text{div } Bv = \sum_{i=1}^n g(\nabla_{e_i} Bv, e_i) = \sum_{i=1}^n g \left(\nabla_{e_i} \left(\frac{F}{\alpha} A_1 v + \mu A_2 v \right), e_i \right),$$

which, using equations (4.4), (4.12), (4.13) and Lemma 4.1, gives

$$\text{div } Bv = -g(A_1^2 v + A_2^2 v, \xi) + n \frac{F}{\alpha} v(\alpha) + n \alpha \mu \|v\|^2 + Fh. \quad (4.14)$$

On using equation (4.12), we get

$$v(F) = \frac{F}{\alpha} v(\alpha) + \alpha \mu \|v\|^2 - \alpha g(A_1 v, \xi), \quad (4.15)$$

and

$$\operatorname{div} Fv = F \operatorname{div} v + v(F) = hF + \frac{F}{\alpha} v(\alpha) + \alpha \mu \|v\|^2 - \alpha g(A_1 v, \xi),$$

and consequently, that

$$n \operatorname{div} Fv + n \alpha g(A_1 v, \xi) = n \frac{F}{\alpha} v(\alpha) + n \alpha \mu \|v\|^2 + nhF. \quad (4.16)$$

Now, using the expression for the Ricci tensor of submanifold, we have

$$\operatorname{Ric}(X, v) = ng(h(v, X), H) - \sum_{i=1}^n g(h(X, e_i), h(v, e_i)),$$

which gives

$$Q(v) = n \alpha A_1 v - A_1^2 v - A_2^2 v. \quad (4.17)$$

Using equations (4.16) and (4.17) in equation (4.14), we get

$$\operatorname{div} Bv = \operatorname{Ric}(\xi, v) + n \operatorname{div} Fv - (n-1)hF,$$

and integrating the above equation we have

$$\int_M \{\operatorname{Ric}(\xi, v) - (n-1)hF\} dv = 0. \quad (4.18)$$

Finally, using equations (4.9) and (4.18) and Lemma 3.2, we get

$$\int_M \operatorname{Ric}(\xi - cv, \xi - cv) dv = \int_M \left\{ (nF^2 - \|B\|^2) + \frac{n-1}{n} [(ch - nF)^2 - n^2] \right\} dv,$$

which, together with the conditions in the statement and the Schwarz inequality $\|B\|^2 \geq nF^2$, gives

$$\|B\|^2 = nF^2, \quad \xi = cv \quad \text{and} \quad |ch - nF| = n.$$

The second equation, together with equation (4.10), gives

$$\nabla_X \xi = \frac{c}{n} hX, \quad X \in \mathfrak{X}(M).$$

This proves that

$$(\mathcal{L}_\xi g)(X, Y) = 2 \frac{c}{n} hg(X, Y),$$

that is, $\xi = \psi^T$ is a conformal vector field. □

References

- [ALO07] H. ALODAN and S. DESHMUKH, Spherical submanifolds in a Euclidean space, *Monatsh. Math.* **152** (2007), 1–11.
- [ALO02] H. ALODAN and S. DESHMUKH, Spherical submanifolds of a Euclidean space, *Q. J. Math.* **53** (2002), 249–256.
- [ALOD02] H. ALODAN and S. DESHMUKH, Submanifolds with parallel mean curvature vector in a real space form, *Int. Math. J.* **2** (2002), 85–100.
- [DES12] S. DESHMUKH, Conformal vector fields and Eigenvectors of Laplacian operator, *Math. Phys. Anal. Geom.* **15** (2012), 163–172.
- [DES08] S. DESHMUKH and F. AL-SOLAMY, Conformal gradient vector fields on a compact Riemannian manifold, *Colloq. Math.* **112** (2008), 157–161.
- [DES10] S. DESHMUKH, Characterizing spheres by conformal vector fields, *Ann. Univ. Ferrara Sez. VII Sci. Mat.* **56** (2010), 231–236.
- [CHE83] B. Y. CHEN, Total Mean Curvature and Submanifolds of Finite Type, *World Scientific, Singapore*, 1984.
- [OBA62] M. OBATA, Certain conditions for a Riemannian manifold to be isometric with a sphere, *J. Math. Soc. Japan* **14** (1962), 333–340.

HANAN ALOHALI
 DEPARTMENT OF MATHEMATICS
 COLLEGE OF SCIENCE
 KING SAUD UNIVERSITY
 P. O. BOX-2455
 RIYADH-11451
 SAUDI ARABIA
E-mail: halohali@ksu.edu.sa

HAILA ALODAN
 DEPARTMENT OF MATHEMATICS
 COLLEGE OF SCIENCE
 KING SAUD UNIVERSITY
 P. O. BOX-2455
 RIYADH-11451
 SAUDI ARABIA
E-mail: halodan1@ksu.edu.sa

SHARIEF DESHMUKH
 DEPARTMENT OF MATHEMATICS
 COLLEGE OF SCIENCE
 KING SAUD UNIVERSITY
 P. O. BOX-2455
 RIYADH-11451
 SAUDI ARABIA
E-mail: shariefd@ksu.edu.sa

(Received August 10, 2016; revised September 19, 2016)