

Functional equations in the spectral theory of random fields III.

By K. LAJKÓ (Debrecen)

1. Introduction

Let $X(t)$ denote a wide-sense homogeneous random field on the n -dimensional Euclidean space $\mathbb{R}^n$. This means that $M|X(t)|^2 < +\infty$ and $MX(t)\overline{X(s)}$ depends only on the difference $t - s$.

Let $SO(n)$ denote the group of rotations of $\mathbb{R}^n$ around the origin. A homogeneous random field $X(t)$ is called isotropic if

$$MX(t)\overline{X(s)} = MX(gt)\overline{X(gs)},$$

for every $g \in SO(n)$.

Let $\mathcal{M}$ denote a set of sufficiently smooth Jordan surfaces. Each such surface ∂D divides $\mathbb{R}^n$ into two parts: D^- -the interior of ∂D (“past”) and D^+ - the exterior of ∂D (“future”).

The random field $X(t)$ is of Markov type relative $\mathcal{M}$ if for arbitrary ∂D from $\mathcal{M}$ and for arbitrary $t_1 \in D^-$, $t_2 \in D^+$ the random variables $X(t_1)$, $X(t_2)$ are conditionally independent given $\{X(t), t \in \partial D\}$.

Consider the case $n = 1$. The Markov random field discussed above represents an analog of a real Gaussian stationary process $X(t)$ with the following property: for any interval (a, b) and any $t_1 \in (a, b)$, $t_2 \notin (a, b)$ the random variables $X(t_1)$ and $X(t_2)$ are conditionally independent given $X(a)$ and $X(b)$.

The correlation function of such process satisfies the functional equation

$$(1) \quad B(t)[1 + B(2a)] = B(a)[B(t + a) + B(t - a)] \quad (t \geq a, t, a \in \mathbb{R}),$$

(see [2]).

This paper presents the general continuous solution of functional equation (1).

2. Continuous solutions of (1)

Lemma 1. *If the continuous function $B : \mathbb{R} \rightarrow \mathbb{R}$ satisfies the functional equation (1) then $B(t) = 0$ ($t \in \mathbb{R}$) or there exists a positive real number r , such that the functions B_e and B_o defined by*

$$(2) \quad B_e(t) = \frac{B(t) + B(-t)}{2}, \quad B_o(t) = \frac{B(t) - B(-t)}{2} \quad (t \in [-2r, 2r]),$$

satisfy the functional equations

$$(3) \quad B_e(t+a) + B_e(t-a) = 2B_e(t)B_e(a), \quad (t, a) \in T,$$

and

$$(4) \quad B_o(t+a) + B_o(t-a) = 2B_o(t)B_o(a), \quad (t, a) \in T,$$

where $T = \{(t, a) | t \leq |a| \leq r\}$.

PROOF. a) If $B(0) \neq 1$, then, by substitution $t = a$, it follows from (1) the equation

$$B(a)[1 + B(2a)] = B(a)[B(2a) + B(0)] \quad (a \in \mathbb{R}),$$

or

$$B(a)(1 - B(0)) = 0 \quad (a \in \mathbb{R}),$$

which implies

$$(5) \quad B(a) = 0 \quad (a \in \mathbb{R})$$

b) If $B(0) = 1$, then, first by $t = 0$ in (1), we get

$$(6) \quad 1 + B(2a) = B(a)[B(a) + B(-a)] \quad (a \leq 0).$$

Combining (6) with (1), we get the functional equation

$$(7) \quad B(a)[B(t+a) + B(t-a) - B(t)(B(a) + B(-a))] = 0 \quad (t \geq a, a \leq 0).$$

Let us write here $-a$ for a , then we see that B satisfies the functional equation

$$(8) \quad B(-a)[B(t+a) + B(t-a) - B(t)(B(a) + B(-a))] = 0 \quad (t \geq -a, a \geq 0).$$

$B(0) = 1$ and the continuity of B gives that there exists an $r > 0$, such that $B(a) > 0$ if $a \in [-r, r]$ and thus it follows from (7) and (8) that

$$(9) \quad \begin{aligned} B(t+a) + B(t-a) &= B(t)[B(a) + B(-a)], \\ (t, a) &\in T_1 = \{(t, a) \mid t \geq a \text{ if } -r \leq a \leq 0 \text{ or } t \geq -a \text{ if } 0 \leq a \leq r\}. \end{aligned}$$

Putting $t \rightarrow -t$, $a \rightarrow -a$ in (9), we get

$$(10) \quad \begin{aligned} B(-(t+a)) + B(-(t-a)) &= B(-t)[B(a) + B(-a)], \\ (t, a) &\in T_2 = \{(t, a) \mid t \leq -a \text{ if } -r \leq a \leq 0 \text{ or } t \leq a \text{ if } 0 \leq a \leq r\}. \end{aligned}$$

Adding and subtracting equations (9) and (10), we find equations (3) and (4), respectively for functions B_e and B_o defined by (2), where $T = T_1 \cap T_2$.

Lemma 2. *If the continuous function B_e , defined by (2), satisfies the functional equation (3), then in case $B_e(0) = 1$*

$$(11) \quad B_e(t) = \cos bt, \quad t \in [-2r, 2r],$$

$$(12) \quad B_e(t) = \operatorname{ch} bt, \quad t \in [-2r, 2r],$$

$$(13) \quad B_e(t) = 1, \quad t \in [-2r, 2r],$$

where $b \in \mathbb{R}_0 = \mathbb{R} \setminus \{0\}$ is an arbitrary constant.

PROOF. (2) shows that B_e is even and $B_e(t) > 0$ if $t \in [-r, r]$.

Following the arguments of [1] in Chapter 8 for D'Alembert's functional equation, we distinguish two cases

$$B_e(r) \leq 1, \quad \text{or} \quad B_e(r) > 1.$$

If $0 < B_e(r) \leq 1$, then there exists an angle $0 \leq \alpha < \frac{\pi}{2}$, for which

$$(14) \quad B_e(r) = \cos \alpha.$$

With $t = a = \frac{x}{2}$ and taking $B_e(0) = 1$ into consideration, from (3) we obtain

$$(15) \quad B_e(x) + 1 = 2B_e^2\left(\frac{x}{2}\right) \quad (x \in [-2r, 2r])$$

Putting $x = r$ here, we get with (14)

$$(16) \quad B_e\left(\frac{r}{2}\right) = \sqrt{\frac{1 + B_e(r)}{2}} = \sqrt{\frac{1 + \cos \alpha}{2}} = \cos \frac{\alpha}{2}.$$

Then, by induction on n we obtain, that

$$(17) \quad B_e\left(\frac{r}{2^n}\right) = \cos \frac{\alpha}{2^n} \quad \text{for all } n = 0, 1, 2, \dots$$

Now substitute

$$t = \frac{r}{2^n}, \quad a = m \frac{r}{2^n} \quad (m = 1, 2, \dots, m \leq 2^n)$$

into (3), then (using that B_e is even)

$$(18) \quad B_e\left(\frac{m+1}{2^n}r\right) + B_e\left(\frac{m-1}{2^n}r\right) = 2B_e\left(\frac{1}{2^n}r\right) B\left(\frac{m}{2^n}r\right).$$

We prove by induction on k , that

$$(19) \quad B_e\left(\frac{k}{2^n}r\right) = \cos\left(\frac{k}{2^n}\alpha\right) \quad \text{for all } k = 0, 1, 2, \dots, n = 0, 1, 2, \dots$$

Indeed, (19) is true for $k = 0$ and for $k = 1$ by $B_e(0) = 1 = \cos 0$ and (17). Suppose that it holds for $k = m - 1$ and $k = m$. Then by (18)

$$\begin{aligned} B_e\left(\frac{m+1}{2^n}r\right) &= 2B_e\left(\frac{1}{2^n}r\right) B_e\left(\frac{m}{2^n}r\right) - B_e\left(\frac{m-1}{2^n}r\right) \\ &= 2 \cos \frac{1}{2^n}\alpha \cos \frac{m}{2^n}\alpha - \cos\left(\frac{m-1}{2^n}\alpha\right) = \cos\left(\frac{m+1}{2^n}\alpha\right). \end{aligned}$$

and (19) is proved.

For all nonnegative dyadic fractions δ , with $\delta \leq 2$ (i.e., $\delta = \frac{m}{2^n}$, where m and n are nonnegative integers)

$$(20) \quad B_e(\delta r) = \cos \delta \alpha.$$

Since both B_e and $\cos$ are continuous on $[0, 2r]$, by putting into (20) a sequence $\{\delta_m\}$ of dyadic fractions tending to the arbitrary positive number $x \in [0, 2]$, we get $B_e(rx) = \cos \alpha x$, or with $\frac{\alpha}{r} = b$, $rx = t$ $B_e(t) = \cos bt$ for all $t \in [0, 2r]$. Finally, $B_e(-t) = B_e(t)$ gives that

$$B_e(t) = \cos bt, \quad t \in [-2r, 2r],$$

i.e., B_e is of the form (11).

In case $B_e(r) > 1$ we get similarly, that

$$B_e(t) = \operatorname{ch} bt, \quad t \in [-2r, 2r].$$

If $b = 0$ in (11) or (12), then we obtain

$$B_e(t) = 1, \quad t \in [-2r, 2r],$$

and the proof is complete.

Lemma 3. *If the odd and continuous function B_o , defined by (2), satisfies the functional equation (4), where B_e is one of the forms (11), (12), (13), then*

$$(21) \quad B_o(t) = C \sin bt, \quad t \in [-2r, 2r],$$

$$(22) \quad B_o(t) = C \operatorname{sh} bt, \quad t \in [-2r, 2r],$$

$$(23) \quad B_o(t) = Ct, \quad t \in [-2r, 2r],$$

respectively, where $C, b \in \mathbb{R}$ are arbitrary constants.

PROOF. Since $B_e(t) > 0$ if $t \in [-r, r]$ and B_e is continuous thus $\int_0^r B_e(a) da = c \neq 0$.

Integrate (4) from 0 to r with respect to a to obtain

$$\int_0^r B_o(t+a) da + \int_0^r B_o(t-a) da = 2cB_o(t), \quad t \in [-r, r],$$

or

$$\int_t^{t+r} B_o(x) dx - \int_t^{t-r} B_o(x) dx = 2cB_o(t), \quad t \in [-r, r].$$

Thus

$$(24) \quad B_o(t) = \frac{1}{2c} \left[\int_t^{t+r} B_o(x) dx - \int_t^{t-r} B_o(x) dx \right], \quad t \in [-r, r].$$

Since B_o is continuous, the right hand side of (24) is differentiable on $[-r, r]$ and so B_o is differentiable on $[-r, r]$ too.

But with $t = a = \frac{x}{2}$ and taking $B_o(0) = 0$ in consideration, we obtain from (4)

$$(25) \quad B_o(x) = 2B_o\left(\frac{x}{2}\right) B_e\left(\frac{x}{2}\right), \quad x \in [-2r, 2r],$$

which implies that B_o is differentiable on $[-2r, 2r]$.

By differentiating (24), we obtain

$$(26) \quad B'_o(t) = \frac{1}{2c} [B(t+r) - B(t-r)], \quad t \in [-r, r].$$

The right hand side of (26) is differentiable on $[-r, r]$ and so B_o is twice differentiable on $[-r, r]$. Finally, (25) gives the differentiability of B_o on the interval $[-2r, 2r]$.

We differentiate (4) twice with respect to a and substitute $a = 0$, then we get:

a) in case (11),

$$B''_o + b^2 B_o(t) = 0, \quad t \in [-2r, 2r].$$

Then $B_o(t) = C \sin bt + C' \cos bt$ with some constants $C, C' \in \mathbb{R}$. Finally, because of B_o is an odd function, we get (21) for B_o .

b) in case (12),

$$B_o''(t) - b^2 B_o(t) = 0, \quad t \in [-2r, 2r],$$

which implies that

$$B_o(t) = C_1 e^{bt} + C_2 e^{-bt}, \quad t \in [-2r, 2r],$$

where $C_1, C_2 \in \mathbb{R}$ are arbitrary constants. But B_o is odd, therefore we get that $C_2 = -C_1$ and so

$$B_o(t) = C_1(e^{bt} - e^{-bt}), \quad t \in [-2r, 2r],$$

i.e., (22) holds with $C = \frac{C_1}{2}$.

c) in case (13),

$$B_o''(t) = 0, \quad t \in [-2r, 2r],$$

which implies (23) for B_o .

Theorem. *The general continuous solution $B : \mathbb{R} \rightarrow \mathbb{R}$ of functional equation (1) is given by*

$$(27) \quad B(t) \equiv 0 \quad (t \in \mathbb{R})$$

$$(28) \quad B(t) = \cos bt + C \sin bt \quad (t \in \mathbb{R})$$

$$(29) \quad B(t) = \operatorname{ch} bt + C \operatorname{sh} bt \quad (t \in \mathbb{R})$$

$$(30) \quad B(t) = 1 + Ct \quad (t \in \mathbb{R})$$

where $b, C \in \mathbb{R}$ are arbitrary constants.

PROOF. Lemma 1 shows that $B(t) \equiv 0$, if $B(0) \neq 1$. If $B(0) = 1$, then we obtain again from Lemma 1 that there exists an $r > 0$, such that functions B_e and B_o , defined by (2), satisfy the functional equations (3) and (4) and because of (2)

$$(31) \quad B(t) = B_e(t) + B_o(t) \quad t \in [-2r, 2r].$$

Using (11), (12), (13), (21), (22), (23), we have (28), (29), (30) for B , if $t \in [-2r, 2r]$. From equation (1), using some well known identities for trigonometric and hyperbolic functions, we get the following equalities:

$$(\cos bt + C \sin bt)[1 + B(2a)] = (\cos bt + C \sin bt)(1 + \cos b2a + C \sin b2a),$$

$$(\operatorname{ch} bt + C \operatorname{sh} bt)[1 + B(2a)] = (\operatorname{ch} bt + C \operatorname{sh} bt)(1 \operatorname{ch} b2a + C \operatorname{sh} b2a),$$

$$(1 + ct)[1 + B(2a)] = (1 + ct)(1 + 1 + c2a)$$

which imply that B has the forms (28), (29), (30) for all $t \in [-4r, 4r]$, respectively and than by induction on n , we get these forms for all $t \in [-2^n r, 2^n r]$ ($n \in N$) and finally for all $t \in \mathbb{R}$.

References

- [1] J. ACZÉL and J. DHOMBRES, Functional Equations in Several Variables, *Cambridge University Press*, 1989.
- [2] M. I. YADRENKO, Spectral Theory of Random Fields, Optimization software, *INC, Publications Division, New York*, 1983.

KÁROLY LAJKÓ
LAJOS KOSSUTH UNIVERSITY,
INSTITUTE OF MATHEMATICS AND INFORMATICS,
4010 DEBRECEN P.O.B. 12.

(Received March 28, 1994)