

On the determinant of general pentadiagonal matrices

By CARLOS M. DA FONSECA (Safat) and LÁSZLÓ LOSONCZI (Debrecen)

Abstract. In this paper, we consider square matrices with two sub-diagonals and two super-diagonals. We provide an algorithm to transform such matrices (by multiplying them with suitable matrices) to tridiagonal matrices. It is known that tridiagonal matrices can be transformed to diagonal ones (again by multiplying them from both sides by suitable matrices). Thus pentadiagonal matrices can be transformed to diagonal ones and in this way their determinants can be calculated. Two examples show how our method works.

CARLOS M. DA FONSECA
KUWAIT COLLEGE OF SCIENCE
AND TECHNOLOGY
DOHA DISTRICT, BLOCK 4
P. O. BOX 27235
SAFAT 13133
KUWAIT
AND
UNIVERSITY OF PRIMORSKA
FAMNIT
GLAGOLJSAŠKA 8
6000 KOPER
SLOVENIA

LÁSZLÓ LOSONCZI
FACULTY OF ECONOMICS
UNIVERSITY OF DEBRECEN
DEBRECEN
HUNGARY

Mathematics Subject Classification: 15A15, 15B99, 65F40, 15A42.

Key words and phrases: determinant, pentadiagonal matrices, tridiagonal matrices.