

## Recollements associated to cotorsion pairs over upper triangular matrix rings

By RONGMIN ZHU (Nanjing), YEYANG PENG (Nanjing)  
and NANQING DING (Nanjing)

**Abstract.** Let  $A, B$  be two rings and  $T = \begin{pmatrix} A & M \\ 0 & B \end{pmatrix}$  with  $M$  an  $A$ - $B$ -bimodule. Suppose that we are given two complete hereditary cotorsion pairs  $(\mathcal{A}_A, \mathcal{B}_A)$  and  $(\mathcal{C}_B, \mathcal{D}_B)$  in  $A$ -Mod and  $B$ -Mod, respectively. We define two cotorsion pairs  $(\Phi(\mathcal{A}_A, \mathcal{C}_B), \text{Rep}(\mathcal{B}_A, \mathcal{D}_B))$  and  $(\text{Rep}(\mathcal{A}_A, \mathcal{C}_B), \Psi(\mathcal{B}_A, \mathcal{D}_B))$  in  $T$ -Mod and show that both of these cotorsion pairs are complete and hereditary. If we are given two cofibrantly generated model structures  $\mathcal{M}_A$  and  $\mathcal{M}_B$  on  $A$ -Mod and  $B$ -Mod, respectively, then using the result above, we investigate when there exists a cofibrantly generated model structure  $\mathcal{M}_T$  on  $T$ -Mod and a recollement of  $\text{Ho}(\mathcal{M}_T)$  relative to  $\text{Ho}(\mathcal{M}_A)$  and  $\text{Ho}(\mathcal{M}_B)$ . Finally, some applications are given in Gorenstein homological algebra.

RONGMIN ZHU &  
YEYANG PENG &  
NANQING DING  
DEPARTMENT OF MATHEMATICS  
NANJING UNIVERSITY  
NANJING, 210093  
P. R. CHINA

---

*Mathematics Subject Classification:* 16E30, 18E30, 18G25.

*Key words and phrases:* upper triangular matrix ring, cotorsion pair, model structure, recollement.