

## Functional analysis behind a family of multidimensional continued fractions. Part II

By ILYA AMBURG (Ithaca) and THOMAS GARRITY (Williamstown)

**Abstract.** This paper is a direct continuation of “Functional analysis behind a family of multidimensional continued fractions. Part I”, in which we started the exploration of the functional analysis behind the transfer operators for triangle partition maps, a family that includes many, if not most, well-known multidimensional continued fraction algorithms. This allows us now to find eigenfunctions of eigenvalue 1 for transfer operators associated with select triangle partition maps on specified Banach spaces. We proceed to prove that the transfer operators, viewed as acting on one-dimensional families of Hilbert spaces, associated with select triangle partition maps are nuclear of trace class zero. We finish by deriving Gauss–Kuzmin distributions associated with select triangle partition maps.

ILYA AMBURG  
CENTER FOR APPLIED MATHEMATICS  
CORNELL UNIVERSITY  
ITHACA, NY 14850  
USA

THOMAS GARRITY  
DEPARTMENT OF MATHEMATICS  
WILLIAMS COLLEGE  
WILLIAMSTOWN, MA 01267  
USA

---

*Mathematics Subject Classification:* 11J70, 37C30.

*Key words and phrases:* continued fractions, number theory, triangle partition maps, functional analysis.