

On two problems concerning the theory of binary relations.

By G. FODOR in Szeged.

Let us consider the closed interval $[0, 1]$. Let us suppose that to each point x of this interval there corresponds a set S_x — called the picture of x — the points of which are outside of the interval $K(x)$ which is symmetrical about x ; the length of $K(x)$ shall be denoted by $f(x)$; we shall suppose only that $f(x) > 0$, for every x . Let us call two points of the interval $[0, 1]$ *independent*, if neither of them belongs to the picture of the other.

Problem 1. Does there exist in $[0, 1]$ a set of positive measure, all pairs of points of which are independent?

Problem 2. Does there exist in $[0, 1]$ a set, having the power of the continuum, all pairs of points of which are independent?

Theorem 1. *If $f(x)$ is a measurable function, there exists in $[0, 1]$ a set of positive measure, the pairs of points of which are independent.*

Proof. The sets

$$E_i = E \left[\frac{1}{i} < f(x) \leq \frac{1}{i-1} \right] \quad (i = 2, 3, \dots)$$

cannot all be of measure zero, because in this case the interval $[0, 1]$ would be the sum of countably many sets of measure zero. Hence there exists a value $i = i_1$ for which E_{i_1} is of positive measure. Let us divide the interval $[0, 1]$ into n subintervals, by means of the points $x_0 = 0 < x_1 < x_2 < \dots < x_n = 1$, in such a manner that the length of each subinterval (x_k, x_{k+1}) should be less than $\frac{1}{2i_1}$. Let us now denote the common part of the set E_{i_1} with the subinterval $[x_k, x_{k+1}]$ by $E_{i_1 k}$ ($k = 0, 1, 2, \dots, n-1$). Among the sets $E_{i_1 k}$ there must be at least one — $E_{i_1 k_1}$ say — having positive measure, because the sum of the sets $E_{i_1 k}$ is of positive measure. Any two points of $E_{i_1 k_1}$ are independent, because if x belongs to $E_{i_1 k_1}$ and y to S_x we have $|y - x| > \frac{f(x)}{2} \geq \frac{1}{2i_1}$ and thus y is outside $E_{i_1 k_1}$.

Theorem 2. *The answer to problem 2 is always in the affirmative.¹⁾*

Proof. Let us define the sets E_i as above ($i = 2, 3, \dots$). If all sets E_i were countable, the interval $[0, 1]$ would be the sum of countably many countable sets. Thus there exists a value $i_1 = i$ for which E_{i_1} has the power of the continuum. Let us now define the points x_k and the sets $E_{i_1 k}$ as above ($k = 0, 1, 2, \dots, n-1$). At least one of the sets $E_{i_1 k}$ — say $E_{i_1 k_1}$ — will have the power of the continuum. It follows exactly as above that any two points of $E_{i_1 k_1}$ are independent.

So far we supposed, that the interval $K(x)$ is symmetrical about x . It is easy to see that our results are true and the proofs essentially unchanged, if we suppose only that x is an interior point of the interval $K(x)$, i. e. $K(x)$ is the interval $[x - f_1(x), x + f_2(x)]$ where $f_1(x)$ and $f_2(x)$ are positive functions; as regards Theorem 1 it must be supposed that both $f_1(x)$ and $f_2(x)$ are mesurable. As a matter of fact, if $f(x) = \min(f_1(x), f_2(x))$, it follows that S_x is outside of the symmetrical interval of length $2f(x)$ about x , and the problem is reduced to that considered above.

(Received May 26, 1950)

¹⁾ D. LÁZÁR, On a problem in the theory of aggregates. *Compositio Math.* 3 (1936), 304. In this paper D. LÁZÁR has proved a similar theorem in the case where each S_x has only a finite number of elements. His method could be applied also to our case, but we prefer to give a different proof.