

On the definition of the determinant as a multilinear antisymmetric function.

By S. N. AFRIAT in Jerusalem.

The properties considered here for the identification of the determinant are the three characteristic properties of WEIERSTRASS.¹⁾ The present object is to show how this identification can be very simply carried out, together with an incidental proof of the multiplication theorem.²⁾

Consider a function $\Delta(\mathbf{a}) = \Delta(\mathbf{a}_1, \dots, \mathbf{a}_n)$, of n vectors $\mathbf{a}_1, \dots, \mathbf{a}_n$ of order n , and thus of a matrix $\mathbf{a}$ of order $n \times n$ whose i, j th element a_{ij} is the i th element of the j th vector $\mathbf{a}_j$; and suppose that it has the following properties:

- I. $\Delta(\mathbf{a}_1, \dots, \lambda \mathbf{x}, \dots, \mathbf{a}_n) = \Delta(\mathbf{a}_1, \dots, \mathbf{x}, \dots, \mathbf{a}_n) \lambda$.
- II. $\Delta(\mathbf{a}_1, \dots, \mathbf{x} + \mathbf{y}, \dots, \mathbf{a}_n) = \Delta(\mathbf{a}_1, \dots, \mathbf{x}, \dots, \mathbf{a}_n) + \Delta(\mathbf{a}_1, \dots, \mathbf{y}, \dots, \mathbf{a}_n)$.
- III. $\Delta(\mathbf{a}_1, \dots, \mathbf{x}, \dots, \mathbf{y}, \dots, \mathbf{a}_n) = -\Delta(\mathbf{a}_1, \dots, \mathbf{y}, \dots, \mathbf{x}, \dots, \mathbf{a}_n)$.

As a consequence of III:

$$(1) \quad \Delta(\mathbf{a}_1, \dots, \mathbf{x}, \dots, \mathbf{x}, \dots, \mathbf{a}_n) = 0;$$

and if π is any permutation, by which $i \rightarrow \pi i$ for $i = 1, \dots, n$ and $\pi i \neq \pi j$ ($i \neq j$), then

$$(2) \quad \Delta(\mathbf{a}_{\pi 1}, \dots, \mathbf{a}_{\pi n}) = \alpha(\pi) \Delta(\mathbf{a}_1, \dots, \mathbf{a}_n),$$

where $\alpha(\pi)$ is the alternating character of π .

Let $\mathbf{b}$ be a matrix of order $n \times n$, with elements b_{ij} and determinant $|\mathbf{b}| = \sum_{\pi} \alpha(\pi) b_{\pi 1, 1} \dots b_{\pi n, n}$. Then for $\Delta(\mathbf{a}\mathbf{b})$ there is the expression

$$\Delta(\mathbf{a}_1 b_{11} + \dots + \mathbf{a}_n b_{n1}, \dots, \mathbf{a}_1 b_{1n} + \dots + \mathbf{a}_n b_{nn}).$$

By repeated application of I and II this ultimately becomes

$$\sum_i \Delta(\mathbf{a}_{i_1}, \dots, \mathbf{a}_{i_n}) b_{i_1, 1}, \dots, b_{i_n, n},$$

¹⁾ WEIERSTRASS, Werke III, 271–286. Other references are to be found in MAC DUFFEE [1].

²⁾ GÁSPÁR [2] takes the multiplication theorem as one of a defining set of axioms.

where the indices in the set $i = (i_1, \dots, i_n)$ range independently in $1, \dots, n$. By (1) and (2), certain of the terms in this sum, those for which $i_1, \dots, i_n$ is not a permutation of $1, \dots, n$ (i. e. there exist $i_j = i_k, j \neq k$), vanish, and the rest are contained in the sum

$$\sum_{\pi} \Delta(a_1, \dots, a_n) \alpha(\pi) b_{\pi 1, 1} \dots b_{\pi n, n} = \Delta(a) |b|.$$

Hence I, II and III imply $\Delta(ab) = \Delta(a) |b|$. By taking $a = \mathbf{1}$, this gives $\Delta(b) = \Delta(\mathbf{1}) |b|$; so that the function Δ is shown to be *identical with the determinant, but for a constant multiplier, given by $\Delta(\mathbf{1})$* . If there is taken the further supposition:

IV. $\Delta(\mathbf{1}) = 1$;

then $\Delta(b) = |b|$; and $\Delta(ab) = \Delta(a) |b|$ then shows that $|ab| = |a| |b|$.

Bibliography.

- [1] C. C. MACDUFFEE, The Theory of Matrices, New York, 1946.
- [2] JULIUS GÁSPÁR, Eine neue Definition der Determinanten, *Publ. Math. Debrecen* 3 (1954), 257—260.

(Received November 1, 1955.)