

On some properties of commutator subsemigroups

J. DÉNES (Budapest)

Dedicated to Professor A. G. Kuroš on the occasion of his sixtieth birthday

A. SUSCHKEWITSCH was the first who introduced the graph representation of a transformation of degree n i.e. a mapping of a set of n elements into itself (see [8]). For a certain generalization see [9].

One can formulate A. Suschkewitsch's graph representation as follows: to every transformation of degree n there may correspond uniquely a directed graph having n labelled vertices in such a way that the vertices are labelled by the natural numbers $1, 2, \dots, n$ and if the transformation maps i to j then the graph has a directed edge from i to j (see [8]).

The same graph representation was rediscovered by O. ORE, F. HARARY and others. (See e.g. [4], [6]).

Several properties of the graph representation have been investigated by the author and some of the results were contained in three of the author's papers (see [1], [2], [3]).

It is easy to see that a directed graph corresponds to a transformation if and only if each of its connected components contains a single cyclically directed circuit and directed rooted trees. Such graphs with n vertices will be called $F(n)$ graphs. A transformation corresponding to a component of an $F(n)$ graph is conveniently called *generalized cycle*. Deleting the trees (apart from their roots) from an $F(n)$ graph one can obtain a special $F(k)$ graph ($k \leq n$), containing circuits only. The transformation corresponding to the $F(k)$ graph is a permutation: it is called the *main-permutation* of the original transformation.

If α denotes an arbitrary transformation of degree n , then the *quasiinverse* will be defined so that the quasiinverse of α is its power α^s with the least exponent s whose main permutation is the inverse of the main permutation of α . Obviously if α is a permutation its quasiinverse is equal to its inverse. Therefore the notation (α^{-1}) of the quasiinverse of α will not be troublesome.

For an arbitrary abstract semigroup S of order n there exists a subsemigroup S' of the symmetric semigroup of degree $n+1$ such that S and S' are isomorphic. If the representation is given by the correspondence

$$a_i \leftrightarrow \begin{pmatrix} a_1 & a_2 & \dots & a_n & a_{n+1} \\ a_1 a_i & a_2 a_i & \dots & a_n a_i & a_i \end{pmatrix}$$

where $a_i \in S$, $i=1, 2, \dots, n$ it is called a *regular representation*. By means of the regular representation one can define the quasiinverse of an arbitrary element

of an abstract semigroup in a similar way as it would be for a transformation. Let us consider the abstract semigroup S and its regular representation T ; if $\alpha \in T$ and $a \in S$ then $\alpha \leftrightarrow a$ implies $\alpha^{-1} \leftrightarrow a^{-1}$ and the *commutator subsemigroup* of S will be defined as the subsemigroup which is generated by the *commutators* $aba^{-1}b^{-1}$ where $a, b \in S$. If A_n denotes the alternating group of degree n , S_n denotes the symmetric group of degree n , and the symmetric semigroup of degree n i.e. the semigroup of all transformations of degree n will be denoted by F_n , then for the commutator subsemigroup K_n of F_n the equality

$$K_n = (F_n \setminus S_n) \cup A_n$$

holds, where $\setminus$ denotes the set-theoretical difference. The proof which we omit here has been published in [2].

Since K_n plays a similar role in F_n as A_n in S_n , K_n will be called *alternating semigroup* of degree n .

It is almost trivial, that $\alpha K_n \alpha^{-1} = K_n$ ($\alpha \in F_n$) holds, i.e. K_n is a *normal subsemigroup* of F_n . Further K_n is a maximal normal subsemigroup, since $\{K_n, \alpha\} = F_n$ if $\alpha \notin K_n$ holds.

When a group coincides with its commutator subgroup it is said to be a *perfect group*. It is well-known that A_n is perfect. A semigroup will be called *perfect* if it coincides with its commutator subsemigroup. To exhibit the strong analogy between A_n and K_n we shall prove that K_n is perfect.

Since A_n is perfect and all the transformations of the form

$$\begin{pmatrix} 1 & 2 & \dots & i-1 & i & i+1 & \dots & n \\ 1 & 2 & \dots & i-1 & j & i+1 & \dots & n \end{pmatrix} \quad i \neq j, i, j = 1, 2, \dots, n$$

(they are called *singular transformations*) are idempotent elements and so commutators, there remains to prove that A_n and the set T_n of all singular transformations generate K_n i.e.

$$K_n = \{T_n, A_n\}$$

holds. It is obvious, that an arbitrary element of K_n whose main permutation is even can be represented as the product of the elements of A_n and T_n .

The proof will be completed when it is pointed out that for any $\alpha \in K_n$ whose main permutation is odd $\alpha \in \{T_n, A_n\}$ holds. This is true since if k is odd then the permutation $\varrho = (1 \ 2 \ \dots \ k) \ (k \leq n)$ is contained in A_n and the singular transformation $\sigma = \begin{pmatrix} 1 & 2 & \dots & k-1 & k \\ 1 & 2 & & k-1 & 1 \end{pmatrix}$ is an element of T_n . Obviously $\varrho\sigma$ is a transformation whose main permutation is odd, since $\varrho\sigma = \begin{pmatrix} 1 & 2 & \dots & k-1 & k \\ 2 & 3 & & 1 & 1 \end{pmatrix}$. By iteration the proof can be easily extended to an arbitrary transformation (which is not a permutation) and whose main permutation is odd.

N. ITO [5] and O. ORE [8] proved: if $n \geq 5$ all the elements of A_n are commutators. The author has the conjecture, that a similar theorem holds for the alternating semigroups i.e. all the elements of K_n ($n \geq 5$) are commutators.

The author hereby wishes to express his thanks to Prof. B. N. SCHEIN who kindly called his attention to [9]. The author is indebted to Prof. Š. SCHWARZ for making possible to have a copy of A. Suschkewitsch's book.

References

- [1] J. DÉNES, Connection between transformation semigroups and graphs. *Actes des Journees Internationales d'etude sur la theorie de graphs*. Rome (1966), 298—303.
- [2] J. DÉNES, On transformations, transformation-semigroups and graphs. *Theory of graphs. Proc. of the Colloquium held at Tihany Hungary 1966*, 65—75.
- [3] J. DÉNES, Some combinatorial properties of transformations and their connections with the theory of graphs. *J. Combinatorial Theory (to appear)*.
- [4] F. HARARY, The number of functional diagraphs. *Math. Ann.* **138** (1959), 203—210.
- [5] N. ITO, A theorem on alternating group. *Math. Japon.* **2** (1951), 59—60.
- [6] O. ORE, Graphs and correspondences. *Festschrift zum 60. Geburtstag von Prof. Andreas Speiser' Zürich* (1945), 184—191.
- [7] O. ORE, Some remarks on commutators. *Proc. Amer. Math. Soc.* **2** (1951), 307—314.
- [8] A. SUSCHKEWITSCH, Untersuchungen über verallgemeinerte substitutionen. *Atti del Congresso Internazionale dei Matematici Bologna 1928*, 147—157.
- [9] M. YOELI, The cascade decomposition of sequential machines. *IRE Trans. EC-10* (1961), 587—592.

(Received December 8, 1967.)