

Inverse limits of the Cantor-manifolds

By IVAN LONČAR (Varaždin)

ABSTRACT. We say that a compact space X is (n, k) -Cantor-manifold if $\dim X = n$, and if for every closed subset $F \subseteq X$ of the dimension $\dim F \leq n - k$ the set $X \setminus F$ is connected.

In the present paper we investigate the following question: Under what conditions the limit of the inverse system of the n -dimensional Cantor-manifolds is (n, k) -Cantor-manifold?

The partial answers for the inverse systems with fully closed monotone bonding mappings and for inverse systems of metric Cantor-manifolds with open bonding mappings are given.

0. Introduction

All spaces in this paper are assumed to be Hausdorff, and this assumption will be used without explicit mention.

We say that a compact space X is an $\text{ind}(\text{Ind}, \text{dim})$ - n -dimensional Cantor-manifold if $\text{ind } X (\text{Ind } X, \text{dim } X) = n \geq 1$ such that no closed subset $F \subseteq X$ satisfying the inequality $\text{ind } F (\text{Ind } F, \text{dim } F) \leq n - 2$ separates the space X i.e. for every such set the complement $X \setminus F$ is connected [7; 91].

If we omit the assumption that X is a compact space, we obtain the definition of a generalized Cantor-manifold.

The cardinality of the set A is denoted by $|A|$. The symbol $\text{cf}(A)$ means the cofinality of the well-ordered set A i.e. the smallest ordinal number which is cofinal in A .

If $f: X \rightarrow Y$ is a mapping, then $f^\# A$ is the set $\{y: f^{-1}(y) \subseteq A\}$ for $A \subseteq X$.

1. Inverse limits of k -dimensional cantor-manifolds

A compact space X is (n, k) -Cantor-manifold if $\dim X = n$ and if $X \setminus F$ is connected for each closed $F \subseteq X$ with $\dim F \leq n - k$.

A k -dimensional Cantor-manifold is $(k, 2)$ -Cantor-manifold.

We start with the next theorem

1.1. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of k -dimensional Cantor-manifolds X_α . If $\underline{X}$ satisfies the property:

Mathematics subject classifications (1980): Primary 54M25; Secondary 57NXX., Key words and phrases: inverse systems and limits, Cantor-manifold.

(P1) For every closed $(n-k)$ -dimensional subset $F \subseteq \varprojlim X$ the set $f_\alpha(F)$, $\alpha \in A$, is $\leq (k-2)$ -dimensional,

(P2) For every open connected $Y_\alpha \subseteq X_\alpha$, $\alpha \in A$, the set $f_\alpha^{-1}(Y_\alpha)$ is connected, then $X = \varprojlim X$ is (n, k) -Cantor-manifold.

PROOF. Let F be a closed subset of X of the dimension $\leq n-k$. By (P1) we have $\dim f_\alpha(F) \leq k-2$. This means that $Y_\alpha = X_\alpha \setminus f_\alpha(F)$ is open and connected. From (P2) it follows that $f_\alpha^{-1}(Y_\alpha)$ is connected. Since $X \setminus F = \bigcup \{Y_\alpha : \alpha \in A\}$ it follows that $X \setminus F$ is connected [6: 435.]. Q.E.D.

1.2. Remark. The property (P2) is satisfied if $f_{\alpha\beta}$ are monotone or open mappings. This follows from [6: 6.1.28. Theorem] and from the fact that if X is an inverse system of connected spaces and open-closed projections, then $\varprojlim X$ is connected.

We say that a mapping $f: X \rightarrow Y$ is fully closed [8] if for every point $y \in Y$ and for each finite cover $\{U_1, U_2, \dots, U_n\}$ of $f^{-1}(y)$ the set $\{y\} \cup (f^\# U_1 \cup \dots \cup f^\# U_n)$ is open [8].

If $f: X \rightarrow Y$ is fully closed then f is closed. If Y has no isolated points and $f: X \rightarrow Y$ is open fully closed, then f is a homeomorphism [8].

1.3. Lemma. [8: Lemma 1.]. If $f: X \rightarrow Y$ is compact mapping of regular space X , then f is fully closed iff f is closed and for each pair F_1, F_2 of disjoint closed subset of X the set $f(F_1) \cap f(F_2)$ is discrete.

From this Lemma it follows

1.4. Lemma. If $f: X \rightarrow Y$ is a fully closed compact mapping of T_3 space X , then $f|F: F \rightarrow f(F)$ is fully closed for every closed subset F of X .

1.5. Lemma. [8] Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system such that $f_{\alpha\beta}$ are fully closed perfect mappings. The projection $f_\alpha: \varprojlim X \rightarrow X_\alpha$ are fully closed iff $f_{\alpha\beta}$ are fully closed.

1.6. Lemma. [8: Teorema 4.]. If $f: X \rightarrow Y$ is fully closed surjection between normal spaces, then $\dim X \leq \dim Y + 1$.

From the preceding Lemmas it follows

1.7. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of the n -dimensional Cantor-manifolds X_α . If the mappings $f_{\alpha\beta}$ are monotone fully closed, then $X = \varprojlim X$ is a $(n, 3)$ -Cantor-manifold.

PROOF. It is known that $\dim X \leq n$. From lemmas 1.6. and 1.5. it follows that $\dim X \leq n-1$. This means that either $\dim X = n$ or $\dim X = n-1$. Consider a closed $F \subseteq X$ with $\dim F \leq n-3$. From Lemmas 1.6. and 1.4. it follows that $\dim f_\alpha(F) \leq \dim F + 1 \leq n-2$. Since X_α is the Cantor-manifold we infer that $X_\alpha \setminus f_\alpha(F)$ is connected. The set $f_\alpha^{-1}(X_\alpha \setminus f_\alpha(F))$ is also connected since f_α are monotone mappings. The properties (P1) and (P2) of Theorem 1.1 are satisfied. The proof is completed.

1.8. Remark. If $\dim X = n-1$, then X is the $(n-1)$ -dimensional Cantor-manifold.

Now we pass to the inverse systems of a generalized Cantor-manifolds.

1.9. Lemma. Let $\underline{X} = \{X_n, f_{nm}, N\}$ be an inverse sequence of the countably compact spaces X_n . If the mappings f_{nm} are fully closed, then the projections $f_n: X = \varprojlim X \rightarrow X_n, n \in N$, are fully closed.

PROOF. Let x_n be a point of X_n and $\{U_1, \dots, U_k\}$ an open cover of the set $f_n^{-1}(x_n)$. We consider the family $\{f_m^\# U_1, \dots, f_m^\# U_k\}, m \geq n$. If the set $Y_m = f_{nm}^{-1}(x_n) \setminus (f_m^\# U_1 \cup \dots \cup f_m^\# U_k)$ is non-empty for every $m \geq n$, then we obtain the inverse system $Y = \{Y_m, f_{pm}/Y_m, n \leq p \leq m\}$ which has a non-empty limit [18]. This is impossible since $f_n^{-1}(x_n) \subseteq (U_1 \cup \dots \cup U_k)$ and $U_i = \bigcup \{f_m^{-1} f_m^\# U_i: m \geq n\}$. Hence, there exist $m_0 \geq n$ such that $Y_{m_0} = \emptyset$ i.e. $f_{nm_0}^{-1}(x_n) \subseteq f_{m_0}^\# U_1 \cup \dots \cup f_{m_0}^\# U_k$. Since f_{nm_0} is fully closed, the set $\{x_n\} \cup (f_{nm_0}^\# f_{m_0}^\# U_1 \cup \dots \cup f_{nm_0}^\# f_{m_0}^\# U_k)$ is open. The proof is completed since

$$f_{nm_0}^\# f_{m_0}^\# U_i = f_n^\# U_i \text{ for every } i \in \{1, \dots, k\}.$$

By the similar method of proof we have

1.10. Lemma. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, \omega_\tau\}$ be an inverse system of $\aleph_\tau$ -compact spaces X_α . If the mappings $f_{\alpha\beta}$ are fully closed, the projections $f_\alpha: \varprojlim X \rightarrow X_\alpha, \alpha \in A$, are fully closed.

1.11. Theorem. Let $\underline{X} = \{X_n, f_{nm}, N\}$ be an inverse sequence of the k -dimensional normal countably compact generalized Cantor-manifolds X_n . If the mappings f_{nm} are monotone fully closed surjections, then $X = \varprojlim X$ is $(k, 3)$ -Cantor-manifold.

PROOF. It suffices in the proof of Theorem 1.7. to apply the fact that $f_n: X \rightarrow X_n, n \in N$, are monotone mappings and X is connected [18].

Similarly, on can prove

1.12. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, \omega_\tau\}$ be an inverse system of the k -dimensional normal $\aleph_\tau$ -compact generalized Cantor-manifolds X_α . If the mappings $f_{\alpha\beta}$ are monotone fully closed surjections, then $X = \varprojlim X$ is $(k, 3)$ -Cantor-manifold.

1.13. Remark. If $\dim X = k - 1$, then X is $(k - 1)$ -dimensional generalized Cantor-manifold.

1.14. Remark. The space X in Theorem 1.11. (in Theorem 1.12.) is normal countably compact (normal $\aleph_\tau$ -compact) [18].

The next part of this Section is devoted to inverse systems of metric Cantor-manifold.

1.15. Lemma. If $f: X \rightarrow Y$ is an open-closed surjection between separable metric spaces such that for every $y \in Y$ the fibre $f^{-1}(y)$ is a discrete subspace of X , then $\text{ind } Z = \text{ind } f(Z)$ for every closed subset $Z \subseteq X$.

PROOF. Let $B = \{U_i: i \in N\}$ be a base for X and let A_i be defined as in the proof of 1.12.5. Lemma in [7]. We have $X = \bigcup \{A_i: i \in N\}$ (see the proof of 1.12.7. Lemma in [7]). The set $A_i \cap Z$ is F_σ relative to Z and $f(A_i \cap Z)$ is F_σ relative to $f(Z)$. Furthermore, Z is separable metric space and $f|_{A_i \cap Z}: A_i \cap Z \rightarrow f(A_i \cap Z)$ is a homeomorphism. This means that $\text{ind}(A_i \cap Z) = \text{ind } f(A_i \cap Z) \leq \text{ind } f(Z)$ and $\text{ind } Z \leq$

$\cong \text{ind}(A_i \cap Z) = \text{ind} f(A_i \cap Z)$. From this relations and the relations $Z = \bigcup \{A_i \cap Z : i \in N\}$, $f(Z) = \bigcup \{f(A_i \cap Z) : i \in N\}$ it follows that $\text{ind} Z = \text{ind} f(Z)$. Q.E.D.

1.16. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of k -dimensional generalized metric Cantor-manifolds X_α and open—closed surjections $f_{\alpha\beta}$. The space $X = \varprojlim X$ is k -dimensional generalized Cantor-manifold if the following conditions are satisfied:

(C1) X is a metric space,

(C2) For every $x_\alpha \in X_\alpha$, $\alpha \in A$, the fiber $f_\alpha^{-1}(x_\alpha)$ is a discrete subspace of X .

PROOF. X is separable metric space of the dimension $\dim X = k$ (Lemma 1.15.). For every $F \subseteq X$ of the dimension $\dim F \leq k-2$ it follows that $\dim f_\alpha(F) = \dim F \leq k-2$ (Lemma 1.15.). Hence, condition (P1) of Theorem 1.1. is satisfied. The condition (P2) follows from the fact that the projections $f_\alpha: X \rightarrow X_\alpha$, $\alpha \in A$, are open-closed. (See 1.2. Remark.). The proof is completed.

1.17. If the spaces X_α are compact (countably compact), then (C2) means that $f_\alpha^{-1}(x_\alpha)$ is finite.

1.18. Lemma. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ is an inverse system such that the cardinality $|f_{\alpha\beta}^{-1}(x_\alpha)| \leq l$ for each fiber $f_{\alpha\beta}^{-1}(x_\alpha)$ and some fixed natural number l . Then $|f_\alpha^{-1}(x_\alpha)| \leq l$ for every $f_\alpha^{-1}(x_\alpha)$, $\alpha \in A$.

PROOF. Trivial.

From Theorem 1.16. and Lemma 1.18. it follows

1.19. Theorem. Let $\underline{X} = \{X_n, f_{nm}, N\}$ be an inverse sequence of metric k -dimensional Cantor-manifolds. If the mappings f_{nm} are open onto mappings with the property that there exists a natural number $l \geq 1$ such that $|f_{nm}^{-1}(x_n)| \leq l$ for every n, m and x_n , then $X = \varprojlim X$ is a metric k -dimensional Cantor-manifold.

By the same method of proof we have the next theorem.

1.20. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of k -dimensional metric Cantor-manifolds. If the mappings $f_{\alpha\beta}$ are open surjections with the property that for every n and x_n there exist a natural numbers m, l such that $|f_{nm}^{-1}(x_n)| \leq l$, then $X = \varprojlim X$ is a Cantor-manifold.

PROOF. The space X is metrizable [3] i.e. separable metric space since X is compact. Theorem 1.16. completes the proof.

1.21. Lemma. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an σ -directed inverse system. If the fiber $f_{\alpha\beta}^{-1}(x_\alpha)$ are finite, then the fiber $f_\alpha^{-1}(x_\alpha)$ are finite.

PROOF. Suppose that $|f_\alpha^{-1}(x_\alpha)| = \{x^{(1)}, x^{(2)}, \dots, x^{(n)}, \dots\}$. For every pair of points $x^{(i)}, x^{(j)}$ there exists $\alpha(i, j)$ such that $f_\beta(x^{(i)}) \neq f_\beta(x^{(j)})$ for each $\beta \cong \alpha(i, j)$. Since $\underline{X}$ is σ -directed, there exist $\gamma \cong \alpha(i, j)$, $i \in N$, $j \in N$. This means that the cardinality of the set $f_\gamma f_\alpha^{-1}(x_\alpha) = f_{\alpha\gamma}^{-1}(x_\alpha)$ is $\aleph_0$. A contradiction!

1.21. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be a σ -directed inverse system of k -dimensional locally connected metric Cantor-manifolds X such that $f_{\alpha\beta}^{-1}(x_\alpha)$ are finite subsets of X_β , then $X = \varprojlim X$ is k -dimensional locally connected metric Cantor-manifold.

PROOF. X is locally connected [9]. This means that $w(X) = \aleph_0$ [22: Theorem 1.] i.e. X is separable metric space. Now, apply Theorem 1.16.

1.22. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be a well-ordered inverse system such that $\text{cf}(A) > \omega_1$. If the fibers $f_{\alpha\beta}^{-1}(x_\alpha)$ are finite, and if X_α are metric k -dimensional Cantor-manifold, then $X = \varprojlim X$ is k -dimensional metric Cantor-manifold.

PROOF. The space X is metric since $w(X) \equiv \aleph_0$ [28]. As in the preceding theorems we infer that X is a Cantor-manifold.

In the non-metric case, we can prove

1.23. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of n -dimensional Cantor-manifold X_α . If the mappings $f_\alpha: X = \varprojlim X \rightarrow X_\alpha, \alpha \in A$, are open with finite fibers, then X is n -dimensional Cantor-manifold.

PROOF. From [14: III. 2. Theorem] it follows that $\dim f_\alpha(F) = \dim F$ for every $\alpha \in A$ and every closed $F \subseteq X$. Hence $\dim F = n$. Furthermore, the property (P1) of Theorem 1.1. is satisfied. For (P2) see 1.2. Remark.

1.24. Remark. From [14: III. 2. Theorem] it follows also that Theorem 1.23. holds for inverse system X of generalized Cantor-manifolds always when f_α are open-closed mappings with finite fibers and normal limit X . This is the case, for example, for inverse sequence of countable compact normal generalized Cantor-manifolds since X is normal countably compact spaces [18]. From [6: 2.7.15(b)] it follows that Theorem 1.23. holds for inverse sequence of perfectly normal generalized Cantor-manifolds with open-closed onto bonding projections which have finite fiber.

1.25. Corollary. If $\underline{X}$ in Theorem 1.23. is σ -directed or $|f^{-1}(x)| \leq l$ for each α, β, x_α and some fixed natural number l , then X is n -dimensional Cantor-manifold.

1.26. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of n -dimensional Cantor-manifold. If there exists a natural number $l \geq 1$ such that $|f_{\alpha\beta}^{-1}(x_\alpha)| \leq l$ for each α, β and x_α , then $X = \varprojlim X$ is $(\dim X, \dim X - n + l + 1)$ -Cantor-manifold.

PROOF. Let F be a closed subset of X such that $\dim F \leq \dim X - (\dim X - n + l + 1) = n - l - 1$. By [1: 450] and Lemma 1.18. we have $\dim f_\alpha(F) \leq \dim F + l - 1 \leq n - l - 1 + l - 1 = n - 2$. The condition (P1) of Theorem 1.1. is satisfied. This means that the set $Y_\alpha = X_\alpha \setminus f_\alpha(F)$ is connected. For every $\beta > \alpha$ the set $Y'_{\alpha\beta} = f_{\alpha\beta}^{-1}f_\alpha(F)$ has the dimension $\leq n - 2$ [1: 452] since $\dim f_\alpha \leq 0$. It follows that the set $Y_{\alpha\beta} = X_\beta \setminus Y'_{\alpha\beta} = f_{\alpha\beta}^{-1}(Y_\alpha)$ is connected. The inverse system $Y = \{Y_{\alpha\beta}, \beta \geq \alpha\}$ has a connected limit [6: 6.1.18. Theorem]. Since $\lim Y = f_\alpha^{-1}(Y_\alpha)$ we infer that X satisfies (P2) of Theorem 1.1. This means that $X \setminus F$ is connected i.e. X is $(\dim X, \dim X - n + l + 1)$ -Cantor-manifold. Q.E.D.

1.27. Remark. If $\dim X = n - l + 1$, then X is $(n - l + 1)$ -dimensional Cantor-manifold.

Now we consider the inverse system with monotone bonding mappings.

1.28. Theorem. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of hereditarily normal Ind - n -dimensional Cantor-manifolds X_α . If $f_{\alpha\beta}$ are monotone mappings such that there

exists a natural number $l \geq 1$ such that $|\text{Fr } f_{\alpha\beta}^{-1}(x_\alpha)| \leq l$ then $X = \varprojlim X$ is $(\text{Ind } X, \text{Ind } X - n + l + 1)$ -Cantor-manifold.

PROOF. It is readily seen that $|\text{Fr } f_\alpha^{-1}(x_\alpha)| \leq l$. From [24: Theorem VII. 8] it follows that $\text{Ind } f_\alpha(F) \leq \text{Ind } F + l - 1 \leq (\text{Ind } X - \text{Ind } X + n - l - 1) + l - 1 = n - 2$ for each closed $F \subseteq X$ and each $\alpha \in A$. This means that $Y_\alpha = Y_\alpha \setminus f_\alpha(F)$ is connected. Furthermore, the set $f_\alpha^{-1}(Y_\alpha)$ is connected since f_α is a monotone mapping. Theorem 1.1. completes the proof.

1.29. Corollary. Let $\underline{X} = \{X_\alpha, f_{\alpha\beta}, N\}$ be an inverse system of metric n -dimensional Cantor-manifolds. If $f_{\alpha\beta}$ are monotone mappings such that $|\text{Fr } f_{\alpha\beta}^{-1}(x_\alpha)| \leq l, l \geq 1$, then $\varprojlim X$ is $(\text{Ind } X, \text{Ind } X - n + l + 1)$ -Cantor-manifold.

1.30. Corollary. If X in the preceding theorem is an inverse sequence, then $\varprojlim X$ is $(\dim X, \dim X - n + l + 1)$ -Cantor-manifold.

1.31. Corollary. If in the preceding Theorems $|\text{Fr } f_{\alpha\beta}^{-1}(x_\alpha)| = 1$, then $\text{Ind } X = n$, $\dim X = n$ respectively. This means that in 1.30. X is n -dimensional Cantor-manifold.

1.32. Remark. The inductively-open mapping were introduced by Arhangel'skii [4: 209]. A mapping $f: X \rightarrow Y$ is said to be inductively-open if there exists a subspace $X_1 \subseteq X$ such that $f(X_1) = Y$ and if $f|_{X_1}: X_1 \rightarrow Y$ is open.

If $f: X \rightarrow Y$ an inductively-open and closed mapping between metric spaces X and Y such that $|f^{-1}(y)| \leq \aleph_0, y \in Y$, then $\dim Y \leq \dim X$ [4: 9.1. Theorem]. If we assume that X and Y are Cantor-manifolds and $|f^{-1}(y)| \leq k$, then $\dim X \leq \dim Y + \dim f = \dim Y$.

1.33. Problem. Is it true that the assumption that $f_{\alpha\beta}$ are open in Theorem 1.16., 1.19., 1.20—1.27. can be replaced by the assumption " $f_{\alpha\beta}$ are inductively-open"?

Acknowledgement. The author wish thanks to the referee for several valuable suggestions.

References

- [1] P. S. ALEKSANDROV, V. A. PASYNKOV, Vvedeniye v teoriju razmesnosti, Nauka, Moskva, 1973.
- [2] A. T. AL-UNI, A. R. PEARNS, On a result of Nagami, *Rafidan J. sci.* **1** (1976), 89—94.
- [3] ARHANGEL'SKII, A theorem on the metrizability of the inverse image of a metric space under an open-closed finite-to-one mapping: Example and unsolved problems (in Russian) *DAN SSSR* **170** (1966), 1258—1262.
- [4] ARHANGEL'SKI, Otobraženija otkrytie i bliskie k okrytim. Svjazi meždu prostranstvami, *UMN* **15** (1966), 181—223.
- [5] ARHANGEL'SKII, On closed maps, increasing dimension, *Čech. Mat. J.* **18** (93) (1968), 38—391.
- [6] R. ENGELKING, General Topology, PWN, Warszawa, 1977.
- [7] R. ENGELKING, Dimension Theory, PWN, Warszawa, 1978.
- [8] V. V. FEDORČUK, Metod razvertivaemyh spektrov i vpol'ne zamkunyh otobraženij v obščej topologii, *UMN* **35** (1980), 112—121.
- [9] G. R. GORDON and S. MARDEŠIĆ, Characterizing local connectedness of inverse limits, *Pacific J. Math.* **58** (1975), 411—417.
- [10] N. HADŽIIVANOV, On infinite-dimensional Cantor-manifolds, *Topics Topology, Amsterdam—London*, 1974, 355—363.
- [11] J. E. JAYNE, C. A. ROGERS, Functions fermées en partie, *C. r. Acad. Sci.* **AB291**, no 13, A667—, A670 (1980).
- [12] C. F. K. JUNG, Mappings on compact metric spaces, *Colloq. Math.* **19** (1968), 74—76.

- [13] H. KATÓ, A note on infinite-dimension under refinable maps, *Proc. Amer. Math. Soc.* **88** (1983) 177—180.
- [14] J. E. KEESLING, Open and closed mappings and compactification, *Fund. Math.* **65** (1969), 73—81.
- [15] J. E. KEESLING, Closed mappings which lower dimension, *Colloq. Math.* **20** (1969), 237—241.
- [16] K. KURATOVSKI, *Topologija I*, Mir, Moskva, 1966.
- [17] K. KURATOVSKI, *Topologija II*, Mir, Moskva, 1969.
- [18] I. LONČAR, Inverse limits for spaces which generalize compact spaces, *Glasnik matematički* **17** (37) (1982), 155—173.
- [19] I. LONČAR, Some results on the dimension of the inverse limit spaces (*to appear in Glasnik matematički*).
- [20] I. LONČAR, Lindelöfov broj i inverzni sistemi, *Zbornik radova Fakulteta organizacije i informatike Varaždin* **7** (1983), 115—123.
- [21] A. LELEK, Dimension inequalities for unions and mappings of separable metric spaces, *Colloq. Math.* **23** (1971), 70—91.
- [22] S. MARDEŠIĆ, Lokalno povezani, uređeni i lančasti kontinuumi, *JAZU Zagreb*, (1960), 147—166.
- [23] K. NAGAMI, Mappings of finite order and dimension theory, *Jap. Journ. of Math.* **30** (1960), 25—54.
- [24] J. NAGATA, Modern dimension theory, *Amsterdam*, 1965.
- [25] V. A. PASYNKOV, Častične topologičke proizvedenija, *Trudy Mosk. mat. obščestva*, **13** (1965), 136—245.
- [26] L. T. POLKOWSKI, Open and closed mappings and infinite dimension, *Gen. Topol. and Relat. Mod. Anal. and Algebra*, **5** Berlin (1983), 561—564.
- [27] E. V. ŠČEPIN, Funktory i nesčetnye stepeni kompakto, *UMN* **36** (1981), 3—62.
- [28] M. G. TKAČENKO, Capi i kardinaly, *DAN SSSR* **239** (1978), 546—549.
- [29] Y. YAJIMA, On the dimension of limits of inverse systems, *Proc. Amer. Mat. Soc.* **91** (1984), 461—466.

FAKULTET ORGANIZACIJE I INFORMATIKE,
VARAŽDIN YUGOSLAVIA

(Received March 20, 1985.)