

Non-convex perturbations of evolution equations with m -dissipative operators in Banach spaces

E. P. AVGERINOS (Thessaloniki) and N. S. PAPAGEORGIOU (Davis)

Abstract. In this paper we establish the existence of integral solutions for a nonlinear, multivalued evolution equation of the form $\dot{x}(t) \in Ax(t) + F(t, x(t))$, where $A : X \rightarrow 2^X$ is an m -dissipative operator and $F(\cdot, \cdot)$ a nonconvex valued perturbation. Our result generalizes a recent existence theorem of Cellina-Marchi (*Israel J. Math* 46 (1983), pp. 1-11).

Key words and phrases : m -dissipative operator, compact semigroup, lower semicontinuous multifunction, Arzela-Ascoli theorem, parabolic equation.

1. Introduction

Evolution equations of the form $-\dot{x}(t) \in Ax(t) + f(t)$ in a Hilbert space, were first studied by BREZIS [4], with A a maximal monotone operator and $f(\cdot)$ an integrable perturbation. The work of BREZIS was extended by ATTOUCH-DAMLAMIAN [1], to systems of the form $-\dot{x}(t) \in Ax(t) + F(t, x(t))$, with $F(\cdot, \cdot)$ being a multivalued perturbation having convex values. ATTOUCH-DAMLAMIAN [1] proved two existence results : one with A being a general maximal monotone operator, but with the underlying state space being $\mathbf{R}^n$ and the other with A being a subdifferential (i.e. $A = \partial\phi$, with ϕ being a proper, closed, convex function) and the underlying state space being any separable Hilbert space. Recently CELLINA-MARCHI [6] proved an existence theorem for the case where the multivalued perturbation has nonconvex values and the state spaces is $\mathbf{R}^n$. The study of those evolution equations in general Banach space (not necessarily Hilbert), was initiated by PAZY [12], who considered the case of A being a densely defined, linear, m -accretive operator and the perturbation was single valued. A nonlinear version of Pazy's theorem was

proved by VRABIE [14], who also considered the case of multivalued perturbations with convex values, extending this way the work of ATTOUCH-DAMLAMIAN [1]. Other interesting works in these or related issues were done by GUTMAN [8], HARAUX [9] and SCHECHTER [13] (he studied the dependence of the solutions on variations of the initial data).

In this note, we extend the result of CELLINA-MARCHI [6] to arbitrary separable Banach spaces, weakening also the hypotheses on the multivalued perturbation $F(t, x)$. Instead of assuming joint Hausdorff continuity for $F(t, x)$, we only require lower semicontinuity in the variable x , a more natural hypothesis in the context of applications.

2. Preliminaries

Let (Ω, Σ) be a measurable space and X a separable Banach space. By $P_f(X)$ we will denote the collection of all nonempty, closed subsets of X . A multifunction $F : \Omega \rightarrow P_f(X)$ is said to be graph measurable, if $GrF = \{(\omega, x) \in \Omega \times X : x \in F(\omega)\} \in \Sigma \times B(X)$, where $B(X)$ is the Borel σ -field of X . Now let $\mu(\cdot)$ be a σ -finite measure on Σ . By S_F^1 we will denote the set of integrable selectors of $F(\cdot)$ i.e. $S_F^1 = \{f \in L^1(X) : f(\omega) \in F(\omega) \mu - \text{a.e.}\}$. Using Aumann's selection theorem, it is easy to check that if $\omega \rightarrow |F(\omega)| = \sup\{\|x\| : x \in F(\omega)\}$ is in L_+^1 (in which case we say that $F(\cdot)$ is integrably bounded), then $S_F^1 \neq \emptyset$. If Y, Z are Hausdorff topological spaces and $G : Y \rightarrow 2^Z \setminus \{\emptyset\}$, then we say that $F(\cdot)$ is lower semicontinuous (l.s.c.), if for all $U \subseteq Z$ open, the set $G^-(U) = \{y \in Y : G(y) \cap U \neq \emptyset\}$ is open in Y . If Y, Z are metric spaces, then the above definition is equivalent to saying that for all $y_n \rightarrow y$ we have $G(y) \subseteq \varliminf G(y_n) = \{z \in Z : z = \lim z_n, z_n \in G(y_n)\}$.

Next let X be any Banach space. Let $J : X \rightarrow 2^{X^*}$ be the duality map of X i.e. $J(x) = \{x^* \in X^* : (x^*, x) = \|x\|^2 = \|x^*\|^2\}$. Clearly the values of $J(\cdot)$ are closed, convex, bounded subsets of X^* , which because of the Hahn-Banach theorem are also nonempty. Recall that if X^* is strictly convex, then $J(\cdot)$ is single valued. Using $J(\cdot)$ we can define the upper semi-inner product (denoted by $(\cdot, \cdot)_+$) and the lower semi-inner product (denoted by $(\cdot, \cdot)_-$) as follows:

$$(x, y)_+ = \sup\{(x^*, y) : x^* \in J(x)\}$$

$$\text{and } (x, y)_- = \inf\{(x^*, y) : x^* \in J(x)\}$$

for all $x, y \in X$. An operator $A : X \rightarrow 2^X$ is said to be dissipative (see BARBU [2]), if $(x - x', y - y')_- \leq 0$ for any $(x, y)(x', y') \in GrA$. We say that A is m -dissipative, if it is dissipative and in addition $R(I - \lambda A) = X$ for all $\lambda > 0$. It is well known that an m -dissipative operator generates a semigroup $\{S(t)\}_{t \geq 0}$ of nonlinear contractions, via the Crandall-Liggett formula

$$S(t)x = \lim_{n \rightarrow \infty} (I - \frac{t}{n}A)^{-n} x \quad t \geq 0, \quad x \in \overline{D(A)}.$$

Now let A be an m -dissipative operator, $f \in L^1(X)$ and $x_0 \in \overline{D(A)}$. Consider the following Cauchy problem on $T = [0, b]$:

$$\left\{ \begin{array}{l} \dot{x}(t) \in Ax(t) + f(t) \\ x(0) = x_0 \end{array} \right\} \quad (*)$$

Following BENILAN [3], we say that a function $x \in C(T, X)$ is an "integral solution" of $(*)$, if $x(0) = x_0$ and

$$\|x(t) - y\|^2 \leq \|x(s) - y\|^2 + 2 \int_s^t (f(r) + z, x(r) - y)_+ dr$$

for all $(y, z) \in Gr A$ and all $0 \leq s \leq t \leq b$.

It is well known that under the above hypotheses Cauchy problem $(*)$ has a unique integral solution. Moreover this unique integral solution depends continuously on the data of the problem. In fact if $x_1(\cdot)$ is the solution of $(*)$ with data $(x_{01}, f_1) \in \overline{D(A)} \times L^1(X)$ and $x_2(\cdot)$ the solution of $(*)$ with data $(x_{02}, f_2) \in \overline{D(A)} \times L^1(X)$, then we have

$$\|x_1(t) - x_2(t)\|^2 \leq \|x_{01} - x_{02}\|^2 + 2 \int_0^t (f_1(r) - f_2(r), x_1(r) - x_2(r))_+ dr, t \in T,$$

or equivalently

$$\|x_1(t) - x_2(t)\| \leq \|x_{01} - x_{02}\| + \int_0^t \|f_1(r) - f_2(r)\| dr.$$

If A is densely defined, linear, m -accretive, then the notion of integral solution coincides with that of mild solution.

Recall that a "strong solution" of $(*)$ is a continuous function $x : T \rightarrow X$ (i.e. $x(\cdot) \in C(T, X)$), for which we have that $x(t) \in D(A)$, is differentiable a.e. on $(0, b)$ and satisfies $(*)$ a.e. with $x(0) = x_0 \in D(A)$.

Every strong solution is an integral solution. The converse is true only if we impose additional hypotheses on X, A and f . We are not going to go into the details of that problem. We only mention that if $X = \mathbf{R}^n$ and A is maximal monotone or if X is a Hilbert space and $A = \partial\phi$, with ϕ being a proper, closed, convex function on X , then every integral solution is also strong for any initial condition $x_0 \in \overline{D(A)}$. For further details we refer to BARBU [2], BREZIS [4] and SCHECHTER [13].

3. The theorem

In this section we will establish the existence of an integral solution for the following multivalued evolution equation:

$$\begin{cases} \dot{x}(t) \in Ax(t) + F(t, x(t)) \\ x(0) = x_0 \end{cases} \quad (**)$$

By an integral solution of (**), we mean a function $x \in C(T, X)$, which is an integral solution (as defined in section 2) of $\dot{x}(t) \in Ax(t) + f(t)$, $x(0) = x_0$ for some $f \in S_{F(\cdot, x(\cdot))}^1$.

Let $T = [0, b]$ and let X be a separable Banach space. We will need the following hypotheses:

H(A): $A : X \rightarrow 2^X$ is an m -dissipative operator, which generates a semigroup of compact nonlinear contractions

(i.e. $S(t) : \overline{D(A)} \rightarrow \overline{D(A)}$ is compact for $t > 0$),

H(F): $F : T \times X \rightarrow P_f(X)$ is a multifunction s.t.

(1) $(t, x) \rightarrow F(t, x)$ is graph measurable,

(2) for every $t \in T$, $x \rightarrow F(t, x)$ is l.s.c.,

(3) $|F(t, x)| = \sup\{\|y\| : y \in F(t, x)\} \leq a(t) + b(t) \|x\|$ a.e. with $a(\cdot), b(\cdot) \in L_+^1$.

H₀ : $x_0 \in \overline{D(A)}$.

We have the following existence result concerning (**).

Theorem. *If hypotheses H(A), H(F) and H₀ hold, then (**) admits an integral solution.*

PROOF. We will start by determining an a priori bound for the integral solutions of (**). So suppose $x(\cdot) \in C(T, X)$ is such a solution of (**). Recalling that $S(t)x_0$ is the integral solution of $\dot{y}(t) \in Ay(t)$, $y(0) = x_0$ and using the inequalities of section 2, we have :

$$\|x(t) - S(t)x_0\| \leq \int_0^t \|f(s)\| ds$$

for all $t \in T$ and some $f \in S_{F(\cdot, x(\cdot))}^1$. Since $t \rightarrow S(t)x_0$ is continuous on T and using hypothesis $H(F)$ (3), we have

$$\|x(t)\| \leq M_1 + \int_0^t [a(s) + b(s)\|x(s)\|] ds$$

for some $M_1 > 0$. Applying Gronwall's inequality, we get that

$$\|x(t)\| \leq K \exp \|b\|_1 = M_2$$

where $K = M_1 + \|a\|_1$. Then define a new multifunction $\hat{F} : T \times X \rightarrow P_f(X)$ as follows:

$$\hat{F}(t, x) = \begin{cases} F(t, x) & \text{if } \|x\| \leq M_2 \\ F\left(t, \frac{M_2 x}{\|x\|}\right) & \text{if } \|x\| > M_2 \end{cases}$$

Observe that $\hat{F}(t, x) = F(t, p_{M_2}(x))$, where $p_{M_2}(\cdot)$ is the M_2 -radial retraction. We have $Gr\hat{F} = \{(t, x, y) \in T \times X \times X : (t, p_{M_2}(x), y) \in GrF\}$. Let $r : T \times X \times X \rightarrow T \times X \times X$ be defined by $r(t, x, y) = (t, p_{M_2}(x), y)$. Recalling that $p_{M_2}(\cdot)$ is 2-Lipschitz, we have that $r(\cdot, \cdot, \cdot)$ is continuous, hence measurable. So, since $GrF \in \Sigma \times B(X) \times B(X)$, we have $r^{-1}(GrF) = Gr\hat{F} \in \Sigma \times B(X) \times B(X)$ i.e. $\hat{F}(\cdot, \cdot)$ is graph measurable. Also since $\hat{F}(t, \cdot)$ is the composition of the Lipschitz function $p_{M_2}(\cdot)$ with the l.s.c. multifunction $F(t, \cdot)$, we have that $\hat{F}(t, \cdot)$ is l.s.c.. Finally note that $|F(t, x)| \leq a(t) + M_2 b(t) = \gamma(t)$ a.e. with $\gamma(\cdot) \in L^1_+$.

In the sequel we will consider the following multivalued Cauchy problem:

$$\left\{ \begin{array}{l} \dot{x}(t) \in Ax(t) + \hat{F}(t, x(t)) \\ x(0) = x_0 \end{array} \right\} \quad (**)'$$

Let $h \in L^1(X)$ and consider the Cauchy problem

$$\left\{ \begin{array}{l} \dot{x}(t) \in Ax(t) + h(t) \\ x(0) = x_0 \end{array} \right\} \quad (***).$$

We know (see section 2), that $(***)$ has a unique integral solution. Let $r : L^1(X) \rightarrow C(T, X)$ be the map that to each $L^1(X)$ -perturbation $h(\cdot)$ assigns the corresponding unique integral solution $r(h)(\cdot) \in C(T, X)$ of $(***)$. Let $B(\gamma) = \{h \in L^1(X) : \|h(t)\| \leq \gamma(t) \text{ a.e.}\}$. Our claim is that $K = r(B(\gamma))$ is relatively compact in $C(T, X)$.

To this end, first we will show that for every $t \in T$, $K(t) = r(B(\gamma))(t) = \{x(t) : x(\cdot) = r(h)(\cdot), h \in B(\gamma)\}$ is compact in X . For $t = 0$, we have $K(0) = \{x_0\}$ and so the claim is automatically verified. Hence let $t > 0, t \in T$. Note that $B(\gamma)$ is a uniformly integrable subset of $L^1(X)$. So given $t \in (0, b]$ and $\epsilon > 0$, we can find $\delta(\epsilon) \in (0, t)$ s.t. for $B \subseteq T$ Lebesgue measurable with $\lambda(B) < \delta$, we have:

$$\int_B \|h(s)\| ds < \epsilon$$

for all $h \in B(\gamma)$. Now consider the following Cauchy problem; on $[t - \delta, t]$:

$$\left\{ \begin{array}{l} \dot{x}(\delta)(s) \in Ax(\delta)(s) \\ x(\delta)(t - \delta) = r(h)(t - \delta) \end{array} \right\}$$

where $h \in B(\gamma)$. From the inequalities of section 2, we have:

$$\|x(\delta)(t) - r(h)(t)\| \leq \int_{t-\delta}^t \|h(s)\| ds < \epsilon$$

for all $h \in B(\gamma)$. Also recall that

$$x(\delta)(t) = S(\delta)r(h)(t - \delta) \subseteq S(\delta)K(t - \delta).$$

and the latter is relatively compact in X , since $K(t - \delta) = \{y(t - \delta) : y(\cdot) \in K\}$ is bounded and $S(\delta)$ is a compact contraction (see hypothesis $H(A)$). Therefore $\overline{S(\delta)K(t - \delta)}$ is compact. So for every $t \in T$, every $\epsilon > 0$ and every $z \in K(t)$, there exists an element z_ϵ in the compact set $\overline{S(\delta)K(t - \delta)}$ s.t. $\|z - z_\epsilon\| < \epsilon \implies \overline{K(t)}$ is compact.

Next, recall that since the semigroup $S(t)$ is compact, for $B \subseteq X$ nonempty, bounded, we have that $t \rightarrow \{S(t)x : x \in B\}$ is equicontinuous on $t > 0$. Hence given $\epsilon > 0$, we can find $\delta_1(\epsilon) > 0$ s.t. for $|t' - t| < \delta_1$ and for all $x \in K(t - \delta)$ we have:

$$\begin{aligned} & \|S(t' - t + \delta)x - S(\delta)x\| < \epsilon \\ \implies & \|S(t' - t + \delta)r(h)(t - \delta) - S(\delta)r(h)(t - \delta)\| < \epsilon \\ \implies & \|x(\delta)(t') - x(\delta)(t)\| < \epsilon. \end{aligned}$$

So finally for $\delta_2 = \min(\delta, \delta_1)$ and for $|t' - t| < \delta_2$, we have

$$\begin{aligned} & \|r(h)(t') - r(h)(t)\| \\ \leq & \|r(h)(t') - x(\delta)(t')\| + \|x(\delta)(t') - x(\delta)(t)\| + \|x(\delta)(t) - r(h)(t)\| < \\ & < \epsilon + \epsilon + \epsilon = 3\epsilon \implies K = r(B(\gamma)) \text{ is equicontinuous at each } t > 0. \end{aligned}$$

Finally for $t = 0$ note that

$$\begin{aligned} \|r(h)(t) - S(t)x_0\| & \leq \|r(h)(t) - S(t)x_0\| + \|S(t)x_0 - x_0\| \\ & \leq \int_0^t \gamma(s)ds + \|S(t)x_0 - x_0\| \end{aligned}$$

So we also have equicontinuity at $t = 0$.

Invoking the Arzela–Ascoli theorem, we conclude that $\overline{K}$ is compact in $C(T, X)$. Thus by Mazur's theorem $\overline{K}_c = \overline{\text{conv}} K$ is compact

Next let $R : \overline{K}_c \rightarrow 2^{L^1(X)}$ be defined by

$$R(x) = S_{\hat{F}(\cdot, x(\cdot))}^1.$$

Since $\hat{F}(\cdot, \cdot)$ is graph measurable, it is easy to check as before, that $t \rightarrow \hat{F}(t, x(t))$ is graph measurable and integrably bounded by $\gamma(\cdot)$ and so $R(\cdot)$ has nonempty values, in fact $R(\cdot)$ is $P_f(L^1(X))$ -valued. Also since $\hat{F}(t, \cdot)$ is l.s.c. and using theorem 4.1 of [11], we have that if $x_n \rightarrow x$ in $\overline{K}_c$, then $R(x) \subseteq s - \lim R(x_n)$, where s indicates the strong topology on $L^1(X)$. So $R(\cdot)$ is l.s.c. (see section 2). Hence we can apply Fryszkowski's

selection theorem [7], to get $v : \overline{K}_c \rightarrow L^1(X)$ continuous s.t. $v(x) \in R(x)$ for all $x \in \overline{K}$. Set $p = \text{rov}$. Clearly $p : \overline{K}_c \rightarrow \overline{K}_c$ is continuous. Apply Schauder's fixed point theorem to get $\hat{x} \in \overline{K}_c$ s.t. $\hat{x} = p(\hat{x}) = r(v(\hat{x}))$. Hence we have that $\hat{x}(\cdot)$ is an integral solution of

$$\left\{ \begin{array}{l} \hat{x}(t) \in A\hat{x}(t) + v(\hat{x})(t) \\ \hat{x}(0) = x_0 \end{array} \right\}$$

with $v(\hat{x})(\cdot) \in S^1_{\hat{F}(\cdot, \hat{x}(\cdot))}$. So $\hat{x}(\cdot) \in C(T, X)$ is an integral solution of $(**)'$. From the definition of $\hat{F}(t, x)$ and hypothesis $H(F)$ (3), we see easily that $|\hat{F}(t, x)| \leq a(t) + b(t)\|x\|$ a.e.. So as before, through Gronwall's inequality, we get $\|\hat{x}(t)\| \leq M_2, t \in T \implies \hat{F}(t, \hat{x}(t)) = F(t, \hat{x}(t)), t \in T \implies \hat{x}(\cdot)$ is the desired integral solution of $(**)$. Q.E.D.

As we mentioned in section 2, when $X = \mathbf{R}^n$, then every integral solution is a strong solution. So we can state as a corollary to our theorem, an extension of the existence result of CELLINA-MARCHI [6].

So let $T = [0, b], X = \mathbf{R}^n$ and make the following hypothesis about A : $H(A)' : A : D(A) \subseteq \mathbf{R}^n \rightarrow 2^{\mathbf{R}^n}$ is a maximal monotone operator.

Then we get as a corollary to our theorem, the following extension of the work of CELLINA-MARCHI [6].

Corollary. *If hypotheses $H(A)', H(F)$ and H_0 hold, then $(**)$ admits a strong solution.*

Remarks. (1) In CELLINA-MARCHI [6], the multivalued perturbation $F(t, x)$ was assumed to be jointly Hausdorff continuous.

(2) Hypotheses $H(F)$ (1) and (2), cover the case where $t \rightarrow F(t, x)$ is graph measurable and $x \rightarrow F(t, x)$ is Hausdorff continuous (see theorem 3.3 of [10]).

4. An example

Let Ω be a bounded open domain in $\mathbf{R}^n$ with smooth boundary $\partial\Omega = \Gamma, n > 2$.

Let $r > (n-2)/n$ and consider the following multivalued, nonlinear, parabolic partial differential equation on $T \times \Omega$:

$$\left\{ \begin{array}{l} \frac{\partial x(t, z)}{\partial t} - \Delta x(t, z) |x(t, z)|^{r-1} \in F(t, z, x(t, z)) \\ x(t, z) = 0 \text{ on } T \times \Gamma \\ x(0, z) = x_0(z) \text{ on } \{0\} \times \Omega \end{array} \right\} \quad (****)$$

Here $F : T \times \Omega \times \mathbf{R} \rightarrow P_f(\mathbf{R})$ is a multifunction which is l.s.c. in the third variable and $(t, y) \rightarrow S_{F(t, \cdot, y(\cdot))}^1$ is graph measurable on $T \times L^1(\Omega)$. It is easy to check that this is the case if $(t, z) \rightarrow F(t, z, r)$ is measurable and $r \rightarrow F(t, z, r)$ is Hausdorff continuous. Also assume that $|F(t, z, r)| = \sup\{\|v\| : v \in F(t, z, r)\} \leq a(t, z) + b(t, z)|r|$ a.e. with $a(\cdot, \cdot) \in L_+^1(T \times \Omega)$ and $b(t, \cdot) \in L^\infty(\Omega)$ while $t \rightarrow \|b(t, \cdot)\|_\infty$ belongs in L_+^1 . Furthermore let $\hat{x}_0 = x_0(\cdot) \in L^1(\Omega)$.

Take $X = L^1(\Omega)$. This is a separable Banach space. Consider the nonlinear operator $A : D(A) \subseteq X \rightarrow X$ defined by $Ax = \Delta x|x|^{r-1}$ with $D(A) = \{x \in X : x|x|^{r-1} \in W_0^{1,1}(\Omega), \Delta x|x|^{r-1} \in L^1(\Omega)\}$. From BREZIS [5] we know that the operator A defined above is m -dissipative and the nonlinear semigroup it generates is compact for $t \in (0, b]$. Also let $\hat{F} : T \times X \rightarrow P_f(L^1(X))$ be defined by $\hat{F}(t, x) = S_{F(t, \cdot, x(\cdot))}^1$. Then $\hat{F}(\cdot, \cdot)$ is graph measurable, $\hat{F}(t, \cdot)$ is l.s.c. (see theorem 4.1 of [11]) and

$$|\hat{F}(t, x)| \leq \hat{a}(t) + \hat{b}(t)\|x\|_1 \text{ a.e.}$$

with $\hat{a}(t) = \|a(t, \cdot)\|_1$ and $\hat{b}(t) = \|b(t, \cdot)\|_\infty$.

Rewrite the initial-boundary value problem (****) as the following abstract multivalued evolution equation:

$$\left\{ \begin{array}{l} \dot{x}(t) \in Ax(t) + \hat{F}(t, x(t)) \\ x(0) = \hat{x}_0 \end{array} \right\} \quad (****)'$$

We see that all hypotheses of our theorem are satisfied and so we know that (****)' has an integral solution $\hat{x}(\cdot) \in C(T, L^1(\Omega))$. Set $x(t, z) = \hat{x}(t)(z)$ $z \in \Omega$. This is a generalized solution of (****).

Acknowledgement. The second author was supported by NSF Grant DMS - 8802688

References

- [1] H. ATTOUCH - A. DAMLAMIAN, On multivalued evolution equations in Hilbert spaces, *Israel J. Math* **12** (1972), 373-390.
- [2] V. BARBU, Nonlinear Semigroups and Differential Equations in Banach Spaces, *Noordhoff International Publishing, Leyden, Netherlands*, 1976.
- [3] PH. BENILAN, Solutions integrales d'equations d'evolution dans un espace de Banach, *C.R. Acad. Sc. Paris*, t. **274** (1972), 47-50.
- [4] H. BREZIS, Operateurs Maximaux Monotones, *Math. Studies, Vol. 50, North Holland, Amsterdam*, 1972.
- [5] H. BREZIS, New results concerning monotone operators and nonlinear semigroups, *Analysis of Nonlinear problems, RIMS Kyoto Univ. No. 258* (1974), 2-27.
- [6] A. CELLINA - M. MARCHI, Nonconvex perturbations of maximal monotone inclusions, *Israel J. Math* **46** (1983), 1-11.

- [7] A. FRYSZKOWSKI, Continuous selections for a class of nonconvex multivalued maps, *Studia Math* **76** (1983), 163–174.
- [8] S. GUTMAN, Evolutions governed by m -accretive plus compact operators, *Nonl. Anal. - T.M.A.* **7** (1983), 707–715.
- [9] A. HARAUX, Nonlinear Evolution Equations : Global Behavior of Solutions, *Lecture Notes in Math* **841** (1981), Springer, Berlin.
- [10] N.S. PAPAGEORGIOU, On measurable multifunctions with applications to random multivalued equations, *Math. Japonica* **32** (1987), 437–464.
- [11] N.S. PAPAGEORGIOU, Convergence theorems for Banach space valued integrable multifunctions, *Intern. J. Math and Math. Sci.* **10** (1987), 433–442.
- [12] A. PAZY, A class of semilinear equations of evolution, *Israel J. Math.* **20** (1975), 23–36.
- [13] E. SCHECHTER, Perturbations of regularizing maximal monotone operators, *Israel J. Math.* **43** (1982), 49–61; and *correction* **47** (1984), 236–240.
- [14] I. VRABIE, A nonlinear version of Pazy's local existence theorem, *Israel J. Math.* **32** (1979), 221–235.

EVGENIOS P. AVGERINOS
UNIVERSITY OF THESSALONIKI
SCHOOL OF TECHNOLOGY
THESSALONIKI 54006, GREECE

NIKOLAOS S. PAPAGEORGIOU
UNIVERSITY OF CALIFORNIA
1015 DEPARTMENT OF MATHEMATICS
DAVIS, CALIFORNIA 95616, U.S.A.

(Received December 30, 1987)