

A generalization of Skiba's problem

By SHUQIN DONG (Huainan), TINGTING QIU (Yancheng), ZHICHAO GAO (Yangzhou),
LONG MIAO (Wuxi) and JINGYU HE (Wuxi)

Abstract. Let H/K be any chief factor of a finite group G , p a prime, and j a nonnegative integer. We say that $G \in \mathfrak{S}_{p^j}^*$ if every chief factor H/K is a p -group or its p -part satisfies $|H/K|_p \leq p^j$. In this paper, we generalize Skiba's problem to the class $\mathfrak{S}_{p^j}^*$ of groups containing some non-solvable groups. We prove that $G \in \mathfrak{S}_{p^j}^*$ if and only if every maximal subgroup of a Sylow p -subgroup of G has a subnormal supplement in $\mathfrak{S}_{p^j}^*$, where $p > j$ or $p = j = 2$.

SHUQIN DONG
SCHOOL OF MATHEMATICAL SCIENCES
HUAIYIN NORMAL UNIVERSITY
223300 HUAIAN
P. R. CHINA

LONG MIAO &
JINGYU HE
BASIC COURSE DEPARTMENT
JIANGSU VOCATIONAL COLLEGE
OF INFORMATION TECHNOLOGY
214153 WUXI
P. R. CHINA

TINGTING QIU
YANCHENG BIOLOGICAL ENGINEERING
HIGHER VOCATIONAL TECHNOLOGY SCHOOL
224000 YANCHENG
P. R. CHINA

ZHICHAO GAO
SCHOOL OF MATHEMATICAL SCIENCES
YANGZHOU UNIVERSITY
225002 YANGZHOU
P. R. CHINA

Mathematics Subject Classification: 20D10, 20D20.

Key words and phrases: Sylow subgroup, subnormal supplement, G -boundary factor.