

## Symbolic solutions of algebraic ODEs: a comparison of methods

By JOHANN J. MITTERAMSKOGLER (Linz) and FRANZ WINKLER (Linz)

**Abstract.** In this paper, the two methods for finding rational general solutions of first-order algebraic ODEs introduced in Ngô and Winkler ([19], [20], [21]) and Vo, Grasegger and Winkler [26] are compared. Both methods assign some affine algebraic set to an algebraic ODE. Provided the assigned algebraic sets are suitably parametrizable, the initial ODE can be reduced to a more fundamental (set of) differential equation(s). The two approaches lead to a common rational parametrization in certain situations, in which case the corresponding derived differential equation(s) are shown to coincide. Finally, a discussion on relations between certain classes of first-order algebraic ODEs with respect to their rational general solvability is provided.

JOHANN JOSEF MITTERAMSKOGLER &  
FRANZ WINKLER  
RESEARCH INSTITUTE FOR  
SYMBOLIC COMPUTATION (RISC)  
JOHANNES KEPLER UNIVERSITY LINZ  
ALTENBERGERSTRASSE 69  
4040 LINZ  
AUSTRIA

---

*Mathematics Subject Classification:* Primary: 34A26, 34A34; Secondary: 14E05, 14H50, 14J26, 68W30.

*Key words and phrases:* ordinary differential equation, rational general solution, algebraic surface, algebraic curve, rational parametrization.