

On Finsler geometry of tangent Lie groups

By MONA ATASHAFROUZ (Tehran), BEHZAD NAJAFI (Tehran)
and AKBAR TAYEBI (Qom)

Abstract. This paper is divided into two main parts. In the first part, we study left-invariant Randers metrics on Lie groups. We characterize the class of left-invariant Randers metrics with isotropic mean Berwald and isotropic Berwald curvatures on Lie groups. This yields an extension of Deng's well-known theorem for left-invariant Randers metrics with isotropic S -curvature. In the second part, we consider the left-invariant Randers metrics on tangent Lie groups. Let G be a Lie group equipped with a left-invariant Randers metric F . Suppose that F^v and F^c denote the vertical and complete lift of F on TG , respectively. First, we find the necessary and sufficient condition under which these metrics are weakly Berwaldian. Then, we prove that these lifting Randers metrics are isotropic Berwald metrics if and only if F reduces to a Berwald metric. Finally, we give the necessary and sufficient conditions under which these metrics are of Douglas-type metrics.

MONA ATASHAFROUZ &
BEHZAD NAJAFI
DEPARTMENT OF MATHEMATICS
AND COMPUTER SCIENCES
AMIRKABIR UNIVERSITY OF TECHNOLOGY
(TEHRAN POLYTECHNIC)
TEHRAN
IRAN

AKBAR TAYEBI
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
UNIVERSITY OF QOM
QOM
IRAN

Mathematics Subject Classification: 53B40, 53C60, 22E60, 22E15.

Key words and phrases: left-invariant metric, isotropic Berwald metric, Berwald metric, weakly Berwald metric, Douglas metric, \mathcal{Z} -Randers metric.