

Padovan squares which are again Padovan

By ERIC F. BRAVO (Popayán) and FLORIAN LUCA (Wits)

Abstract. The integer sequence defined by $P_{n+1} = P_{n-1} + P_{n-2}$ with initial values $P_0 = P_1 = P_2 = 1$ is known as the Padovan sequence $(P_n)_{n \in \mathbb{Z}}$. In this note, we solve the Diophantine equations $P_{-n} = \pm P_m^2$, $P_n = P_{-m}^2$, and $P_{-n} = \pm P_{-m}^2$ in positive integers n, m .

ERIC F. BRAVO
DEPARTAMENTO DE MATEMÁTICAS
UNIVERSIDAD DEL CAUCA
CALLE 5 NO 4-70
POPAYÁN
COLOMBIA

FLORIAN LUCA
SCHOOL OF MATHEMATICS
UNIVERSITY OF THE WITWATERSRAND
PRIVATE BAG 3, WITS 2050
SOUTH AFRICA

AND

CENTRO DE CIENCIAS MATEMÁTICAS
UNIVERSIDAD NACIONAL AUTÓNOMA
DE MÉXICO (UNAM)
MORELIA
MÉXICO

Mathematics Subject Classification: 11B39, 11J86.

Key words and phrases: Padovan number, linear form in logarithms, Baker's method.