

## Characterizations of Riemannian maps between Kaehler manifolds by certain curves

By TUNAHAN TURHAN (Isparta), GÖZDE ÖZKAN TÜKEL (Isparta)  
and BAYRAM ŞAHİN (Izmir)

**Abstract.** Riemannian maps between Kaehler manifolds are studied with respect to certain curves. For this purpose, the geometric properties of the Riemannian maps themselves, the total manifold and the base manifold are investigated depending on the behavior of the geodesic, circle and Kaehler Frenet curves on the total manifold under Riemannian maps.

TUNAHAN TURHAN  
DIVISION OF ELEMENTARY  
MATHEMATICS EDUCATION  
SÜLEYMAN DEMIREL UNIVERSITY  
ISPARTA  
TURKEY

GÖZDE ÖZKAN TÜKEL  
DEPARTMENT OF BASIC SCIENCES  
ISPARTA UNIVERSITY OF APPLIED SCIENCES  
ISPARTA  
TURKEY

BAYRAM ŞAHİN  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCE, EGE UNIVERSITY  
IZMIR  
TURKEY

---

*Mathematics Subject Classification:* 53C15, 53B20, 53C43.

*Key words and phrases:* Kaehler manifolds, Kaehler Frenet curve, Kaehler circle, second fundamental form, isotropic Riemannian map.