

On function SX of additive complements

By JIN-HUI FANG (Nanjing) and CSABA SÁNDOR (Budapest)

Abstract. Two sets A, B of nonnegative integers are called *additive complements*, if all sufficiently large integers can be expressed as the sum of two elements from A and B . We call A, B *perfect additive complements* if every nonnegative integer can be uniquely expressed as the sum of two elements from A and B . Let $A(x)$ be the counting function of A . In this paper, we focus on the function SX , where $SX = \limsup_{x \rightarrow \infty} \frac{\max\{A(x), B(x)\}}{\sqrt{x}}$ was introduced by Erdős and Freud in 1984. As a main result, we determine the value of SX for perfect additive complements, and further fix the infimum. We also give the absolute lower bound of SX for additive complements.

JIN-HUI FANG
SCHOOL OF MATHEMATICAL SCIENCES
NANJING NORMAL UNIVERSITY
NANJING 210023
P. R. CHINA

CSABA SÁNDOR
DEPARTMENT OF STOCHASTICS
INSTITUTE OF MATHEMATICS
BUDAPEST UNIVERSITY
OF TECHNOLOGY AND ECONOMICS
MŰEGYETEM RKP. 3
H-1111 BUDAPEST
HUNGARY

AND

HUN-REN ALFRÉD RÉNYI
INSTITUTE OF MATHEMATICS
REÁLTANODA UTCA 13-15
H-1053 BUDAPEST
HUNGARY

AND

MTA–HUN-REN RI LENDÜLET “MOMENTUM”
ARITHMETIC COMBINATORICS RESEARCH GROUP
REÁLTANODA UTCA 13–15
H-1053 BUDAPEST
HUNGARY

Mathematics Subject Classification: Primary: 11B13; Secondary: 11B34.
Key words and phrases: additive complements, perfect, infimum.