

Pseudo-Ricci–Bourguignon solitons in the complex hyperbolic two-plane Grassmannians

By YOUNG JIN SUH (Daegu) and CHANGHWA WOO (Pusan)

Abstract. By using the notion of pseudo-anti-commuting Ricci tensor, we have investigated a Hopf real hypersurface in the complex hyperbolic two-plane Grassmannian $G_2^*(\mathbb{C}^{m+2})$ which admits a pseudo-Ricci–Bourguignon soliton. In addition to this, we have proved that a non-trivial gradient pseudo-Ricci–Bourguignon soliton $(M, Df, \eta, \Omega, \theta, \gamma, g)$ on real hypersurfaces with isometric Reeb flow in the complex hyperbolic two-plane Grassmannian $G_2^*(\mathbb{C}^{m+2})$ does not exist. In the class of contact hypersurfaces in $G_2^*(\mathbb{C}^{m+2})$ except a tube with certain radius $r = \coth^{-1}(\sqrt{3})$ over the totally geodesic and totally real quaternionic hyperbolic space $\mathbb{H}H^n$ in $G_2^*(\mathbb{C}^{m+2})$, $m = 2n$, it has been also proved that there does not exist a non-trivial gradient pseudo-Ricci–Bourguignon soliton in $G_2^*(\mathbb{C}^{m+2})$.

YOUNG JIN SUH
DEPARTMENT OF MATHEMATICS AND RIRC
KYUNGPOOK NATIONAL UNIVERSITY
DAEGU 41566
REPUBLIC OF KOREA

CHANGHWA WOO
DEPARTMENT OF APPLIED MATHEMATICS
PUKYUNG NATIONAL UNIVERSITY
PUSAN 48513
REPUBLIC OF KOREA

Mathematics Subject Classification: Primary: 53C40; Secondary: 53C55.

Key words and phrases: pseudo-Ricci–Bourguignon soliton, non-trivial solution, gradient pseudo-Ricci–Bourguignon soliton, isometric Reeb flow, contact hypersurfaces, complex hyperbolic two-plane Grassmannian.