

On homologies of Klingenberg projective spaces over special commutative local rings

By MAREK JUKL (Olomouc)

Abstract. In this article homologies of Klingenberg projective spaces over commutative local rings of a special type are investigated. Especially sets of invariant points are studied.

1. Introduction

Klingenberg projective spaces (KPS) over a local ring were described by F. MACHALA [6]. Let us have in the KPS P a hyperplane $\mathbf{H}$ and a point $\mathbf{C}$ non-neighbour with $\mathbf{H}$ and let there exist a homology of P such that every point of $\mathbf{H}$ and the point $\mathbf{C}$ are invariant. In the case of projective spaces over fields (which is a special case of a KPS over a local ring) the set of invariant points of the considered homology is just $\mathbf{H} \cup \{\mathbf{C}\}$. However in the case of KPS over a local ring there exist certain invariant points which do not belong to $\mathbf{H} \cup \{\mathbf{C}\}$. If we consider KPS over local rings of the following special type then we may introduce the notion “*degree of neighbourhood*” of two points and by this notion we shall describe the sets of invariant points.

In this paper we shall consider the local commutative ring $\mathbf{A}$ the maximal ideal a of which has the following properties:

- (1) $\exists m \in \mathbb{N} : (a^m = 0) \wedge (a^{m-1} \neq 0)$,
- (2) $a = \eta \mathbf{A}^1$.

Mathematics Subject Classification: 51C05.

Key words and phrases: local ring, $\mathbf{R}$ -space, Klingenberg projective space, neighbour points, homology, invariant point.

¹The factor ring of polynomials $\mathbf{R}[x]/(x^m)$ is an example of the ring of this type.

Throughout the paper the capital $\mathbf{A}$ always denotes the ring of described type.

Clearly, for every $\tau \in \mathbf{A}$, $\tau \neq 0$, there exists a unit τ' and an integer k , $0 \leq k \leq m-1$, such that $\tau = \eta^k \tau'$. The number k is called an *order of* τ . The order of $\tau = 0$ is equal to m .

Let $\mathbf{M}$ be a free finite dimensional module over $\mathbf{A}$. It is well known that all bases of $\mathbf{M}$ have the same number of elements and from every system of generators of $\mathbf{M}$ we may select a basis of $\mathbf{M}$.

Moreover, in our case the module $\mathbf{M}$ has the following properties (proved in [4]):

1. Any linearly independent system can be completed to a basis of $\mathbf{M}$.
2. A submodule of $\mathbf{M}$ is a free module if and only if it is a direct summand of $\mathbf{M}$.

Remark. Free finite dimensional modules over a local ring $\mathbf{R}$ are called $\mathbf{R}$ -spaces (see e.g. [3]) and their direct summands $\mathbf{R}$ -subspaces.

We get that in our case the $\mathbf{A}$ -subspaces of an $\mathbf{A}$ -space $\mathbf{M}$ are just all the free submodules of $\mathbf{M}$.

For $\mathbf{A}$ -subspaces of $\mathbf{M}$ we have (proved in [4]):

3. Let K, L be $\mathbf{A}$ -subspaces of an $\mathbf{A}$ -space $\mathbf{M}$. Then $K + L$ is an $\mathbf{A}$ -subspace if and only if $K \cap L$ is an $\mathbf{A}$ -subspace. In this case the dimensions of the $\mathbf{A}$ -subspaces fulfil the following relation:

$$\dim(K + L) + \dim(K \cap L) = \dim K + \dim L.$$

Lemma 1. *Let $\mathbf{M}$ be an $\mathbf{A}$ -space and let $\bar{\mathbf{M}}$ be a vector space $\mathbf{M}/a\mathbf{M}$. Then the elements $\mathbf{u}_1, \dots, \mathbf{u}_k$ form a linearly independent system in $\mathbf{M}$ if and only if cosets $\bar{\mathbf{u}}_1, \dots, \bar{\mathbf{u}}_k$ form a linearly independent system in $\bar{\mathbf{M}}$.*

PROOF. It follows from the first property that a system $\{\mathbf{u}_1, \dots, \mathbf{u}_k\} \subseteq \mathbf{M}$ is linearly independent iff it may be completed to a basis $\{\mathbf{u}_1, \dots, \mathbf{u}_k, \mathbf{u}_{k+1}, \dots, \mathbf{u}_{n+1}\}$ of $\mathbf{M}$. This is (according to the Theorem I.2 of [3]) equivalent to the fact that the cosets $\{\bar{\mathbf{u}}_1, \dots, \bar{\mathbf{u}}_k, \bar{\mathbf{u}}_{k+1}, \dots, \bar{\mathbf{u}}_{n+1}\}$ form a basis of the vector space $\bar{\mathbf{M}} = \mathbf{M}/a\mathbf{M}$ which means that $\bar{\mathbf{u}}_1, \dots, \bar{\mathbf{u}}_k$ are linearly independent vectors.

Lemma 2. *Let $\mathbf{x}$ as well as $\mathbf{y}$ be a linearly independent element of an $\mathbf{A}$ -space $\mathbf{M}$. If $\xi\mathbf{x} + \vartheta\mathbf{y} = \mathbf{o}$ then ξ, ϑ have the same order.*

PROOF. If $\xi = 0$ then the linear independence of $\mathbf{y}$ implies $\vartheta = 0$. Analogously $\vartheta = 0$ implies $\xi = 0$.

Let $\xi, \vartheta \neq 0$. Then we may write $\xi = \eta^k \xi'$, $\vartheta = \eta^h \vartheta'$ where ξ', ϑ' are units and $0 \leq k, h \leq m-1$. If $k \neq h$ (e.g. $h = k + r$, $r \in \mathbb{N}$) then multiplying the equality $\xi\mathbf{x} + \vartheta\mathbf{y} = \mathbf{o}$ by η^{m-k-r} we obtain $\eta^{m-r}\mathbf{x} = \mathbf{o}$, which contradicts the linear independence of $\mathbf{x}$.

2. Klingenberg projective spaces over local rings

According to [6] we define:

Definition 1. Let $\mathbf{R}$ be a local ring with the maximal ideal r . Let us denote $\mathbf{M} = \mathbf{R}^{n+1}$, $\bar{\mathbf{M}} = \mathbf{M}/r\mathbf{M}$, $\bar{\mathbf{R}} = \mathbf{R}/r$, and let μ be a natural homomorphism $\mathbf{M} \rightarrow \bar{\mathbf{M}}$.

Then an incidence structure P_R such that

- (1) the points are just all submodules $[\mathbf{x}]$ of $\mathbf{M}$ such that $\mu(\mathbf{x})$ is a non-zero element of $\bar{\mathbf{M}}$,
 - (2) the lines are just all submodules $[\mathbf{x}, \mathbf{y}]$ of $\mathbf{M}$ such that $[\mu(\mathbf{x}), \mu(\mathbf{y})]$ is a two-dimensional subspace of $\bar{\mathbf{M}}$,
 - (3) the incidence relation is inclusion,
- is called *an n -dimensional Klingenberg coordinate space over the ring $\mathbf{R}$* .

Two points $P = [\mathbf{p}]$, $Q = [\mathbf{q}]$ such that $[\mu(\mathbf{p})] = [\mu(\mathbf{q})]$ are called *neighbour points* or *neighbours*. In the contrary case they are called *non-neighbours*.

If $X = [\mathbf{x}]$ is a point of P_R , then $\mathbf{x}$ will be called an *arithmetical representative* of X .

A submodule $\mathbf{H}$ of $\mathbf{M}$ is called a *hyperplane* of P_R if $\mathbf{H} = [\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n]$ so that $[\mu(\mathbf{x}_1), \mu(\mathbf{x}_2), \dots, \mu(\mathbf{x}_n)]$ is an n -dimensional subspace of $\bar{\mathbf{M}}$.

We obtain from Definition 1 and Lemma 1 the following corollaries:

Corollary 1. *The points of the Klingenberg space P_A are just all one-dimensional $\mathbf{A}$ -subspaces of $\mathbf{M}$. The lines of the Klingenberg space P_A are just all two-dimensional $\mathbf{A}$ -subspaces of $\mathbf{M}$. The hyperplanes of the Klingenberg space P_A are just all n -dimensional $\mathbf{A}$ -subspaces.*

Corollary 2. *Two points of the Klingenberg space P_A are neighbours if and only if their arithmetical representatives form a linearly dependent subset of $\mathbf{M}$.*

The correctness of the following definition follows from Lemma 2.

Definition 2. Let $X = [\mathbf{x}]$, $Y = [\mathbf{y}]$ be points of the Klingenberg space P_A and let k , $k \geq 0$, be an integer fulfilling the following conditions

- (1) $\eta^k \mathbf{x} \in [\mathbf{y}]$
- (2) $(\eta^{k-1} \mathbf{x} \notin [\mathbf{y}]) \vee (k = 0)$.

Then the integer $r = m - k$ is called a *degree of neighbourhood of the points X and Y* .

Remark. It follows from Corollary 2 that for non-neighbour points we have $r = 0$, for different neighbour points we have $0 < r < m$ and for identical points we have $r = m$.

Lemma 3. *Let X be a point of P_A . Then for every integer r , $0 \leq r \leq m$, there exists at least one point $Y \in P_A$ such that the degree of neighbourhood of the points X, Y is r .*

PROOF. Let $X = [\mathbf{x}]$. Then (according to 1 of Section 1) there exists $\mathbf{z} \in \mathbf{M}$ such that $\mathbf{x}, \mathbf{z}$ form a linearly independent couple. If $r = 0$ then the lemma holds ($Y = [\mathbf{z}]$) as well as in the case $r = m$. Let $0 < r < m$. Considering a point $Y = [\mathbf{y}]$, $\mathbf{y} = \mathbf{x} + \eta^{m-r} \mathbf{z}$, we get $\eta^r \mathbf{y} \in [\mathbf{x}]$ and $\eta^{r-1} \mathbf{x} + \eta^{m-1} \mathbf{z} = \eta^{r-1} \mathbf{y} \notin [\mathbf{x}]$.

3. Homologies of Klingenberg projective spaces

Definition 3. An automorphism of the incidence structure P_A such that there exist an hyperplane $\mathbf{H}$ of invariant points and an invariant point C which is non-neighbour with $\mathbf{H}$ (it means C is not neighbour with any point of $\mathbf{H}$) is called an $(\mathbf{H}, C)$ -homology of P_A and the point C the *center of homology*.

Remark. It follows from Corollary 2 (as $\mathbf{H}$ is an $\mathbf{A}$ -subspace) that $C = [\mathbf{c}]$ is non-neighbour with $\mathbf{H}$ just if $[\mathbf{c}] \cap \mathbf{H} = \{\mathbf{o}\}$.

The following theorem is proved by BACON [2] for the case when P_A is a plane and $\mathbf{A}$ is an arbitrary local ring.

Theorem 1. *Let $\mathbf{H}$ be an hyperplane of the Klingenberg projective space P_A , C a point of P_A non-neighbour with $\mathbf{H}$. If X, Y are points of P_A such that*

- (1) *X and Y are non-neighbour with C and with $\mathbf{H}$*
- (2) *C, X, Y are collinear points*

then there exists exactly one $(C, \mathbf{H})$ -homology of P_A such that $X \mapsto Y$.

PROOF. Since $[\mathbf{c}] \cap \mathbf{H} = \{\mathbf{o}\}$ we get (according to the 3 of Section 1) $[\mathbf{c}] \oplus \mathbf{H} = \mathbf{M}$.

An arbitrary $(C, \mathbf{H})$ -homology F^* on P_A will be induced by an automorphism F of $\mathbf{M}$ in a very natural way:

$$\forall X = [\mathbf{x}] \in P_A : F^*(X) = [F(\mathbf{x})].$$

As $F^*|_{\mathbf{H}}$ is an identity mapping we may infer that there exists a unit $\lambda \in \mathbf{A}$ such that $F|_{\mathbf{H}} = \lambda \cdot id$. Without loss of generality we can assume that $F|_{\mathbf{H}} = id$.

Any $\mathbf{x} \in \mathbf{M}$ may be uniquely expressed in the form

$$(1) \quad \mathbf{x} = \mathbf{x}' + \xi \mathbf{c}, \quad \mathbf{x}' \in \mathbf{H}, \quad \xi \in \mathbf{A}.$$

The automorphism F is given by

$$(2) \quad \mathbf{x} = \mathbf{x}' + \xi \mathbf{c} \mapsto F(\mathbf{x}) = \mathbf{x}' + \alpha \xi \mathbf{c},$$

where α is a unit.

Now $\mathbf{x}$ is linearly independent (equivalently, $\mathbf{x}$ represents a point $[\mathbf{x}]$ of P_A) if and only if $\mathbf{x}'$ is linearly independent or $\xi \notin a$.

Moreover $\mathbf{x}$ represents a point non-neighbour with C and with $\mathbf{H}$ if and only if $\mathbf{x}'$ is linearly independent and $\xi \notin a$.

Considering $Y = [\mathbf{y}]$, $Y \in XC$, we get $\mathbf{y} = \gamma \mathbf{x} + \sigma \mathbf{c}$. As Y, C are not neighbour points, γ is a unit and the arithmetical representative $\mathbf{y}$ may be expressed by

$$\mathbf{y} = \mathbf{x} + \sigma \mathbf{c}.$$

Using the expression (1) of $\mathbf{x}$ we obtain from this

$$(3) \quad \mathbf{y} = \mathbf{x}' + (\xi + \sigma) \mathbf{c}.$$

The automorphism F^* maps X to Y just if $F(\mathbf{x}) = \varepsilon \mathbf{y}$, where ε is a unit. Using (2) and (3) we get

$$(1 - \varepsilon)\mathbf{x}' + (\alpha\xi - (\xi + \sigma)\varepsilon) = \mathbf{c} = \mathbf{o}.$$

As $[\mathbf{c}] \cap \mathbf{H}$ is trivial, $\varepsilon = 1$, $\alpha\xi = -(\xi + \sigma)\varepsilon = 0$, hence $\alpha = \xi^{-1}(\xi + \sigma)$. This means that α as well as the homology F^* is determined uniquely.

Remark. Every $(C, \mathbf{H})$ -homology of P_A , $C = [\mathbf{c}]$ may be expressed by the following formula:

$$(4) \quad X = [\mathbf{x}' + \xi \mathbf{c}] \mapsto F^*(X) = [\mathbf{x}' + \alpha \xi \mathbf{c}], \quad \mathbf{x}' \in \mathbf{H}, \alpha \notin a.$$

The element α will be called the *coefficient of the homology* F^* .

Proposition 1. *Let F^* be a $(C, \mathbf{H})$ -homology of P_A and α the coefficient of F^* . If $(1 - \alpha)$ has order r then*

- (1) *for any point $X \in P_A$, X and $F^*(X)$ are neighbours of order at least r .*
- (2) *there exists $X \in P_A$ such that X and $F^*(X)$ are neighbours just of degree r .*

PROOF. As usual let $C = [\mathbf{c}]$. Let F be the automorphism of $\mathbf{M}$ inducing F^* and let F be given by (2). Suppose that $(1 - \alpha) = \alpha_0 \eta^r$, where α_0 is a unit. Using (2) we have

$$F(\mathbf{x}) = (\mathbf{x} - \xi \mathbf{c}) + \xi \alpha \mathbf{c} = \mathbf{x} + \xi = (\alpha - 1)\mathbf{c} = \mathbf{x} - \alpha_0 \eta^r \xi \mathbf{c}.$$

Therefore $\eta^{m-r} F(\mathbf{x}) = \eta^{m-r} \mathbf{x}$. This means that the degree of neighbourhood of the points $X = [\mathbf{x}]$ and $F^*(X)$ is (at least) r .

If ξ is a unit and $[\mathbf{x}]$, C are not neighbours then

$$\eta^{m-r-1} F(\mathbf{x}) = \eta^{m-r-1} \mathbf{x} - (\eta^{m-1} \xi \alpha_0) \mathbf{c} \text{ where } \eta^{m-1} \xi \alpha_0 \neq 0.$$

Thus $\eta^{m-r-1} F(\mathbf{x}) \notin [\mathbf{c}]$ which means that X and $F^*(X)$ are neighbours of precisely degree r .

Remark. In particular, if $1 - \alpha$ has order m (i.e. $\alpha = 1$) the F^* is an identity and all the $X, F^*(X)$ are neighbours of degree (at least) m , i.e. identical.

If $1 - \alpha$ is a unit then there exists X such that $X, F^*(X)$ are neighbours of degree zero i.e. non-neighbours.

Proposition 2. *Let F^* be a $(C, \mathbf{H})$ -homology of P_A and α the coefficient of F^* . Let X be an arbitrary point non-neighbour with C as well as $\mathbf{H}$ and let the degree of neighbourhood of X and $F^*(X)$ be r . Then the following hold*

- (1) $(1 - \alpha)$ has order r .
- (2) *If Y is a point non-neighbour with C as well as with $\mathbf{H}$ then Y and $F^*(Y)$ are neighbours of degree r .*

PROOF. Let F^* be given by the formula (4) and let F be an automorphism of $\mathbf{M}$ given by (2). If X and $F^*(X)$ are neighbours of order r then (according to Lemma 2)

$$(5) \quad \eta^{m-r} F(\mathbf{x}) = \eta^{m-r} \varepsilon \mathbf{x}, \quad \varepsilon \notin a.$$

It follows from (2) that $F(\mathbf{x}) = \mathbf{x} + \xi(\alpha - 1)\mathbf{c}$. Using this and (5) we obtain

$$\eta^{m-r}(1 - \varepsilon)\mathbf{x} = \eta^{m-r}\xi(1 - \alpha)\mathbf{c}.$$

Because X, C are non-neighbours we get

$$(6) \quad \eta^{m-r}(1 - \varepsilon)\mathbf{x} = \eta^{m-r}\xi(1 - \alpha)\mathbf{c} = \mathbf{o}.$$

Since X is non-neighbour with $\mathbf{H}$ we have that ξ is a unit. Thus $(1 - \alpha)$ has order at least r . If the order of $(1 - \alpha)$ is greater than r then (according to the previous proposition) we get that U and $F^*(U)$ are neighbours of degree at least $r + 1$ for every point U and this contradicts our assumption. Therefore the order of $(1 - \alpha)$ is (precisely) r .

Suppose there exists Y (non-neighbour with $\mathbf{H}$ and with C) such that $Y, F^*(Y)$ are neighbours of degree $r + 1$. This implies (by (6)) that $(1 - \alpha)$ has order (at least) $r + 1$ which is not possible.

Theorem 2. *Let F^* be a $(C, \mathbf{H})$ -homology of P_A , α the coefficient of F^* , r the order of $(1 - \alpha)$ and let X be a point of P_A . Then X is F^* -invariant if and only if it is neighbour of degree at least $m - r$ with C or some point of $\mathbf{H}$.*

PROOF. Let us remark that $\mathbf{H} \oplus [\mathbf{c}] = \mathbf{M}$. Let the homology F^* be given by the formula (4).

In case $(1 - \alpha)$ is of order m (i.e. $\alpha = 1$) we get that F^* is an identity and the proposition holds. Now suppose $r \leq m - 1$.

Let $X = [\mathbf{x}]$ be an invariant point of F^* and let F be an automorphism of $\mathbf{M}$ given by (2). Then $F(\mathbf{x}) = \varepsilon\mathbf{x}$, $\varepsilon \notin a$. Using this and (2) we get

$$(1 - \varepsilon)\mathbf{x}' = \xi(\varepsilon - \alpha)\mathbf{c}.$$

As C is not neighbour with $\mathbf{H}$ this yields

$$(7) \quad (1 - \varepsilon)\mathbf{x}' = \xi(\varepsilon - \alpha)\mathbf{c} = \mathbf{o}.$$

Because $\mathbf{x}$ is linearly independent $\mathbf{x}'$ is also linearly independent or ξ is a unit.

(a) If $\mathbf{x}'$ is linearly independent then (by (7)) $\varepsilon = 1$, thus $\xi(1 - \alpha) = 0$. Supposing $(1 - \alpha)$ has order r we have $\xi = \xi_0\eta^{m-r}$. Now we obtain $\mathbf{x} = \mathbf{x}' + \xi_0\eta^{m-r}\mathbf{c}$ and thus $\eta^r\mathbf{x} = \eta^r\mathbf{x}' \in \mathbf{H}$. This means that the degree of neighbourhood of X and $\mathbf{H}$ is at least $m - r$.

(b) If ξ is a unit then (by (7)) $\varepsilon = \alpha$ which implies that $(1 - \varepsilon) = (1 - \alpha)$ has order r . As $(1 - \varepsilon)\mathbf{x}' = \mathbf{o}$, $\mathbf{x}'$ may be written as $\mathbf{x}' = \eta^{m-r}\mathbf{y}$, $\mathbf{y} \in \mathbf{H}$. Now we get $\mathbf{x} = \eta^{m-r}\mathbf{y} + \xi\mathbf{c}$ and therefore $\eta^r\mathbf{x} = \eta^r\xi\mathbf{c}$, $\xi \notin a$. This means that the degree of neighbourhood of X and C is at least $m - r$.

Now, suppose that degree of neighbourhood of $X = [\mathbf{x}]$ and $\mathbf{H}$ is at least $m - r$. This implies that $\eta^r\mathbf{x} \in \mathbf{H}$. As $\eta^r\mathbf{x}' + \eta^r\xi\mathbf{c} = \eta^r\mathbf{x} \in \mathbf{H}$ we have $\eta^r\xi = 0$ and thus $\xi = \xi_0\eta^{m-r}$. Supposing $1 - \alpha = \alpha_0\eta^r$ we get

$$\begin{aligned} F(\mathbf{x}) &= \mathbf{x}' + \xi_0\alpha\eta^{m-r}\mathbf{c} = \mathbf{x}' + \xi_0\eta^{m-r}(1 - \alpha_0\eta^r)\mathbf{c} \\ &= \mathbf{x}' + \xi\mathbf{c} - \xi_0\alpha_0\eta^m\mathbf{c} = \mathbf{x}, \end{aligned}$$

i.e. X is an invariant point.

Let us suppose that C and $X = [\mathbf{x}]$ are neighbours of degree at least $m - r$. Then $\eta^r\mathbf{x} \in [\mathbf{c}]$ which means that $\eta^r\mathbf{x}' + \eta^r\xi\mathbf{c} = \eta^r\mathbf{x} \in [\mathbf{c}]$. This gives $\eta^r\mathbf{x}' = \mathbf{o}$ i.e. $\mathbf{x}' = \eta^{m-r}\mathbf{y}$, $\mathbf{y} \in \mathbf{H}$.

Therefore $\mathbf{x} = \eta^{m-r}\mathbf{y} + \xi\mathbf{c}$ and $F(\mathbf{x}) = \eta^{m-r}\mathbf{y} + \xi\alpha\mathbf{c} = (1 + \alpha_0\eta^r)\eta^{m-r}\mathbf{y} + \xi\alpha\mathbf{c} = \alpha\mathbf{x}' + \alpha\xi\mathbf{c} = \alpha\mathbf{x}$. This means that X is an invariant point.

Remark. By this theorem (in view of Lemma 3) the set of F^* -invariant points uniquely determines the order of $1 - \alpha$.

The following theorem is a consequence of Proposition 1, 2 and of Theorem 2:

Theorem 3. *Let F^* be a $(C, \mathbf{H})$ -homology of P_A and let X be an arbitrary point non-neighbour with $\mathbf{H}$ as well as with C . Then the points X , $F^*(X)$ are neighbours of degree r if and only if the F^* -invariant points are just all the points which are neighbours with C or with $\mathbf{H}$ of degree at least $m - r$.*

References

- [1] F. W. ANDERSON and F. K. FULLER, Rings and categories of modules, *Springer-Verlag, New York*, 1973.
- [2] P. Y. BACON, An introduction to Klingenberg planes, *Florida*, 1976.
- [3] B. R. McDONALD, Geometric algebra over local rings, *Pure and Applied Mathematics, New York*, 1976.
- [4] M. JUKL, Grassmann formula for certain type of modules, *Acta UP Olomouc, Fac. rer. nat., Mathematica* **34** (1995), 69–74.
- [5] W. KLINGENBERG, Projektive Geometrien mit Homomorphismus, *Math. Annalen* **132** (1956), 180–200.
- [6] F. MACHALA, Fundamentalsätze der projektiven Geometrie mit Homomorphismus, *Rozprawy SAV, ada matem.a prodnch vd, 90, seit 5, Praha, Academia*, 1980.

MAREK JUKL
DEPARTMENT OF ALGEBRA AND GEOMETRY
FACULTY OF SCIENCE
PALACKY UNIVERSITY
OLOMOUC TOMKOVA 40
CZECH REPUBLIC

E-mail: jukl@risc.upol.cz

*(Received November 17, 1997; revised February 27, 1998;
accepted December 8, 1998)*