

Integer points on a family of elliptic curves

By ANDREJ DUJELLA (Zagreb) and ATTILA PETHŐ (Debrecen)

*Dedicated to Professor Kálmán Győry
on the occasion of his 60th birthday*

Abstract. Let the sequence (c_k) be given by the recursion

$$c_0 = 0, \quad c_1 = 8, \quad c_{k+2} = 14c_{k+1} - c_k + 8, \quad k \geq 0.$$

Let the elliptic curve E_k be defined by the equation $y^2 = (x+1)(3x+1)(c_kx+1)$. We prove in this paper that if the rank of $E_k(\mathbb{Q})$ is equal to two, or $k \leq 40$, with the possible exceptions $k = 23$ and $k = 37$, then all integer points on E_k are given by

$$(x, y) \in \{(-1, 0), \quad (0, \pm 1), \quad (c_{k-1}, \pm s_{k-1}t_{k-1}(2c_k - s_k t_k)), \\ (c_{k+1}, \pm s_{k+1}t_{k+1}(2c_k + s_k t_k))\}.$$

where $c_k + 1 = s_k^2$ and $3c_k + 1 = t_k^2$.

1. Introduction

A set D of m positive integers is called a *Diophantine m -tuple* if the product of any two distinct elements of D increased by 1 is a perfect square. The first example of a Diophantine quadruple – $\{1, 3, 8, 120\}$ – was found by Fermat (see [6, p. 517]). In 1969, BAKER and DAVENPORT [2] proved that if d is a positive integer such that $\{1, 3, 8, d\}$ is a Diophantine quadruple, then d has to be 120.

Mathematics Subject Classification: 11D09, 11D25, 11G05, 11Y50.

Key words and phrases: systems of Pell equations, Diophantine tuples, elliptic curves. Research of the second author partially supported by Hungarian National Foundation for Scientific Research Grants No. 25157 and 29330.

Recently, in [9], we generalized this result to all Diophantine triples of the form $\{1, 3, c\}$. The fact that $\{1, 3, c\}$ is a Diophantine triple implies that $c = c_k$ for some positive integer k , where the sequence (c_k) is given by

$$c_0 = 0, \quad c_1 = 8, \quad c_{k+2} = 14c_{k+1} - c_k + 8, \quad k \geq 0.$$

Let $c_k + 1 = s_k^2$, $3c_k + 1 = t_k^2$ with positive integers s_k, t_k . It is easy to check that

$$c_{k \pm 1}c_k + 1 = (2c_k \pm s_k t_k)^2.$$

The main result of [9] is the following theorem.

Theorem 1. *Let k be a positive integer. If d is an integer which satisfies the system of equations*

$$(1) \quad d + 1 = x_1^2, \quad 3d + 1 = x_2^2, \quad c_k d + 1 = x_3^2,$$

then $d \in \{0, c_{k-1}, c_{k+1}\}$.

Eliminating d from the system (1) we obtain the following system of Pellian equations

$$(2) \quad x_3^2 - c_k x_1^2 = 1 - c_k$$

$$(3) \quad 3x_3^2 - c_k x_2^2 = 3 - c_k.$$

We used the theory of Pellian equations and some congruence relations to reformulate the system (2) and (3) to four equations of the form $v_m = w_n$, where (v_m) and (w_n) are binary recursive sequences. After that, a comparison of the upper bound for the solutions obtained from the theorem of BAKER and WÜSTHOLZ [3] with the lower bound obtained from the congruence condition modulo c_k^2 finished the proof for $k \geq 76$. The statement for $1 \leq k \leq 75$ was proved by a variant of the reduction procedure due to BAKER and DAVENPORT [2].

Similar results are proved in [7] and [8] for Diophantine triples of the form $\{k-1, k+1, 4k\}$ and $\{F_{2k}, F_{2k+2}, F_{2k+4}\}$. In the second triple F_n denotes the n -th Fibonacci number.

It is clear that every solution $(d, x_1, x_2, x_3) \in \mathbb{Z}^4$ of (1) induce an integer point on the elliptic curve

$$(4) \quad E_k : \quad y^2 = (x+1)(3x+1)(c_k x+1),$$

with $y = x_1 x_2 x_3$ and $x = d$. The purpose of the present paper is to prove that the converse of this statement is true, provided the rank of $E_k(\mathbb{Q})$ is equal to 2. As we will see in Proposition 2, for all $k \geq 2$ the rank of $E_k(\mathbb{Q})$ is always ≥ 2 . Our main result is

Theorem 2. *Let k be a positive integer. If $\text{rank}(E_k(\mathbb{Q})) = 2$ or $k \leq 40$, with the possible exceptions $k = 23$ and $k = 37$, then all integer points on E_k are given by*

$$(x, y) \in \{(-1, 0), (0, \pm 1), (c_{k-1}, \pm s_{k-1}t_{k-1}(2c_k - s_k t_k)), \\ (c_{k+1}, \pm s_{k+1}t_{k+1}(2c_k + s_k t_k))\}.$$

2. Torsion group

Under the substitution $x \leftrightarrow 3c_k x$, $y \leftrightarrow 3c_k y$ the curve E_k is transformed into the following Weierstraß form

$$E'_k : y^2 = x^3 + (4c_k + 3)x^2 + (3c_k^2 + 12c_k)x + 9c_k^2 = (x + 3c_k)(x + c_k)(x + 3).$$

There are three rational points on E'_k of order 2, namely

$$A_k = (-3c_k, 0), \quad B_k = (-c_k, 0), \quad C_k = (-3, 0),$$

and also other two, more or less obvious, rational points on E'_k , namely

$$P_k = (0, 3c_k), \quad R_k = (s_k t_k + 2s_k + 2t_k + 1, (s_k + t_k)(s_k + 2)(t_k + 2)).$$

Note that if $k = 1$, then $R_1 = C_1 - P_1$.

Lemma 1. $E'_k(\mathbb{Q})_{\text{tors}} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

PROOF. From [17, Main Theorem 1] it follows immediately that $E'_k(\mathbb{Q})_{\text{tors}} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ or $E'_k(\mathbb{Q})_{\text{tors}} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$, and the later is possible iff there exist integers α and β such that $\frac{\alpha}{\beta} \notin \{-2, -1, -\frac{1}{2}, 0, 1\}$ and

$$c_k - 3 = \alpha^4 + 2\alpha^3\beta, \quad 3c_k - 3 = 2\alpha\beta^3 + \beta^4.$$

Now, we have

$$(5) \quad 4c_k - 6 = (\alpha^2 + \alpha\beta + \beta^2)^2 - 3\alpha^2\beta^2.$$

Since c_k is even, the left hand side of (5) is $\equiv 2 \pmod{8}$. If α and β are both even then the right hand side of (5) is divisible by 8, and if α and β are both odd then the right hand side of (5) is $\equiv 6 \pmod{8}$, a contradiction. Hence, $E'_k(\mathbb{Q})_{\text{tors}} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. $\square$

3. The independence of P_k and R_k

In this section we will often use the following 2-descent proposition (see [12, 4.1, p. 37]).

Proposition 1. *Let $P = (x', y')$ be a $\mathbb{Q}$ -rational point on E , an elliptic curve over $\mathbb{Q}$ given by the equation*

$$y^2 = (x - \alpha)(x - \beta)(x - \gamma),$$

where $\alpha, \beta, \gamma \in \mathbb{Q}$. Then there exists a $\mathbb{Q}$ -rational point $Q = (x, y)$ on E such that $2Q = P$ iff $x' - \alpha, x' - \beta, x' - \gamma$ are all $\mathbb{Q}$ -rational squares.

Lemma 2. $P_k, P_k + A_k, P_k + B_k, P_k + C_k \notin 2E'_k(\mathbb{Q})$.

PROOF. We have:

$$\begin{aligned} P_k + A_k &= (-c_k - 2, -2c_k + 2), \\ P_k + B_k &= (-3c_k + 6, 6c_k - 18), \\ P_k + C_k &= (c_k^2 - 4c_k, -c_k^3 + 4c_k^2 - 3c_k). \end{aligned}$$

It follows immediately from Proposition 1 that $P_k, P_k + A_k, P_k + B_k \notin 2E'_k(\mathbb{Q})$. If $P_k + C_k \in 2E'_k(\mathbb{Q})$, then $c_k^2 - c_k = \square$, which is impossible. $\square$

Lemma 3. $R_k, R_k + A_k, R_k + B_k, R_k + C_k \notin 2E'_k(\mathbb{Q})$.

PROOF. We have:

$$\begin{aligned} R_k &= (s_k t_k + 2s_k + 2t_k + 1, (t_k + s_k)(s_k + 2)(t_k + 2)), \\ R_k + A_k &= (2s_k - 2t_k - s_k t_k + 1, (s_k - t_k)(s_k + 2)(t_k - 2)), \\ R_k + B_k &= (2t_k - 2s_k - s_k t_k + 1, (t_k - s_k)(s_k - 2)(t_k + 2)), \\ R_k + C_k &= (s_k t_k - 2s_k - 2t_k + 1, (t_k + s_k)(2 - s_k)(t_k - 2)). \end{aligned}$$

Since $2s_k - 2t_k - s_k t_k + 4 = (s_k + 2)(2 - t_k) < 0$ and $2t_k - 2s_k - s_k t_k + 4 = (t_k + 2)(2 - s_k) < 0$, we have $R_k + A_k, R_k + B_k \notin 2E'_k(\mathbb{Q})$.

If $R_k \in 2E'_k(\mathbb{Q})$, then $(t_k + s_k)(t_k + 2) = \square$ and $(t_k + s_k)(s_k + 2) = \square$. Let $d = \gcd(t_k + s_k, t_k + 2, s_k + 2)$. Then d divides $(t_k + 2) + (s_k + 2) - (t_k + s_k) = 4$, and since s_k and t_k are odd, we conclude that $d = 1$. Hence, we have

$$(6) \quad t_k + s_k = \square, \quad t_k + 2 = \square, \quad s_k + 2 = \square.$$

Consider the sequence $(t_k + s_k)_{k \in \mathbb{N}}$. It follows easily by induction that $t_k + s_k = 2a_{k+1}$, where

$$(7) \quad a_0 = 0, \quad a_1 = 1, \quad a_{k+2} = 4a_{k+1} - a_k, \quad k \geq 0.$$

Thus, (6) implies $a_{k+1} = 2\Box$, and this is impossible by a theorem of MIGNOTTE and PETHŐ [14] (see also [16]) which says that $a_k = \Box, 2\Box, 3\Box$ or $6\Box$ implies $k \leq 3$.

If $R_k + C_k \in 2E'_k(\mathbb{Q})$, then $(t_k + s_k)(t_k - 2) = \Box$ and $(t_k + s_k)(s_k - 2) = \Box$. This implies $t_k + s_k = \Box$ and we obtain a contradiction as above. $\square$

Lemma 4. *If $k \geq 2$, then $R_k + P_k, R_k + P_k + A_k, R_k + P_k + B_k, R_k + P_k + C_k \notin 2E'_k(\mathbb{Q})$.*

PROOF. As in the proof of Lemmas 2 and 3, we use Proposition 1.

If $R_k + P_k + A_k \in 2E'_k(\mathbb{Q})$ then $0 > c_k(s_k + 2)(s_k - t_k) = \Box$, and if $R_k + P_k + B_k \in 2E'_k(\mathbb{Q})$ then $0 > c_k(s_k - 2)(s_k - t_k) = \Box$. Hence, $R_k + P_k + A_k, R_k + P_k + B_k \notin 2E'_k(\mathbb{Q})$.

If $R_k + P_k \in 2E'_k(\mathbb{Q})$ then

$$(8) \quad \begin{aligned} 3c_k(t_k + s_k)(t_k + 2) &= \Box, \\ c_k(t_k + s_k)(s_k + 2) &= \Box, \\ 3(s_k + 2)(t_k + 2) &= \Box. \end{aligned}$$

Substituting $2c_k = (t_k + s_k)(t_k - s_k)$ in (8) we obtain

$$\begin{aligned} (t_k - s_k)(t_k + 2) &= 6\Box, \\ (t_k - s_k)(s_k + 2) &= 2\Box, \\ (s_k + 2)(t_k + 2) &= 3\Box. \end{aligned}$$

Let $d = \gcd(s_k + 2, t_k + 2)$. Then the relation $t_k^2 - 3s_k^2 = -2$ implies $d|6$. Since $t_k + 2$ is odd, we have $d \in \{1, 3\}$. Hence we obtain

$$(9) \quad t_k - s_k = 6\Box \quad \text{or} \quad t_k - s_k = 2\Box.$$

But $t_k - s_k = 2a_k$, where (a_k) is defined by (7). Thus (9) implies $a_k = \Box$ or $3\Box$. According to [14], this is possible only if $k = 2$. But $(s_2, t_2) = (11, 19)$ and $(s_2 + 2)(t_2 + 2) \neq 3\Box$.

If $R_k + P_k + C_k \in 2E'_k(\mathbb{Q})$ then

$$3c_k(t_k + s_k)(t_k - 2) = \square,$$

$$c_k(t_k + s_k)(s_k - 2) = \square,$$

$$3(s_k - 2)(t_k - 2) = \square.$$

Arguing as before, we obtain

$$(t_k - s_k)(t_k - 2) = 6\square,$$

$$(t_k - s_k)(s_k - 2) = 2\square,$$

$$(s_k - 2)(t_k - 2) = 3\square,$$

and conclude that

$$t_k - s_k = 6\square \quad \text{or} \quad t_k - s_k = 2\square.$$

As we have already seen, it is possible only for $(s_2, t_2) = (11, 19)$, but then $(s_2 - 2)(t_2 - 2) \neq 3\square$. $\square$

Proposition 2. *If $k \geq 2$, then the points P_k and R_k generate a subgroup of rank 2 in $E'_k(\mathbb{Q})/E'_k(\mathbb{Q})_{\text{tors}}$.*

PROOF. We have to prove that $mP_k + nR_k \in E'_k(\mathbb{Q})_{\text{tors}}$, $m, n \in \mathbb{Z}$, implies $m = n = 0$.

Assume $mP_k + nR_k = T \in E'_k(\mathbb{Q})_{\text{tors}} = \{\mathcal{O}, A_k, B_k, C_k\}$ with $(m, n) \neq (0, 0)$. If m and n are not both even, then $T \equiv P_k, R_k$ or $P_k + R_k \pmod{2E'_k(\mathbb{Q})}$, which is impossible by Lemmas 2, 3 and 4. Hence, m and n are even, say $m = 2m_1$, $n = 2n_1$, and since by Lemma 1 $A_k, B_k, C_k \notin 2E'_k(\mathbb{Q})$,

$$2m_1P_k + 2n_1Q_k = \mathcal{O}.$$

Thus we obtain $m_1P_k + n_1R_k \in E'_k(\mathbb{Q})_{\text{tors}}$. Arguing as above, we obtain that m_1 and n_1 are even, and continuing this process we finally conclude that $m = n = 0$. $\square$

4. Proof of Theorem 2 ($\text{rank}(E_k(\mathbb{Q})) = 2$)

Let $E'_k(\mathbb{Q})/E'_k(\mathbb{Q})_{\text{tors}} = \langle U, V \rangle$ and $X \in E'_k(\mathbb{Q})$. Then there exist integers m, n and a torsion point T such that $X = mU + nV + T$. Also $P_k = m_P U + n_P V + T_P$, $R_k = m_R U + n_R V + T_R$ with integers m_P, n_P, m_R, n_R and with $T_P, T_R \in E'_k(\mathbb{Q})_{\text{tors}}$. Let $\mathcal{U} = \{\mathcal{O}, U, V, U + V\}$. There exist $U_1, U_2 \in \mathcal{U}$, $T_1, T_2 \in E'_k(\mathbb{Q})_{\text{tors}}$ such that $P_k \equiv U_1 + T_1 \pmod{2E'_k(\mathbb{Q})}$, $R_k \equiv U_2 + T_2 \pmod{2E'_k(\mathbb{Q})}$. Let $U_3 \in \mathcal{U}$ such that $U_3 \equiv U_1 + U_2 \pmod{2E'_k(\mathbb{Q})}$. Then $P_k + R_k \equiv U_3 + (T_1 + T_2) \pmod{2E'_k(\mathbb{Q})}$. Now Lemmas 2, 3 and 4 imply that $U_1, U_2, U_3 \neq \mathcal{O}$ and accordingly $\{U_1, U_2, U_3\} = \{U, V, U + V\}$. Therefore $X \equiv X_1 \pmod{2E'_k(\mathbb{Q})}$, where

$$\begin{aligned} X_1 \in \mathcal{S} = \{ & \mathcal{O}, A_k, B_k, C_k, P_k, P_k + A_k, P_k + B_k, P_k + C_k, \\ & R_k, R_k + A_k, R_k + B_k, R_k + C_k, R_k + P_k, \\ & R_k + P_k + A_k, R_k + P_k + B_k, R_k + P_k + C_k \}. \end{aligned}$$

Let $\{a, b, c\} = \{3, c_k, 3c_k\}$. By [13, 4.6, p. 89], the function $\varphi : E'_k(\mathbb{Q}) \rightarrow \mathbb{Q}^*/\mathbb{Q}^{*2}$ defined by

$$\varphi(X) = \begin{cases} (x+a)\mathbb{Q}^{*2}, & \text{if } X = (x, y) \neq \mathcal{O}, (-a, 0), \\ (b-a)(c-a)\mathbb{Q}^{*2}, & \text{if } X = (-a, 0), \\ \mathbb{Q}^{*2}, & \text{if } X = \mathcal{O} \end{cases}$$

is a group homomorphism.

This fact and Theorem 1 imply that it is sufficient to prove that for all $X_1 \in \mathcal{S} \setminus P_k$, $X_1 = (3c_k u, 3c_k v)$, the system

$$(11) \quad x + 1 = \alpha \square, \quad 3x + 1 = \beta \square, \quad c_k x + 1 = \gamma \square$$

has no integer solution, where $\square$ denotes a square of a rational number, and α, β, γ are defined by $u + 1 = \alpha$, $3u + 1 = \beta$, $c_k u + 1 = \gamma$ if all those numbers are $\neq 0$, and if e.g. $u + 1 = 0$ then we choose $\alpha = \beta\gamma$ (so that $\alpha\beta\gamma = \square$). Note that for $X_1 = P_k$ we obtain the system $x + 1 = \square$, $3x + 1 = \square$, $c_k x + 1 = \square$, which is completely solved in Theorem 1.

For $X_1 \in \{A_k, B_k, P_k + A_k, P_k + B_k, R_k + A_k, R_k + B_k, R_k + P_k + A_k, R_k + P_k + B_k\}$ exactly two of the numbers α, β, γ are negative and thus the system (11) has no integer solution.

The rest of the proof falls naturally into 7 parts. By a' we will denote the square free part of an integer a .

1) $X_1 = \mathcal{O}$:

We have

$$(12) \quad x + 1 = 3c_k \square, \quad 3x + 1 = c_k \square, \quad c_k x + 1 = 3\square.$$

From the second equation in (12) we see that $3 \nmid c'_k$ and thus the first and second equations imply that c'_k divides $3x + 1$ and $x + 1$. Accordingly, $c'_k \mid 3(x + 1) - (3x + 1) = 2$ and we conclude that $c'_k = 1$ or 2 . Hence,

$$c_k = \square, \quad \text{or} \quad c_k = 2\square.$$

However, $c_k = s_k^2 - 1 = \square$ is obviously impossible, while $c_k = 2w^2$ leads to the system of Pellian equations

$$s_k^2 - 2w^2 = 1, \quad t_k^2 - 6w^2 = 1.$$

This system is solved by ANGLIN [1], and the only positive solution is $(s_k, t_k, w) = (3, 5, 2)$ which corresponds to $c_k = c_1 = 8$, contradicting our assumption that $k \geq 2$. (Note that for $c_1 = 8$ there is also no solution because in this case the first and the third equations in (12) imply $3 \mid 7$.)

2) $X_1 = C_k$:

We have

$$\begin{aligned} x + 1 &= c_k(c_k - 1)\square, \\ 3x + 1 &= c_k(c_k - 3)\square, \\ c_k x + 1 &= (c_k - 1)(c_k - 3)\square. \end{aligned}$$

If $3 \nmid c_k$ then, as in 1), we obtain $c'_k = 1$ or 2 , and $c_k = \square$ or $2\square$, which is impossible.

If $c_k = 3e_k$ then e'_k divides $3x + 1$ and $3x + 3$ and thus $e'_k = 1$ or 2 . Hence,

$$c_k = 3\square, \quad \text{or} \quad c_k = 6\square.$$

The relation $c_k = 3\square$ is impossible since it implies $t_k^2 - 1 = 9\square$, while $c_k = 6w^2$ leads to the system of Pellian equations

$$s_k^2 - 6w^2 = 1, \quad t_k^2 - 18w^2 = 1$$

which has no positive solution according to [1].

3) $X_1 = P_k + C_k :$

We have

$$\begin{aligned} x + 1 &= 3(c_k - 1)\square, \\ 3x + 1 &= (c_k - 3)\square, \\ c_k x + 1 &= 3(c_k - 1)(c_k - 3)\square. \end{aligned}$$

Since $c_k = s_k^2 - 1$, we see that $c_k \not\equiv 1 \pmod{3}$, and thus $x \equiv -1 \pmod{3}$. From the second equation we have that $(c_k - 3)'$ is not divisible by 3, and then the third equation gives $c_k x + 1 \equiv 0 \pmod{3}$. This implies $c_k \equiv 1 \pmod{3}$, a contradiction.

4) $X_1 = R_k :$

We have

$$\begin{aligned} x + 1 &= 6(t_k - s_k)(t_k + 2)\square, \\ 3x + 1 &= 2(t_k - s_k)(s_k + 2)\square, \\ c_k x + 1 &= 3(s_k + 2)(t_k + 2)\square. \end{aligned}$$

From the relation $t_k^2 - 3s_k^2 = -2$ it follows that $\gcd(t_k - s_k, s_k + 2) = \gcd(t_k - s_k, t_k + 2) = 1$ or 3 .

If $3 \nmid t_k - s_k$ then $[2(t_k - s_k)]'$ divides $x + 1$ and $3x + 1$, and thus $[2(t_k - s_k)]' = 1$ or 2 . Accordingly,

$$t_k - s_k = 2\square \quad \text{or} \quad t_k - s_k = \square.$$

As we have already seen in the proof of Lemma 4, this implies

$$a_k = \square \quad \text{or} \quad a_k = 2\square,$$

and [14] implies again that $k = 2$. Now we obtain $120x + 1 = 91\square$, which is impossible modulo 4.

If $t_k - s_k = 3z_k$ then $(2z_k)'$ divides $x + 1$ and $9x + 3$. Hence $(2z_k)'$ divides 6, which implies $a_k = \square, 2\square, 3\square$ or $6\square$, and this is possible only if $k = 2$. But for $k = 2$, $t_k - s_k = 8 \not\equiv 0 \pmod{3}$.

5) $X_1 = R_k + C_k :$

We have

$$\begin{aligned} x + 1 &= 6(t_k - s_k)(t_k - 2)\square, & 3x + 1 &= 2(t_k - s_k)(s_k - 2)\square, \\ c_k x + 1 &= 3(s_k - 2)(t_k - 2)\square. \end{aligned}$$

This case is completely analogous to the case 4).

6) $X_1 = R_k + P_k :$

We have

$$\begin{aligned} x + 1 &= (t_k + s_k)(t_k + 2)\square, & 3x + 1 &= (t_k + s_k)(s_k + 2)\square, \\ c_k x + 1 &= (s_k + 2)(t_k + 2)\square. \end{aligned}$$

As in 4), we obtain that if $3 \nmid t_k + s_k$ then $(t_k + s_k)'$ divides 2, and if $t_k + s_k = 3z_k$ then z_k' divides 6. Hence, we have $a_{k+1} = \square, 2\square, 3\square$ or $6\square$, which is impossible for $k \geq 2$.

7) $X_1 = R_k + P_k + C_k :$

We have

$$\begin{aligned} x + 1 &= (t_k + s_k)(t_k - 2)\square, & 3x + 1 &= (t_k + s_k)(s_k - 2)\square, \\ c_k x + 1 &= (s_k - 2)(t_k - 2)\square. \end{aligned}$$

This case is completely analogous to the case 6). $\square$

Remark 1. It is easy to check that $\text{rank}(E_1(\mathbb{Q})) = 1$, and from the proof of the first statement of Theorem 2 (parts 1), 2) and 3)) it is clear that all integer points on E_1 are given by $(x, y) \in \{(-1, 0), (0, \pm 1), (120, \pm 6479)\}$. Hence Theorem 2 is true for $k = 1$.

Remark 2. As the coefficients of E_k grow exponentially, the computation of the rank of E_k for large k is difficult. The following values of $\text{rank}(E_k(\mathbb{Q}))$ were computed using the programs SIMATH ([18]) and *mwrank* ([5]):

k	1	2	3	4	5	7	8*	9	10*
$\text{rank}(E_k(\mathbb{Q}))$	1	2	3	3	2	4	4	3	3

In the cases $k = 8, 10$, the rank is computed assuming the Parity Conjecture. For $k = 6, 11, 12$, under the same conjecture, we obtained that $\text{rank}(E_k(\mathbb{Q}))$ is equal to 2 or 4. We also verified by SIMATH that for $k = 3$ and $k = 4$ (when $\text{rank}(E_k(\mathbb{Q})) > 2$) all integer points on E_k are given by the values from Theorem 2.

Remark 3. Let us mention that BREMNER, STROEKER and TZANAKIS [4] proved recently a similar result as the first statement of our Theorem 2 for the family of elliptic curves

$$C_k: \quad y^2 = \frac{1}{3}x^3 + \left(k - \frac{1}{2}\right)x^2 + \left(k^2 - k + \frac{1}{6}\right)x,$$

under the assumptions $\text{rank}(C_k(\mathbb{Q})) = 1$ and $C_k(\mathbb{Q})/C_k(\mathbb{Q})_{\text{tors}} = \langle(1, k)\rangle$.

5. Proof of Theorem 2 ($3 \leq k \leq 40$)

We pointed out in Remark 2 that the coefficients of E_k are growing very fast. Therefore, using SIMATH¹ we were able to compute the integer points of $E_k(\mathbb{Q})$ only for $k \leq 4$. However, the following elementary argument gives us the proof of the second statement of Theorem 2.

Notice the following relations

$$(13) \quad c_0 = 0, \quad c_1 = 8, \quad c_{k+2} = 14c_{k+1} - c_k + 8, \quad \text{if } k \geq 0,$$

$$(14) \quad t_0 = 1, \quad t_1 = 5, \quad t_{k+2} = 4t_{k+1} - t_k, \quad \text{if } k \geq 0,$$

$$(15) \quad s_0 = 1, \quad s_1 = 3, \quad s_{k+2} = 4s_{k+1} - s_k, \quad \text{if } k \geq 0,$$

$$(16) \quad c_k + 1 = s_k^2 \implies c_k = (s_k + 1)(s_k - 1),$$

$$(17) \quad 3c_k + 1 = t_k^2 \implies 3c_k = (t_k + 1)(t_k - 1),$$

$$(18) \quad 3(c_k - 1) = (t_k + 2)(t_k - 2),$$

$$(19) \quad c_k - 3 = (s_k + 2)(s_k - 2).$$

We have $8 \mid c_k$ for any $k \geq 0$ by (13). Hence s_k and t_k are odd. We have further $3 \nmid c_k - 1$ by (16).

Assume that $(x, y) \in \mathbb{Z}^2$ is a solution of (4). Put $D_1 = \gcd(x + 1, 3x + 1)$, $D_2 = \gcd(x + 1, c_k x + 1)$ and $D_3 = \gcd(3x + 1, c_k x + 1)$. As $D_1 = \gcd(x + 1, 3x + 1) = \gcd(x + 1, 2)$, we have $D_1 = 1$ if $x + 1$ is odd, and $D_1 = 2$ if $x + 1$ is even. We have further $D_2 = \gcd(x + 1, c_k x + 1) =$

¹SIMATH is presently the only available computer algebra system which is capable to compute all integer points of elliptic curves. There is implemented the algorithm of GEBEL, PETHŐ and ZIMMER [10].

$\gcd(x+1, c_k-1)$ and $D_3 = \gcd(3x+1, c_kx+1) = \gcd(3x+1, c_k-3)$.
Hence D_1, D_2 and D_3 are pairwise relatively prime.

Assume first $D_1 = 1$. Then there exist $x_1, x_2, x_3 \in \mathbf{Z}$ such that

$$\begin{aligned} x+1 &= D_2x_1^2 \\ 3x+1 &= D_3x_2^2 \\ c_kx+1 &= D_2D_3x_3^2. \end{aligned}$$

Eliminating x we obtain the following system of equations

$$\begin{aligned} 3D_2x_1^2 - D_3x_2^2 &= 2 \\ c_kx_1^2 - D_3x_3^2 &= \frac{c_k-1}{D_2}. \end{aligned}$$

Similarly, if $D_1 = 2$, then (4) implies

$$\begin{aligned} x+1 &= 2D_2x_1^2 \\ 3x+1 &= 2D_3x_2^2 \\ c_kx+1 &= D_2D_3x_3^2, \end{aligned}$$

from which we obtain

$$\begin{aligned} 3D_2x_1^2 - D_3x_2^2 &= 1 \\ 2c_kx_1^2 - D_3x_3^2 &= \frac{c_k-1}{D_2}. \end{aligned}$$

Hence, to find all integer solutions of (4), it is enough to find all integer solutions of the systems of equations

$$(20) \quad d_1x_1^2 - d_2x_2^2 = j_1,$$

$$(21) \quad d_3x_1^2 - d_2x_3^2 = j_2,$$

where

- $d_1 = 3D_2$, D_2 is a square-free divisor of $c_k-1 = (t_k+2)(t_k-2)/3$,
- $d_2 = D_3$, D_3 is a square-free divisor of $c_k-3 = (s_k+2)(s_k-2)$, which is not divisible by 3,

- $(d_3, j_1, j_2) = (c_k, 2, \frac{c_k-1}{D_2})$ or $(d_3, j_1, j_2) = (2c_k, 1, \frac{c_k-1}{D_2})$.

We expect that most of the systems (20)–(21) are not solvable. To exclude as early as possible the unsolvable systems we considered the equations (20) and (21) separately modulo appropriate prime powers.

As $8 \mid c_k$ and $c_k \mid d_3$, and d_2 and j_2 are odd, the equation (21) is solvable modulo 8 only if $-d_2 j_2 \equiv 1 \pmod{8}$.

Assume that equation (20) is solvable. Let p be an odd prime divisor of d_2 . Then (20) implies

$$d_1 x_1^2 \equiv j_1 \pmod{p},$$

hence

$$(d_1 x_1)^2 \equiv j_1 d_1 \pmod{p},$$

i.e. $\left(\frac{j_1 d_1}{p}\right) = 1$, where $\left(\frac{\cdot}{p}\right)$ denotes the Legendre symbol. Similarly, (21) implies $\left(\frac{j_2 d_3}{p}\right) = 1$. If q and r are odd prime divisors of d_1 and d_3 respectively, then we obtain the following conditions for the solvability of (20) and (21): $\left(\frac{-j_1 d_2}{q}\right) = 1$ and $\left(\frac{-j_2 d_2}{r}\right) = 1$.

Let finally p_1 be an odd prime divisor of j_2 , such that $\text{ord}_{p_1}(j_2)$ is odd. Then a necessary condition for solvability of equation (21) is: $\left(\frac{d_2 d_3}{p_1}\right) = 1$.

We performed this test for $3 \leq k \leq 40$ and we found that, apart from the systems listed in the following table, all are unsolvable except those of the form

$$3x_1^2 - x_2^2 = 2,$$

$$c_k x_1^2 - x_3^2 = c_k - 1,$$

and this system is equivalent to the system (2) and (3) which is completely solved by Theorem 1.

We considered in the case $k = 19$ equations (20) and (21), with the values of d_1, d_2, d_3, j_1, j_2 given in the table, modulo 5. We obtained

$$x_1^2 - 4x_2^2 \equiv 2 \pmod{5},$$

$$3x_1^2 - 4x_3^2 \equiv 1 \pmod{5}.$$

k	d_1, d_2, d_3, j_1, j_2
19	251210975091, 44809, 3371344269872647091408, 2, 40261110431
23/1	380631510488414383527682077, 11263976658479, 253754340325609589018454720, 1, 1
23/2	19509779867757, 11263976658479, 25375430325609589018454720, 1, 19509779867761
23/3	58529339603283, 1, 126877170162804794509227360, 2, 6503259955919
35	20288310329233162249058888791445649852717, 2254256703248129138784320976827294428079, 13525540219488774832705925860963766568480, 1, 1
37	187060083, 1489467623820555129, 1311942540724389723505929002667880175005208, 2, 21040446251556347115048521645334887

The first congruence implies $x_1^2 \equiv 1, 2$ or $3 \pmod{5}$, and the second congruence implies $x_1^2 \equiv 0, 2$ or $4 \pmod{5}$. Hence, $x_1^2 \equiv 2 \pmod{5}$, which is a contradiction.

In the cases $k = 23/3$ and $k = 35$ we used arithmetical properties of some real quadratic number fields.

In the case $k = 23/3$ we have $d_3 = 126877170162804794509227360$. The fundamental unit of the order $\mathbb{Z}[\sqrt{d_3}] = \mathbb{Z}[\sqrt{d_2 d_3}]$ is $\varepsilon = 11263976658481 + \sqrt{d_3}$. By a theorem of NAGELL [15, Theorem 108a] the base solution of the equation

$$x_3^2 - 126877170162804794509227360x_1^2 = -6503259955919$$

satisfies $0 < x_1^{(0)} < 1$, which is impossible.

In the case $k = 35$ the fundamental unit of the order $\mathbb{Z}[\sqrt{d_1 d_2}]$ is $u + \sqrt{d_1 d_2}$, where $u = 6762770109744387416352962930481883284238$. A necessary condition for the solvability of the equation $d_1 x_1^2 - d_2 x_2^2 = 1$ is that $2d_1 | (u + 1)$ (see [11]). But $\frac{u+1}{2d_1} = \frac{1}{6}$, and hence the last equation has no solution. $\square$

In the remaining three cases $k = 23/1, 23/2$ and 37 all our methods fail to work.

References

- [1] W. A. ANGLIN, Simultaneous Pell equations, *Math. Comp.* **65** (1996), 355–359.
- [2] A. BAKER and H. DAVENPORT, The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$, *Quart. J. Math. Oxford Ser. (2)* **20** (1969), 129–137.
- [3] A. BAKER and G. WÜSTHOLZ, Logarithmic forms and group varieties, *J. Reine Angew. Math.* **442** (1993), 19–62.
- [4] A. BREMNER, R. J. STROEGER and N. TZANAKIS, On sums of consecutive squares, *J. Number Theory* **62** (1997), 39–70.
- [5] J. E. CREMONA, Algorithms for Modular Elliptic Curves, *Cambridge Univ. Press*, 1997.
- [6] L. E. DICKSON, History of the Theory of Numbers, Vol. 2, *Chelsea, New York*, 1992.
- [7] A. DUJELLA, The problem of the extension of a parametric family of Diophantine triples, *Publ. Math. Debrecen* **51** (1997), 311–322.
- [8] A. DUJELLA, A proof of the Hoggatt–Bergum conjecture, *Proc. Amer. Math. Soc.* **127** (1999), 1999–2005.
- [9] A. DUJELLA and A. PETHŐ, A generalization of a theorem of Baker and Davenport, *Quart. J. Math. Oxford Ser. (2)* **49** (1998), 291–306.
- [10] J. GEBEL, A. PETHŐ and H. G. ZIMMER, Computing integral points on elliptic curve, *Acta Arith.* **68** (1994), 171–192.
- [11] A. GRELAK, A. GRZYTCZUK, On the diophantine equation $ax^2 - by^2 = c$, *Publ. Math. Debrecen* **44** (1994), 291–299.
- [12] D. HUSEMÖLLER, Elliptic Curves, *Springer-Verlag, New York*, 1987.
- [13] A. KNAPP, Elliptic Curves, *Princeton Univ. Press*, 1992.
- [14] M. MIGNOTTE and A. PETHŐ, Sur les carrés dans certaines suites de Lucas, *Théor. Nombres Bordeaux* **5** (1993), 333–341.
- [15] T. NAGELL, Introduction to Number Theory, *Almqvist, Stockholm; Wiley, New York*, 1951.
- [16] K. NAKAMULA and A. PETHŐ, Squares in binary recurrence sequences, Number Theory, Diophantine, Computational and Algebraic Aspects (K. Győry, A. Pethő and V. T. Sós, eds.), *Walter de Gruyter, Berlin*, 1998, 409–421.
- [17] K. ONO, Euler’s concordant forms, *Acta Arith.* **78** (1996), 101–123.
- [18] SIMATH manual, *Saarbrücken*, 1997.

ANDREJ DUJELLA
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF ZAGREB
 BIJENIČKA CESTA 30
 10000 ZAGREB
 CROATIA
E-mail: duje@math.hr

ATTILA PETHŐ
 INSTITUTE OF MATHEMATICS
 COMPUTER SCIENCE
 LAJOS KOSSUTH UNIVERSITY
 H-4010 DEBRECEN P.O. BOX 12
 HUNGARY
E-mail: pethoe@math.klte.hu

(Received November 18, 1998; revised April 19, 1999)