

## Graded Lie algebra associated to a SODE

By ZOLTÁN MUZSNAY (Debrecen)

**Abstract.** In this paper we introduce a graded Lie algebra associated to a second order differential equation which gives a powerful tool to the study of the inverse problem of the calculus of variations. We give effective generalizations of Douglas' criteria for the existence of a regular Lagrangian associated to a SODE.

### 1. Introduction

The inverse problem of the calculus of variations is an old problem of Differential Geometry consisting of the characterization of second order ordinary differential equations (SODE) derivable from a variational principle. In this problem one wants to determine whether a given SODE expresses that the unknown is the critical point of a functional.

One of the most important contribution to this problem is a paper of J. DOUGLAS [4], where he classifies systems of variational differential equations of second order in the two-dimensional case. He showed that the Euler–Lagrange partial differential system

$$(1.1) \quad \frac{d}{dt} \frac{\partial E}{\partial y^i} - \frac{\partial E}{\partial x^i} = 0,$$

---

*Mathematics Subject Classification:* 49N45, 58E30, 70H35.

*Key words and phrases:* inverse problem, calculus of variations, Euler–Lagrange equation, variational multiplier, spray, connection, derivations.

This research was partially supported by OTKA grant 32058.

(where the unknown function is  $E$ ) associated to a SODE  $\ddot{x}^i = f^i(x, \dot{x})$ ,  $i = 1, \dots, n$ , is equivalent to the first order partial differential system

$$\begin{aligned}
 (1.2) \quad & \frac{d}{dt}g_{ij} + \frac{1}{2}\frac{\partial f^k}{\partial y^j}g_{ik} + \frac{1}{2}\frac{\partial f^k}{\partial y^i}g_{jk} = 0, \\
 & A_j^k g_{ik} - A_i^k g_{jk} = 0, \\
 & \frac{\partial g_{ij}}{\partial y^k} - \frac{\partial g_{ik}}{\partial y^j} = 0, \\
 & g_{ij} - g_{ji} = 0, \\
 & \det(g_{ij}) \neq 0,
 \end{aligned}$$

where the unknown functions are  $g_{ij}$ ,  $i, j = 1, \dots, n$ , and  $A_j^i$  are the components of the Douglas tensor (called also Jacobi endomorphism). A solution  $E$  of (1.1) gives a solution of (1.2) by taking

$$(1.3) \quad g_{ij} = \frac{\partial^2 E}{\partial y^i \partial y^j}$$

and conversely, for every solution of (1.2) there exists a regular a solution  $E$  of (1.1) so that (1.3) holds. A solution of (1.2) is called *variational multiplier*.

Obstructions to the existence of a variational principle, i.e. integrability conditions of the Euler–Lagrange PDE, are in general very complex. However, in his article [4] DOUGLAS gives a simple criteria on the existence of the variational multiplier and therefore on the existence of a variational principle for SODEs on 2-dimensional manifolds. This criteria can be easily carried over to the  $n$ -dimensional case (see [2], [8], [11]). In [12] the authors found a double hierarchy of algebraic conditions for the variational multiplier which is determined by the Douglas tensor, the curvature tensor and their derivatives.

In this paper we introduce a graded Lie-algebra associated in a natural way with the SODE using a differential algebraic characterization of connections and derivations. It contains algebraic conditions on the variational multiplier and in generic cases it gives a significant part of the obstruction to the existence of a variational principle (Theorem 2). This concept is of particular interest when the dimension of the base manifold is large, because we are able to obtain new information about the structure of the obstructions (Theorems 4 and 5).

## 2. Frölicher–Nijenhuis theory

In this paper we use extensively the Frölicher–Nijenhuis' theory of the derivation associated to vector valued forms. A complete description can be found in [5] or [10]. We recall here only the basic elements of this theory.

We denote by  $\Lambda(M)$  (resp.  $\Psi(M)$ ) the  $C^\infty(M)$  modulus of the scalar (resp. vector valued) forms. The Frölicher–Nijenhuis theory gives a complete description of the derivation of  $\Lambda(M)$  with the help of  $\Psi(M)$ .

*Definition 1.* A morphism  $D : \Lambda(M) \longrightarrow \Lambda(M)$  is a derivation of  $\Lambda(M)$  of degree  $r$  if it satisfies the following conditions:

- a)  $D(\Lambda^p(M)) \subset \Lambda^{p+r}(M)$ ,
- b)  $D(a\omega + b\omega') = aD\omega + bD\omega'$ ,  $a, b \in \mathbb{R}$
- c)  $D(\omega \wedge \pi) = D\omega \wedge \pi + (-1)^{r \deg \omega} \omega \wedge D\pi$ .

The bracket of two derivations  $D_1$  and  $D_2$  is defined by

$$[D_1, D_2] = D_1 D_2 - (-1)^{(\deg D_1)(\deg D_2)} D_2 D_1.$$

*Definition 2.* A derivation is called of  $i_*$  type or algebraic, if its action is trivial on  $\Lambda^o(M)$ , and of  $d_*$  type, if it commutes with the operator  $d$ .

Every derivation is determined by his action on  $\Lambda^o(M)$  and  $\Lambda^1(M)$ . So the  $i_*$  type derivations are those which are completely determined by their action on  $\Lambda^1(M)$ , and the  $d_*$  type derivations are determined by their action on  $C^\infty(M)$ .

An  $i_*$  and a  $d_*$  type derivation can be associated to a vector valued  $l$ -form  $L \in \Psi^l(M)$ , denoted by  $i_L$  and  $d_L$ , in the following way:

1. if  $\deg L = 0$ , (i.e.  $L \in \mathfrak{X}(M)$  is a vector field on  $M$ ):

$$i_L \omega := \omega(L), \text{ and}$$

$$d_L \omega := \mathcal{L}_L \omega;$$

2. if  $\deg L = l > 1$ :

$$i_L \omega(X_1, \dots, X_l) := \omega(L(X_1, \dots, X_l)), \quad \text{where } \omega \in \Lambda^1(M);$$

$$d_L f(X_1, \dots, X_l) := df(L(X_1, \dots, X_l)), \quad \text{where } f \in \Lambda^o(M),$$

where  $\omega \in \Lambda^1(TM)$ , and  $\mathcal{L}_L$  denotes the Lie derivation with respect to  $L$ . Conversely, it is easy to show that every  $i_*$  or  $d_*$  type derivation can be written in the above form with the help of some vector valued form. Therefore we arrive at the following:

**Proposition 3.** *Let  $L$  and  $M$  be vector valued differential  $l$ - and  $m$ -forms. Then there exists a unique vector valued  $(l + m)$ -form (denoted by  $[L, M]$ ) which satisfies the equation*

$$[d_L, d_M] = d_{[L, M]}.$$

By this bracket,  $\Psi(M)$  is a graded Lie algebra.

### 3. Inverse problem of the calculus of variations

We turn our attention to the inverse problem of the variational calculus. A coordinate free formulation of this problem can be given by the notion of sprays introduced by KLEIN in [9].

Let  $J$  be the canonical vertical endomorphism and  $C$  the canonical vertical vector field. In the local coordinate system  $(x^i)$  on  $M$  and  $(x^i, y^i)$  on  $TM$ ,

$$J = dx^\alpha \otimes \frac{\partial}{\partial y^\alpha}, \quad C = y^\alpha \frac{\partial}{\partial y^\alpha}.$$

A vector field  $S$  on  $TM$  is a spray if  $JS = C$ . A curve  $\gamma : I \rightarrow M$  is associated to the spray  $S$  if  $\gamma'$  is an integral curve of  $S$  i.e.  $S_{\gamma'} = S\gamma''$ . In a coordinate system the expression of a spray is

$$(3.1) \quad S = y^\alpha \frac{\partial}{\partial x^\alpha} + f^\alpha(x, y) \frac{\partial}{\partial y^\alpha},$$

and the path  $(x^\alpha(t))$  is associated to the spray (3.1) if and only if the second order differential equation

$$(3.2) \quad \frac{d^2 x^\alpha}{dt^2} = f^\alpha \left( x, \frac{dx}{dt} \right)$$

holds. Therefore a spray is a coordinate free presentation of a SODE on the manifold  $M$ .

*Definition 4.* The Lagrangian function  $E : TM \rightarrow \mathbb{R}$  is called regular, if the 2-form  $\Omega_E := dd_J E$  is symplectic.

Using a coordinate system, a Lagrangian  $E$  is regular if and only if the matrix  $\left( \frac{\partial^2 E}{\partial y^\alpha \partial y^\beta} \right)$  is not singular. It is well known that to any Lagrangian which defines a regular variational problem, a spray (i.e. a SODE) can be associated, in the following way:

**Theorem 1** [6]. *Let  $E \in C^\infty(TM)$  be a regular Lagrangian. The vector field  $S$  defined by the equation*

$$(3.3) \quad i_S \Omega_E = d(E - \mathcal{L}_C E)$$

*is a spray, and the paths associated to the spray  $S$  are solutions of the corresponding variational problem.*

So the spray  $S$  is variational if there exists a regular function  $E \in C^\infty(TM)$ , so that the equation (3.3) holds.

*Remark.* Let  $E$  be an arbitrary Lagrangian on  $M$  and  $S$  be a spray. The associated semi-basic 1-form

$$(3.4) \quad \omega_E := i_S \Omega_E + d\mathcal{L}_C E - dE$$

is called *Euler–Lagrange* form. The solution of the inverse problem for a given SODE is a regular Lagrangian such that the equation  $\omega_E = 0$  holds.

#### 4. Identities satisfied by variational sprays

From now on we shall work on  $TM$ , the tangent manifold of  $M$ . Where there is no possibility of confusion,  $TTM$ ,  $T^*TM$  and  $T^vTM$  will be noted as  $T$ ,  $T^*$  and  $T^v$  respectively.

The space of semi-basic scalar (resp. vectorial)  $l$ -forms is denoted by  $\Lambda_v^l T_v^*$  (resp.  $\Psi^l T_v^*$ ). We recall that a  $p$ -form  $\omega \in \Lambda^p T^*$  is semi-basic if  $\omega(X_1, \dots, X_p) = 0$  when one of the vectors  $X_i$  is vertical, and a vector valued  $l$ -form  $L$  is semi-basic if  $L(X_1, \dots, X_p) = 0$  when one of the vectors  $X_i$  is vertical and its value is vertical.

A connection  $\Gamma$  can be associated to every spray  $S$  defined by the formula  $\Gamma := [J, S]$  (see [7]): it is easy to check that  $\Gamma^2 = I$  and the eigenspace corresponding to the eigenvalue  $-1$  is the vertical space  $T^v$ . If we denote the eigenspace corresponding to the eigenvalue  $+1$  by  $T^h$ , then  $TTM$  can be decomposed as

$$TTM = T^h \oplus T^v.$$

Let  $h$  and  $v$  be the corresponding projectors:  $h := \frac{1}{2}(I + \Gamma)$ ,  $v := \frac{1}{2}(I - \Gamma)$ . The curvature of the connection  $\Gamma$  is the vector valued 2-form

$$R := -\frac{1}{2}[h, h].$$

The almost complex structure  $F$  associated to  $\Gamma$  which exchanges the horizontal and the vertical space is defined by

$$F = h[S, h] - J.$$

**Property 5.** *Let  $E$  be a Lagrangian,  $\Gamma$  a connection on  $M$ ,  $h$  its associated horizontal projection, and  $F$  the associated almost-complex structure. The following properties are equivalent:*

- a)  $i_\Gamma \Omega_E = 0$
- b)  $i_F \Omega_E = 0$
- c)  $\Omega_E(hX, hY) = 0 \ \forall X, Y \in TTM$  (i.e. the horizontal distribution is Lagrangian).

A connection is called *Lagrangian* with respect to  $E$ , if it satisfies the above conditions.

Indeed, we have:

$$i_\Gamma \Omega_E(hX, hY) = 2\Omega_E(hX, hY)$$

$$i_\Gamma \Omega_E(hX, JY) = \Omega_E(hX, JY) - \Omega_E(hX, JY) = 0$$

$$i_\Gamma \Omega_E(JX, JY) = -2\Omega_E(JX, JY) = 0,$$

so a)  $\iff$  c). On the other hand:

$$i_F \Omega_E(hX, hY) = -\Omega_E(JX, hY) - \Omega_E(hX, JY) = -i_J \Omega_E(hX, hY) = 0$$

$$i_F \Omega_E(hX, JY) = -\Omega_E(JX, JY) + \Omega_E(hX, hY) = \Omega_E(hX, hY)$$

$$i_F \Omega_E(JX, JY) = \Omega_E(hX, JY) + \Omega_E(JX, hY) = 0,$$

so b)  $\iff$  c).

**Property 6.** *Let  $E$  be a Lagrangian on the manifold  $M$ . Then*

$$(4.1) \quad d_J \omega_E = i_\Gamma \Omega_E.$$

Consequently, if the spray  $S$  is variational and  $E$  is a Lagrangian associated to  $S$ , then the horizontal distribution associated to the spray  $S$  must be Lagrangian with respect to the symplectic 2-form  $\Omega_E$ .

PROOF. The Euler–Lagrange form can be written in the following form:

$$\begin{aligned}\omega_E &= i_S dd_J E + d\mathcal{L}_S E - dE = \mathcal{L}_S d_J E - dE \\ &= d_J \mathcal{L}_S E - i_{[J, S]} dE = d_J \mathcal{L}_S E - 2d_h E.\end{aligned}$$

Since the vertical distribution is integrable, we get  $[J, J] = 0$ , we have  $d_J^2 = d_J \circ d_J = d_{[J, J]} = 0$ . So

$$\begin{aligned}d_J \omega_E &= -2d_J d_h E = 2d_h d_J E = 2(i_h dd_J E - di_h d_J E) \\ &= 2i_h \Omega_E - 2\Omega_E = i_\Gamma \Omega_E.\end{aligned}$$

If the spray is variational and  $E$  is a Lagrangian associated with  $S$ , we have  $\omega_E = 0$ , then  $i_\Gamma \Omega_E = 0$ , so the connection associated to the spray is Lagrangian.  $\square$

*Definition 7.* Let  $S$  be a spray on  $M$ ,  $L \in \Psi_v(TM)$ . The semi-basic derivation of  $L$  with respect to the spray  $S$  is

$$(4.2) \quad L' := h^* v[S, L]$$

where  $h^* L(X_1, \dots, X_l) := L(hX_1, \dots, hX_l)$ .

We have the following

**Proposition 8.** Let  $S$  be a spray on  $M$  and  $L \in \Psi_v(TM)$ . We have the formula

$$(4.3) \quad L' = [S, L] + FL - L\bar{\wedge}F.$$

In particular, suppose that  $S$  is variational,  $E$  being a Lagrangian associated to  $S$ . If the equation  $i_L \Omega_E = 0$  holds, then the equations

$$(4.4) \quad i_{L'} \Omega_E = 0, \quad i_{L''} \Omega_E = 0, \quad i_{L'''} \Omega_E = 0, \quad \text{etc.}$$

hold too.

PROOF. To show the first expression, we note that

$$\begin{aligned}
L'(X_1, \dots, X_l) &= v[S, L](hX_1, \dots, hX_l) \\
&= v[S, L(X_1, \dots, X_l)] - \sum_{i=1}^l L(X_1, \dots, [S, hX_i], \dots, X_l) \\
&= [S, L(X_1, \dots, X_l)] - h[S, L(X_1, \dots, X_l)] \\
&\quad - \sum_{i=1}^l L(X_1, \dots, [S, h]X_i, \dots, X_l) - \sum_{i=1}^l L(X_1, \dots, [S, X_i], \dots, X_l) \\
&= [S, L](X_1, \dots, X_l) + FL(X_1, \dots, X_l) \\
&\quad - \sum_{i=1}^l L(X_1, \dots, h[S, h]X_i, \dots, X_l).
\end{aligned}$$

Using the identity  $h[S, h] = F + J$  and the hypothesis that  $L$  is semi-basic, we obtain (4.3). Secondly, by the formula (4.3) we have

$$\begin{aligned}
i_{L'}\Omega_E &= i_{[S, L]}\Omega_E + i_{FL}\Omega_E - i_F\bar{\wedge}\Omega_E = i_{[S, L]}\Omega_E + i_F i_L \Omega_E - i_L i_F \Omega_E \\
&= \mathcal{L}_S i_L \Omega_E - d_L \omega_E + i_F i_L \Omega_E - i_L i_F \Omega_E.
\end{aligned}$$

When  $S$  is variational and the function  $E$  is a Lagrangian associated to  $S$ , then  $\omega_E = 0$  and the connection  $\Gamma$  is Lagrangian, so we have  $i_F \Omega_E = 0$  (Properties 5 and 6). If the equation  $i_L \Omega_E = 0$  holds, we have also  $i_{L'} \Omega_E = 0$  and recursively we obtain (4.4).  $\square$

*Definition 9.* Let  $h$  be the horizontal projection associated to the connection  $\Gamma = [J, S]$ , and  $L \in \Psi_v^l(TM)$  a vector valued semi-basic  $l$ -form. The operator

$$(4.5) \quad d^h L := [h, L]$$

is the *semi-basic derivation of  $L$  with respect to  $h$* .

**Proposition 10.** *Let  $L$  be a semi-basic vector valued  $l$ -form. Then  $d^h L$  is semi-basic. Moreover assume that  $S$  is variational, and  $E$  is a Lagrangian associated to  $S$ . If the equation  $i_L \Omega_E = 0$  holds, then the equation  $i_{d^h L} \Omega_E = 0$  holds too.*

PROOF. It is not difficult to check that if  $L$  is a semi-basic vector valued  $l$ -form, then  $d^h L$  is also semi-basic. Let us show the second part

of the proposition. Let us assume that  $S$  is variational,  $E$  is a Lagrangian associated to  $S$ , and  $L$  is a vector-valued semi-basic  $l$ -form. By the relation

$$(-1)^l i_{[h,L]} = i_h d_L - d_L i_h - d_L \bar{\wedge} h$$

and taking into account that  $L \bar{\wedge} h = l L$ , because  $L$  is semi-basic, we have

$$(-1)^l i_{d^h L} \Omega_E = (-1)^l i_{[h,L]} dd_J E = i_h d_L dd_J E - d_L i_h dd_J E - l d_L dd_J E.$$

If the equation  $i_L \Omega_E = 0$  holds, then

$$\begin{aligned} (-1)^l i_{d^h L} \Omega_E &= i_h di_L dd_J E - d_L i_{\frac{1}{2}(I+\Gamma)} dd_J E - l di_L dd_J E \\ &= -l di_L dd_J E - \frac{1}{2} d_L i_\Gamma dd_J E = 0. \end{aligned} \quad \square$$

## 5. Graded Lie algebra associated to a SODE

*Definition 11.* The graded Lie algebra  $\mathcal{A}_S$  associated to the spray  $S$  is the graded Lie sub-algebra of the vector-valued forms spanned by the vertical endomorphism  $J$ , the Douglas tensor  $A := v[h, S]$ , and generated by the action of the semi-basic derivation defined in (4.2), the derivation  $d^h$ , and the Frölicher–Nijenhuis bracket  $[\ , \ ]$ . The graduation of  $\mathcal{A}_S$  is given by

$$(5.1) \quad \mathcal{A}_S = \oplus_{k=1}^n \mathcal{A}_S^k$$

where  $\mathcal{A}_S^k := \mathcal{A}_S \cap \Psi^k(TM)$ .

*Remark.* Note that  $J$  and  $A$  are semi-basic and that, as we showed in the preceding paragraph, the space of semi-basic forms is stable by semi-basic derivation defined in (4.2), by the derivation  $d^h$ , and by the Frölicher–Nijenhuis bracket  $[\ , \ ]$ . It follows that  $\mathcal{A}_S$  is a graded Lie sub-algebra of the vector-valued *semi-basic* forms.

The importance of the graded Lie algebra associated to a spray is given by the following:

**Theorem 2.** *Let  $S$  be a variational spray and  $E$  a Lagrangian associated to  $S$ . Then for every element  $L$  of  $\mathcal{A}_S$  the equation*

$$(5.2) \quad i_L \Omega_E = 0$$

*holds. Therefore elements of  $\mathcal{A}_S$  give algebraic conditions on the variational multiplier.*

PROOF. To prove Theorem 2 we will first show that  $J$  and  $A$  satisfy the equation (5.2). Then we will prove that all the vector-valued forms obtained from  $J$  and  $A$  by a finite number of successive operations which define  $\mathcal{A}_S$ , also satisfy the equation (5.2).

1. From  $[J, J] = 0$  we can easily obtain :

$$(5.3) \quad i_J \Omega_E = i_J dd_J E = d_J^2 E = d_{[J, J]} E = 0,$$

so the equation (5.2) holds for  $J$ .

2. For the Douglas tensor we find

$$\begin{aligned} i_A \Omega_E &= i_{[h, S]} \Omega_E + i_F \Omega_E = i_h \mathcal{L}_S \Omega_E - \mathcal{L}_S i_h \Omega_E + i_F \Omega_E \\ &= i_h d\omega_E - \mathcal{L}_S \left( \Omega_E + \frac{1}{2} d_J \omega_E \right) + i_F \Omega_E \\ &= i_h d\omega_E - d\omega_E - \frac{1}{2} \mathcal{L}_S d_J \omega_E + i_F \Omega_E \\ &= d_h \omega_E - \frac{1}{2} \mathcal{L}_S d_J \omega_E + i_F \Omega_E \end{aligned}$$

If  $S$  is variational and  $E$  is a Lagrangian associated to  $S$ , then  $\omega_E = 0$  and the connection  $\Gamma$  is Lagrangian. Therefore every term vanishes, and the equation (5.2) also holds for  $A = L$ .

3. From Propositions 8 and 10, respectively we know that if  $i_L \Omega_E = 0$  holds for  $L \in \mathcal{A}_S$  then  $i_{L'} \Omega_E = 0$ ,  $i_{d^h L} \Omega_E = 0$ , hold too.
4. Let  $K \in \mathcal{A}_S^k$ ,  $L \in \mathcal{A}_S^l$  be semi-basic vector-valued forms, such that  $i_K \Omega_E = 0$  and  $i_L \Omega_E = 0$ . Since  $K$  and  $L$  are semi-basic, we have  $L \bar{\wedge} K \equiv 0$  and hence

$$\begin{aligned} (-1)^l i_{[K, L]} \Omega_E &= (i_K d_L - (-1)^{l(m-1)} d_L i_K - d_{L \bar{\wedge} K}) \Omega_E \\ &= i_K (i_L d - d i_L) dd_J E - (-1)^{l(m-1)} d_L i_K dd_J E - d_{L \bar{\wedge} K} dd_J E \\ &= i_K d i_L \Omega_E - (-1)^{l(m-1)} d_L i_K \Omega_E = 0. \end{aligned}$$

On the other side, it is easy to see that (5.2) gives algebraic condition on the variational multiplier. Indeed, if the spray is variational and  $E$  is a regular Lagrangian associated with  $S$ , then locally one has

$$(5.4) \quad \Omega_E = \frac{1}{2} \left( \frac{\partial^2 E}{\partial x^\alpha \partial y^\beta} - \frac{\partial^2 E}{\partial y^\alpha \partial x^\beta} \right) dx^\alpha \wedge dx^\beta - \frac{\partial^2 E}{\partial y^\alpha \partial y^\beta} dx^\alpha \wedge dy^\beta.$$

If  $L \in \Psi^l(TM)$  is semi-basic, then

$$i_L \Omega_E = \frac{1}{l!} \sum_{\alpha \in \mathfrak{S}_{l+1}} \varepsilon(\alpha) L_{\alpha_1 \dots \alpha_l}^\beta \frac{\partial^2 E}{\partial y^\beta \partial y^{\alpha_{l+1}}} dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_{l+1}},$$

where  $\mathfrak{S}_{p+l-1}$  denotes the  $(p+l-1)!$ -order symmetric group and  $\varepsilon(\alpha)$  the sign of  $\alpha$ . Then the equation  $i_L \Omega_E = 0$  is an algebraic equation

$$(5.5) \quad \sum_{\alpha \in \mathfrak{S}_{l+1}} \varepsilon(\alpha) L_{\alpha_1 \dots \alpha_l}^\beta g_{\beta \alpha_{l+1}} = 0$$

in the variational multiplier  $g_{\alpha\beta} = \frac{\partial^2 E}{\partial y^\alpha \partial y^\beta}$ .  $\square$

*Remark.* From the construction of  $\mathcal{A}_S$  it is clear that  $\mathcal{A}_S$  contains the Douglas tensor and its semi-basic derivatives with respect to  $S$ . On the other hand, the curvature tensor  $R$  is related to  $A$  by the equation  $R = \frac{1}{3}[J, A]$ , so  $\mathcal{A}_S$  also contains  $R$  and its semi-basic derivatives. Therefore  $\mathcal{A}_S$  contains the double hierarchy of algebraic conditions for the variational multiplier founded in [12].

**Theorem 3.** *If at  $x \in TM$  one has*

$$\text{rank}\{J, A, A', \dots, A^{(k)}, \dots\}_{k \in \mathbb{R}} > \frac{n(n+1)}{2},$$

*then  $S$  is not variational in the neighborhood of  $x$ .*

PROOF. Let us suppose that  $S$  is variational,  $E$  is an associated regular Lagrangian, and  $g_{ij} = \frac{\partial^2 E}{\partial y^i \partial y^j}$  is a variational multiplier. For every  $L \in \mathcal{A}_S^1$  the condition  $i_L \Omega_E = 0$  gives

$$g_{ik} L_j^k = g_{jk} L_i^k.$$

i.e.  $L$  is symmetric with respect to  $g$ . Since the tensors  $J, A, A', A'', \dots, A^{(\frac{n(n+1)}{2}-1)}$  are elements of  $\mathcal{A}_S$ , we have  $i_{A^{(k)}} \Omega_E = 0$ . Therefore, if

the spray is variational, then the tensors  $J, A, A', A'', \dots, A^{(\frac{n(n+1)}{2}-1)}$  are self-adjoint with respect to  $g$ . But the space of the  $(1-1)$  tensors which are self-adjoint with respect to a regular matrix is  $\frac{n(n+1)}{2}$ -dimensional. Consequently if the spray is variational, then  $J, A, A', A'', \dots, A^{(\frac{n(n+1)}{2}-1)}$  are linearly dependent.  $\square$

If  $\dim M = 2$ , then  $\mathcal{A}_S$  only contains  $J, A$  and the hierarchy given by its semi-basic derivatives  $A', A''$ , etc. However, if  $\dim M > 2$ , then we find other hierarchies in  $\mathcal{A}_S$  which give, in the generic case, new necessary conditions for the variational multipliers. We arrive at the following generalization of the Theorem 3:

**Theorem 4.** *Let  $S$  be a spray and  $x \in TM$ . If there exists an integer  $k < n$  for which*

$$(5.6) \quad \dim \mathcal{A}_S^k(x) > k \binom{n+1}{k+1},$$

*then the spray is not variational in a neighborhood of  $x$ .*

PROOF. Let  $S$  be a spray and  $E$  a regular Lagrangian. We consider for every  $k = 1, \dots, n$  the morphism

$$\begin{aligned} \Lambda^k T_v^* \otimes T^v &\xrightarrow{\psi_k} \Lambda^{k+1} T_v^* \\ L &\longrightarrow i_L \Omega_E. \end{aligned}$$

By the regularity of  $E$  the 2-form  $\Omega_E$  is symplectic, and the morphism  $\psi_k$  is onto. Indeed, it is easy to see that if  $\{X_1, \dots, X_n\}$  is a basis of  $T^v$ , then  $\alpha_1, \dots, \alpha_n \in T^*$  defined by  $\alpha_i = i_{X_i} \Omega_E$ , gives a basis of  $T_v^*$ . Consequently

$$(5.7) \quad \{\alpha_{i_1} \wedge \dots \wedge \alpha_{i_k} \wedge \alpha_{i_{k+1}}\}_{1 \leq i_1 < \dots < i_{k+1} \leq n}$$

is a basis of  $\Lambda^{k+1} T_v^*$  and

$$(5.8) \quad \{\alpha_{i_1} \wedge \dots \wedge \alpha_{i_k} \otimes X_{i_{k+1}}\}_{1 \leq i_1 < \dots < i_k \leq n, 1 \leq i_{k+1} \leq n}$$

gives a basis of  $\Lambda^k T_v^* \otimes T^v$ . Moreover, if the components of  $\Lambda \in \Lambda^{k+1} T_v^*$  with respect to the basis (5.7) are  $\Lambda^{i_1 \dots i_{k+1}}$ , then  $\Lambda = \psi_k(L)$  where  $L = \Lambda_{i_1 \dots i_{k+1}} \alpha_{i_1} \wedge \dots \wedge \alpha_{i_k} \otimes X_{i_{k+1}}$ . This proves that  $\psi_k$  is onto. Therefore

$$\text{rank } \psi_k = \dim \Lambda^{k+1} T_v^* = \binom{n}{k+1} = \frac{n!}{(k+1)!(n+1-k)!},$$

and

$$(5.9) \quad \dim \operatorname{Ker} \psi_k = n \cdot \binom{n}{k} - \binom{n}{k+1} \\ = \frac{k(n+1)n!}{(k+1)!(n-k)!} = k \cdot \binom{n+1}{k+1}$$

On the other hand using Theorem 2 we have

$$\mathcal{A}_S^k \subset \operatorname{Ker} \psi_k.$$

But if the inequality (5.6) holds, then  $\dim \mathcal{A}_S^k > \dim \operatorname{Ker} \psi_k$ , and consequently the spray is not variational.  $\square$

*Definition 12.* Let  $S$  be a spray  $x \in TM$ , and let us consider the system of linear equations

$$(5.10) \quad \left\{ \sum_{i \in \mathfrak{S}_{l+1}} \varepsilon(i) L_{i_1 \dots i_l}^j x_{j i_{l+1}} = 0 \mid L \in \mathcal{A}_S(x) \right\}$$

in the symmetric variables  $x_{ij}$  ( $x_{ij} = x_{ji}$ ) where  $L_{i_1 \dots i_l}^j$  are the components of  $L \in \mathcal{A}_S(x)$ . The rank of the linear equations (5.10) is called the *rank of the spray* at  $x$ .

*Remark.* As equation (5.5) shows, the rank of a spray gives the number of independent equations satisfied by the variational multipliers. Consequently, if the system (5.10) does not have a solution with  $\det(x_{ij}) \neq 0$ , then there is no variational multiplier for  $S$ , and therefore the spray is non-variational. Thus we arrive at

**Theorem 5.** *If at  $x \in TM$  we have*

$$\operatorname{rank} S(x) \geq \frac{n(n+1)}{2}$$

*then  $S$  is non-variational in a neighborhood of  $x$ .*

*Acknowledgement.* The author gratefully acknowledges the suggestions and encouragement of Professor JOSEPH GRIFONE.

## References

- [1] I. M. ANDERSON, Aspects of the inverse problem to the calculus of variations, *Arch. Math. Brno* **24** no. 4 (1988), 181–202.
- [2] I. M. ANDERSON and G. THOMPSON, The Inverse Problem of the Calculus of Variations for Ordinary Differential Equations, *Mem. AMS* **98** (1992), 473.
- [3] M. CRAMPIN, W. SARLET, E. MARTINEZ, G. B. BYRNES and G. E. PRINCE, Towards a geometrical understanding of Douglas's solution of the inverse problem of the calculus of variations, *Inv. Probl.* **10** (1994), 245–260.
- [4] J. DOUGLAS, Solution to the inverse problem of the calculus of variations, *Trans. Amer. Math. Soc.* **50** (1941), 71–128.
- [5] A. FRÖLICHER and A. NIJENHUIS, Theory of vector-valued differential forms, *Proc. Kon. Ned. Akad. A* **59** (1956), 338–359.
- [6] C. GODBILLON, Géométrie différentielle et mécanique analytique, *Hermann, Paris*, 1969.
- [7] J. GRIFONE, Structure presque-tangente et connexions I, II, *Ann. Inst. Fourier* **XXII** (1) (1972), 287–334; **XXII** (3), 291–338.
- [8] J. GRIFONE and Z. MUZSNAY, On the inverse problem of the variational calculus: existence of Lagrangians associated with a spray in the isotropic case, *Ann. Inst. Fourier* **49** (4) (1999), 1387–1421.
- [9] J. KLEIN, On variational second order differential equations: polynomial case, *Diff. Geom. Appl. Proc. Conf. Aug. 24–28, 1993, Silesian Univ. Opava, 1993*, 449–459.
- [10] I. KOLÁŘ, P. W. MICHOR and JAN SLOVÁK, Natural Operations in Differential Geometry, *Springer, Berlin*, 1993.
- [11] W. SARLET, The Helmholtz conditions revisited. A new approach to the inverse problem of Lagrangian dynamics, *J. Phys. A* **15** (1982), 1503–1517.
- [12] W. SARLET, M. CRAMPIN and E. MARTINEZ, The integrability conditions in the inverse problem of the calculus of variations for second-order ordinary differential equations, *Acta Appl. Math.* **54** (1998), 233–273.

ZOLTÁN MUZSNAY  
 INSTITUTE OF MATHEMATICS AND INFORMATICS  
 UNIVERSITY OF DEBRECEN  
 H-4010 DEBRECEN, P.O. BOX 12  
 HUNGARY

E-mail: muzsnay@math.klte.hu

(Received December 30, 1999; revised August 31, 2000)