

Single-valued and multi-valued Caristi type operators

By ADRIAN PETRUȘEL (Cluj-Napoca)
and ALINA SÎNTĂMĂRIAN (Cluj-Napoca)

Abstract. New results in connection with some single-valued and multi-valued Caristi type operators are established.

1. Introduction

Caristi's fixed point theorem states that each operator f from a complete metric space (X, d) into itself satisfying the condition: there exists a proper lower semi-continuous function $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ such that:

$$(1.1) \quad d(x, f(x)) + \varphi(f(x)) \leq \varphi(x), \quad \text{for each } x \in X$$

has at least a fixed point $x^* \in X$, i.e. $x^* = f(x^*)$ (see [5]). There are several extensions and generalizations of this important principle of the nonlinear analysis (see for example [4], [6], etc.). One of the latest, asserts that if (X, d) is a complete metric space, $x_0 \in X$, $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ is a proper lower semi-continuous function and $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function such that $\int_0^\infty \frac{ds}{1+h(s)} = +\infty$, then each single-valued operator f from X to itself satisfying the condition:

$$(1.2) \quad \text{for each } x \in X, \quad \frac{d(x, f(x))}{1 + h(d(x_0, x))} + \varphi(f(x)) \leq \varphi(x),$$

has at least a fixed point (see [15]). For the multi-valued case, if F is an operator of the complete metric space X into the space of all nonempty

Mathematics Subject Classification: 47H10, 54H25.

Key words and phrases: fixed point, Reich type operator, entropy.

subsets of X and there exists a proper lower semi-continuous function $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ such that

$$(1.3) \quad \text{for each } x \in X, \text{ there is } y \in F(x) \text{ so that } d(x, y) + \varphi(y) \leq \varphi(x),$$

then the multi-valued map F has at least a fixed point $x^* \in X$, i.e. $x^* \in F(x^*)$ (see for example [1]). Moreover, if F satisfies the stronger condition:

$$(1.4) \quad \text{for each } x \in X \text{ and each } y \in F(x) \text{ we have } d(x, y) + \varphi(y) \leq \varphi(x),$$

then the multi-valued map F has at least a strict fixed point $x^* \in X$, i.e. $\{x^*\} = F(x^*)$ (see [1]). On the other hand, if F is a multi-valued operator with nonempty closed values and $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ is a proper lower semi-continuous function such that the following condition holds:

$$(1.5) \quad \text{for each } x \in X, \quad \inf\{d(x, y) + \varphi(y) : y \in F(x)\} \leq \varphi(x),$$

then F has at least a fixed point (see [9]). In this framework, let us remark that if we replace (1.5) by a weaker condition (see (1.6) below), then the conjecture stated by J.-P. PENOT in [11] as follows: *Let (X, d) be a complete metric space, $\varphi : X \rightarrow \mathbb{R}_+$ be a lower semicontinuous function and F be a multivalued operator of X into the family of all nonempty closed subsets of X satisfying the following condition:*

$$(1.6) \quad \inf\{d(x, y) : y \in F(x)\} + \inf\{\varphi(y) : y \in F(x)\} \leq \varphi(x),$$

for each $x \in X$. Then F has at least a fixed point. is false (see [9] for a counterexample). It is easy to see that (1.4) $\Rightarrow$ (1.3) $\Rightarrow$ (1.5) and (1.5) $\Rightarrow$ (1.3) provided that F has nonempty compact values.

The purpose of this paper is to present several new results in connection with the above mentioned single-valued and multi-valued Caristi type operators in complete metric spaces.

2. Main results

Let (X, d) be a metric space and $\mathcal{P}(X)$ the space of all subsets of X . We denote by $P(X)$ the space of all nonempty subsets of X and by $P_p(X)$ the set of all nonempty subsets of X having the property “ p ”, where “ p ” could be: cl = *closed*, b = *bounded*, cp = *compact*, cv =

convex (for normed spaces X), etc. We consider the following (generalized) functionals:

$$D : P(X) \times P(X) \rightarrow \mathbb{R}_+, \quad D(A, B) = \inf\{d(a, b) \mid a \in A, b \in B\}$$

$$\delta : P(X) \times P(X) \rightarrow \mathbb{R}_+ \cup \{+\infty\}, \quad \delta(A, B) = \sup\{d(a, b) \mid a \in A, b \in B\}$$

$$\rho : P(X) \times P(X) \rightarrow \mathbb{R}_+ \cup \{+\infty\}, \quad \rho(A, B) = \sup\{D(a, B) \mid a \in A\}$$

$$H : P(X) \times P(X) \rightarrow \mathbb{R}_+ \cup \{+\infty\}, \quad H(A, B) = \max\{\rho(A, B), \rho(B, A)\}$$

It is well-known that if (X, d) is a complete metric space, then $(P_{b,cl}(X), H)$ is also a complete metric space. We denote by $\text{Fix}(F)$ the fixed points set of F and by $S\text{Fix}(F)$ the strict fixed points set of F . Let (X, d) be a metric space and $F : X \rightarrow P(X)$ be a multi-valued map.

Definition 2.1. A proper function $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ is called:

- (i) a weak entropy of F if the condition (1.3) holds;
- (ii) an entropy of F if the condition (1.4) holds.

The map $F : X \rightarrow P(X)$ is said to be *weakly dissipative* iff there exists a weak entropy of F and F is said to be *dissipative* iff there is an entropy of it.

Definition 2.2. Let (X, d) be a metric space and $F : X \rightarrow P(X)$ be a multi-valued operator. Then F is said to be:

- (i) a -contraction iff there exists $a \in [0, 1[$ such that

$$H(F(x), F(y)) \leq ad(x, y), \quad \text{for each } x, y \in X;$$

- (ii) *Reich type operator* iff there exist $a, b, c \in \mathbb{R}_+$, with $a + b + c < 1$ such that

$$H(F(x), F(y)) \leq ad(x, y) + bD(x, F(x)) + cD(y, F(y)),$$

for each $x, y \in X$;

- (iii) δ -Reich type operator iff there exist $a, b, c \in \mathbb{R}_+$, with $a + b + c < 1$ such that

$$\delta(F(x), F(y)) \leq ad(x, y) + b\delta(x, F(x)) + c\delta(y, F(y)),$$

for each $x, y \in X$.

Let us remark now, that if f is a (single-valued) a -contraction in a complete metric space X , then f satisfies condition (1.1) with $\varphi(x) = (1-a)^{-1} d(x, f(x))$, for each $x \in X$, so that part of the Banach contraction principle which says about the existence of a fixed point can be obtained by Caristi's theorem. For the multi-valued case we have the following result:

Theorem 2.1. *Let (X, d) be a metric space and $F : X \rightarrow P_{cl}(X)$ be an a -contraction. Then:*

- a) *F satisfies the condition (1.5) with $\varphi(x) = (1-a)^{-1} D(x, F(x))$, for each $x \in X$.*
- b) *If, in addition $F(x) \in P_{cp}(X)$, for each $x \in X$, then F is weakly dissipative with a weak entropy given by the formula $\varphi(x) = (1-a)^{-1} D(x, F(x))$, for each $x \in X$.*

PROOF. a) is Corollary 1 in [9] and b) follows immediately from a) and the conditions (1.3) $\iff$ (1.5). $\square$

Remark 2.1. It is an open question if a multi-valued a -contraction is dissipative.

First main result of this paper is

Theorem 2.2. *Let (X, d) be a metric space and $F : X \rightarrow P_{cl}(X)$ be a Reich type multi-valued map. Then there exists $f : X \rightarrow X$ a selection of F satisfying the Caristi type condition (1.1).*

PROOF. Let $\varepsilon > 0$ such that $a < \varepsilon < 1 - b - c$. We denote by $U_x = \{y \in F(x) \mid \varepsilon d(x, y) \leq (1 - b - c)D(x, F(x))\}$, for each $x \in X$. Obviously, for each $x \in X$, the set U_x is nonempty (otherwise, if $x \in X$ is not a fixed point of F and we suppose that for each $y \in F(x)$ we have $\varepsilon d(x, y) > (1 - b - c)D(x, F(x))$, then we reach the contradiction $\varepsilon D(x, F(x)) \geq (1 - b - c)D(x, F(x))$; if $x \in X$ is a fixed point of F , then clearly $U_x \neq \emptyset$). We can choose a single-valued mapping $f : X \rightarrow X$ such that $f(x) \in U_x$, i.e. $f(x) \in F(x)$ and $\varepsilon d(x, f(x)) \leq (1 - b - c)D(x, F(x))$, for each $x \in X$. Then we have successively:

$$\begin{aligned} D(f(x), F(f(x))) &\leq H(F(x), F(f(x))) \\ &\leq ad(x, f(x)) + bD(x, F(x)) + cD(f(x), F(f(x))) \end{aligned}$$

and hence

$$(1 - c)D(f(x), F(f(x))) - bD(x, F(x)) \leq ad(x, f(x)).$$

In view of this we obtain:

$$\begin{aligned}
 d(x, f(x)) &= (\varepsilon - a)^{-1} [\varepsilon d(x, f(x)) - ad(x, f(x))] \\
 &\leq (\varepsilon - a)^{-1} [(1 - b - c)D(x, F(x)) \\
 &\quad - (1 - c)D(f(x), F(f(x))) + bD(x, F(x))] \\
 &= (1 - c)/(\varepsilon - a) [D(x, F(x)) - D(f(x), F(f(x)))].
 \end{aligned}$$

If we define $\varphi : X \rightarrow \mathbb{R}_+$ by $\varphi(x) = (1 - c)/(\varepsilon - a)D(x, F(x))$, for each $x \in X$, then it is easy to see that

$$d(x, f(x)) \leq \varphi(x) - \varphi(f(x)), \quad \text{for each } x \in X.$$

Moreover, for each $x \in X$ and each $y \in F(x)$ we have:

$$\begin{aligned}
 |\varphi(x) - \varphi(y)| &= (1 - c)/(\varepsilon - a) |D(x, F(x)) - D(y, F(y))| \\
 &\leq (1 - c)/(\varepsilon - a) |d(x, y) + H(F(x), F(y))| \\
 &\leq (1 - c)/(\varepsilon - a) [d(x, y) + ad(x, y) + bD(x, F(x)) + cD(y, F(y))] \\
 &= (1 - c)(1 + a)/(\varepsilon - a) d(x, y) + b(1 - c)/(\varepsilon - a) D(x, F(x)) \\
 &\quad + c(1 - c)/(\varepsilon - a) D(y, F(y)),
 \end{aligned}$$

proving the fact that the selection f is a kind of single-valued Reich type operator. $\square$

Remark 2.2. If the multi-valued operator $F : X \rightarrow P_{cl}(X)$ is an upper semicontinuous Reich type operator, then φ is a lower semicontinuous entropy of f (because the map $x \mapsto D(x, F(x))$ is lower semicontinuous).

Remark 2.3. If in Theorem 2.2 we take $b = c = 0$, then we obtain Theorem 5 in [10]. Moreover, we get that a multivalued a -contraction ($0 \leq a < 1$) is weakly dissipative.

Theorem 2.3. Let (X, d) be a metric space and $F : X \rightarrow P(X)$ be a δ -Reich type operator. Then the multi-valued operator F is dissipative.

PROOF. Let $\varepsilon > 0$ such that $a < \varepsilon < 1 - b - c$. Let $x \in X$ and $y \in F(x)$. It is not difficult to see that

$$\varepsilon d(x, y) \leq (1 - b - c)\delta(x, F(x)).$$

Using the fact that $y \in F(x)$ and the condition from hypothesis we have

$$\delta(y, F(y)) \leq \delta(F(x), F(y)) \leq ad(x, y) + b\delta(x, F(x)) + c\delta(y, F(y)).$$

It follows that

$$-ad(x, y) \leq b\delta(x, F(x)) - (1 - c)\delta(y, F(y)).$$

So, we have

$$\begin{aligned} d(x, y) &= (\varepsilon - a)^{-1}[\varepsilon d(x, y) - ad(x, y)] \\ &\leq (\varepsilon - a)^{-1}[(1 - b - c)\delta(x, F(x)) + b\delta(x, F(x)) - (1 - c)\delta(y, F(y))] \\ &= (1 - c)/(\varepsilon - a)[\delta(x, F(x)) - \delta(y, F(y))]. \end{aligned}$$

We define $\varphi(x) : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ as follows: $\varphi(x) = (1 - c)/(\varepsilon - a)\delta(x, F(x))$, for each $x \in X$ and we get

$$d(x, y) + \varphi(y) \leq \varphi(x), \quad \text{for each } x \in X \text{ and for all } y \in F(x),$$

i.e. the multi-valued operator F is dissipative. $\square$

An extension of Proposition 1 in [9] is as follows:

Theorem 2.4. *Let (X, d) be a complete metric space, $x_0 \in X$ be arbitrarily, $\varphi : X \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ a proper lower semi-continuous function and $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ a continuous non-decreasing function such that $\int_0^\infty \frac{ds}{1+h(s)} = +\infty$. Let $F : X \rightarrow P_{cl}(X)$ be a multi-valued operator such that:*

$$\inf \left\{ \frac{d(x, y)}{1 + h(d(x_0, x))} + \varphi(y) : y \in F(x) \right\} \leq \varphi(x),$$

for each $x \in X$. Then F has at least a fixed point.

PROOF. We shall prove that for each $x \in X$ there exists $f(x) \in F(x)$ such that:

$$\frac{d(x, f(x))}{1 + h(d(x_0, x))} + 2\varphi(f(x)) \leq 2\varphi(x).$$

If $D(x, F(x)) = 0$ then $x \in F(x)$ and we put $x = f(x)$. If $D(x, F(x)) > 0$ then we get successively:

$$\begin{aligned} &\frac{D(x, F(x))}{1 + h(d(x_0, x))} + \inf \left\{ \frac{d(x, y)}{1 + h(d(x_0, x))} + 2\varphi(y) : y \in F(x) \right\} \\ &\leq 2 \inf \left\{ \frac{d(x, y)}{1 + h(d(x_0, x))} + \varphi(y) : y \in F(x) \right\} \leq 2\varphi(x), \end{aligned}$$

for each $x \in X$. It follows that:

$$\inf \left\{ \frac{d(x, y)}{1 + h(d(x_0, x))} + 2\varphi(y) : y \in F(x) \right\} < 2\varphi(x)$$

and hence there exists $f(x) \in F(x)$ such that:

$$\frac{d(x, f(x))}{1 + h(d(x_0, x))} + 2\varphi(f(x)) \leq 2\varphi(x).$$

If we define $\psi(t) = 2\varphi(t)$ we get that f satisfies the hypothesis of Lemma 1.2 in [15] and hence there exists $x^* \in X$ such that $x^* = f(x^*) \in F(x^*)$. $\square$

In what follows we shall study the data dependence of the fixed points set of multi-valued operators which satisfy the Caristi type condition (1.3) and the data dependence of the strict fixed points set of multi-valued operators which satisfy the Caristi type condition (1.4), by giving some results.

Theorem 2.5. *Let (X, d) be a complete metric space and $F_1, F_2 : X \rightarrow P(X)$ be two multi-valued operators. We suppose that:*

- (i) *there exist two lower semi-continuous functions $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}_+$ such that for all $x \in X$, there exists $y \in F_i(x)$ so that*

$$d(x, y) \leq \varphi_i(x) - \varphi_i(y), \quad i \in \{1, 2\};$$

- (ii) *there exists $c_i \in]0, +\infty[$ such that*

$$\varphi_i(x) \leq c_i d(x, y), \quad \text{for each } x \in X \text{ and for all } y \in F_i(x), \quad i \in \{1, 2\};$$

- (iii) *there exists $\eta > 0$ such that*

$$H(F_1(x), F_2(x)) \leq \eta, \quad \text{for all } x \in X.$$

Then

$$H(\text{Fix}(F_1), \text{Fix}(F_2)) \leq \eta \max\{c_1, c_2\}.$$

PROOF. From the condition (i) we have that $\text{Fix}(F_i) \neq \emptyset$, $i \in \{1, 2\}$. Let $\varepsilon \in]0, 1[$ and $x_0 \in \text{Fix}(F_1)$. It follows, from Ekeland variational principle (see for example [8]), that there exists $x^* \in X$ such that

$$\varepsilon d(x_0, x^*) \leq \varphi_2(x_0) - \varphi_2(x^*)$$

and

$$\varphi_2(x^*) - \varphi_2(x) < \varepsilon d(x, x^*), \quad \text{for each } x \in X \setminus \{x^*\}.$$

For $x^* \in X$, there exists $y \in F_2(x^*)$ so that

$$d(x^*, y) \leq \varphi_2(x^*) - \varphi_2(y).$$

If we suppose that $y \neq x^*$, then we reach the contradiction

$$d(x^*, y) \leq \varphi_2(x^*) - \varphi_2(y) < \varepsilon d(y, x^*).$$

So $y = x^*$ and therefore $x^* \in F_2(x^*)$, i.e. $x^* \in \text{Fix}(F_2)$. Let $q \in \mathbb{R}$, $q > 1$. Then, there exists $x_1 \in F_2(x_0)$ such that

$$d(x_0, x_1) \leq qH(F_1(x_0), F_2(x_0)).$$

Taking into account the conditions (ii) and (iii) we are able to write

$$\begin{aligned} \varepsilon d(x_0, x^*) &\leq \varphi_2(x_0) - \varphi_2(x^*) = \varphi_2(x_0) \leq c_2 d(x_0, x_1) \\ &\leq c_2 q H(F_1(x_0), F_2(x_0)) \leq c_2 q \eta. \end{aligned}$$

Hence

$$d(x_0, x^*) \leq \eta c_2 q / \varepsilon.$$

Analogously, for all $y_0 \in \text{Fix}(F_2)$, there exists $y^* \in \text{Fix}(F_1)$ such that

$$d(y_0, y^*) \leq \eta c_1 q / \varepsilon.$$

Using the last two inequalities, we obtain

$$H(\text{Fix}(F_1), \text{Fix}(F_2)) \leq \eta q \varepsilon^{-1} \max\{c_1, c_2\}.$$

From this, letting $q \searrow 1$ and $\varepsilon \nearrow 1$, the conclusion follows. $\square$

Remark 2.4. In the condition (ii) of the Theorem 2.5 it is sufficient to ask that $\varphi_i(x) = 0$, for all $x \in \text{Fix}(F_i)$ and the existence of $c_i \in]0, +\infty[$ such that

$$\varphi_i(x) \leq c_i d(x, y),$$

for each $x \in \text{Fix}(F_j)$ and for all $y \in F_i(x)$, $i, j \in \{1, 2\}$, $i \neq j$.

Theorem 2.6. Let (X, d) be a complete metric space and $F : X \rightarrow P(X)$ be a multi-valued operator. We suppose that:

- (i) there exists $\varphi : X \rightarrow \mathbb{R}_+$ a lower semicontinuous function such that

$$d(x, y) \leq \varphi(x) - \varphi(y), \quad \text{for each } x \in X \text{ and for all } y \in F(x);$$

- (ii) there exists $c \in]0, +\infty[$, such that

$$\varphi(x) \leq cd(x, y), \quad \text{for each } x \in X \text{ and for all } y \in F(x).$$

Then

$$\text{Fix}(F) = S\text{Fix}(F) \neq \emptyset.$$

PROOF. From the condition (i) we have that $S\text{Fix}(F) \neq \emptyset$. Let $x^* \in \text{Fix}(F)$ and $y \in F(x^*)$. It follows that

$$d(x^*, y) \leq \varphi(x^*) - \varphi(y) = -\varphi(y) \leq 0.$$

Hence $d(x^*, y) = 0$ and therefore $y = x^*$. So $F(x^*) = \{x^*\}$, i.e. $x^* \in S\text{Fix}(F)$ and thus we are able to write that $\text{Fix}(F) \subseteq S\text{Fix}(F)$. $\square$

Remark 2.5. In the condition (ii) of the Theorem 2.6 it is sufficient to ask that $\varphi(x) = 0$, for all $x \in \text{Fix}(F)$.

Example 2.1. Let $F : [0, 1] \rightarrow P([0, 1])$, $F(x) = [x/3, x/2]$, for each $x \in [0, 1]$ and $\varphi : X \rightarrow \mathbb{R}_+$, $\varphi(x) = kx$, for each $x \in [0, 1]$, where $k \in \mathbb{R}$, $k \geq 1$. It is not difficult to see that $|x - y| \leq \varphi(x) - \varphi(y)$, for each $x \in [0, 1]$ and for all $y \in F(x)$ and there exists $c = 2k > 0$ such that $\varphi(x) \leq c|x - y|$ for each $x \in [0, 1]$ and for all $y \in F(x)$. From Theorem 2.6 we have $\text{Fix}(F) = S\text{Fix}(F) \neq \emptyset$.

Theorem 2.7. Let (X, d) be a complete metric space and $F_1, F_2 : X \rightarrow P(X)$ be two multi-valued operators. We suppose that:

- (i) there exist two lower semicontinuous functions $\varphi_1, \varphi_2 : X \rightarrow \mathbb{R}_+$ such that

$$d(x, y) \leq \varphi_i(x) - \varphi_i(y),$$

for each $x \in X$ and for all $y \in F_i(x)$, $i \in \{1, 2\}$;

(ii) there exists $c_i \in]0, +\infty[$ such that

$$\varphi_i(x) \leq c_i d(x, y), \quad \text{for each } x \in X \text{ and for all } y \in F_i(x), \quad i \in \{1, 2\};$$

(iii) there exists $\eta > 0$ such that

$$H(F_1(x), F_2(x)) \leq \eta, \quad \text{for all } x \in X.$$

Then

$$H(\text{Fix}(F_1), \text{Fix}(F_2)) = H(S \text{Fix}(F_1), S \text{Fix}(F_2)) \leq \eta \max\{c_1, c_2\}.$$

PROOF. From the Theorem 2.6 we have $\text{Fix}(F_i) = S \text{Fix}(F_i) \neq \emptyset$, $i \in \{1, 2\}$ and applying Theorem 2.5 the conclusion follows. $\square$

We mention that the Remark 2.4 can be made for this theorem, too.

Example 2.2. Let $F_1, F_2 : [0, 1] \rightarrow P([0, 1])$, $F_1(x) = [x/3, x/2]$, for each $x \in [0, 1]$ and $F_2(x) = [(x+1)/2, (x+2)/3]$, for each $x \in [0, 1]$. Let $\varphi_1, \varphi_2 : [0, 1] \rightarrow \mathbb{R}_+$, $\varphi_1(x) = x$, for each $x \in [0, 1]$ and $\varphi_2(x) = 1 - x$, for each $x \in [0, 1]$. By an easy calculation we get that $|x - y| \leq \varphi_i(x) - \varphi_i(y)$, for each $x \in [0, 1]$ and for all $y \in F_i(x)$, $i \in \{1, 2\}$ and there exist $c_1 = 2$ and $c_2 = 2$ such that $\varphi_i(x) \leq c_i |x - y|$, for each $x \in [0, 1]$ and for all $y \in F_i(x)$, $i \in \{1, 2\}$. Also, there exists $\eta = 2/3 > 0$ so that $H(F_1(x), F_2(x)) \leq \eta$, for all $x \in [0, 1]$. Then, from Theorem 2.7 we have $H(\text{Fix}(F_1), \text{Fix}(F_2)) = H(S \text{Fix}(F_1), S \text{Fix}(F_2)) \leq 4/3$.

References

- [1] J.-P. AUBIN, Optima and equilibria, An introduction to nonlinear analysis, *Springer-Verlag, Berlin*, 1993.
- [2] J.-P. AUBIN and H. FRANKOWSKA, Set-valued analysis, Systems and control, *Birkhäuser, Basel*, 1990.
- [3] J.-P. AUBIN and J. SIEGEL, Fixed points and stationary points of dissipative multivalued maps, *Proc. A. M. S.* **78** (1980), 391–398.
- [4] J. S. BAE, E. W. CHO and S. H. YEOM, A generalization of the Caristi–Kirk fixed point theorem and its applications to mapping theorem, *J. Korean Math. Soc.* **31** (1994), 29–48.
- [5] J. CARISTI, Fixed points theorems for mappings satisfying inwardness conditions, *Trans. A. M. S.* **215** (1976), 241–251.

- [6] L. B. CIRIC, A generalization of Caristi's fixed point theorem, *Math. Pannonica* **3** (1992), 51–57.
- [7] K. DEIMLING, Multivalued differential equations, *Walter de Gruyter*, 1992.
- [8] D. G. DE FIGUEIREDO, The Ekeland variational principle with applications and detours, *Springer-Verlag, Berlin*, 1989.
- [9] L. VAN HOT, Fixed point theorems for multivalued mappings, *Comment. Math. Univ. Carolinae* **23** (1982), 137–145.
- [10] J. R. JACHYMSKI, Caristi's fixed point theorem and selections of set-valued contractions, *J. Math. Anal. Appl.* **227** (1998), 55–67.
- [11] J.-P. PENOT, A short constructive proof of Caristi's fixed point theorem, (*preprint*).
- [12] A. PETRUȘEL and A. SÎNTĂMĂRIAN, On Caristi-type operators, Tiberiu Popoviciu Itinerant Seminar of Functional Equations, Approximation and Convexity, *Cluj Napoca, Romania*, May 22–26, 2001, 181–190.
- [13] I. A. RUS, Generalized contractions and applications, *Cluj University Press, Cluj Napoca*, 2001.
- [14] I. A. RUS and S. MUREȘAN, Data dependence of the fixed points set of some weakly Picard operators, Tiberiu Popoviciu Itinerant Seminar of Functional Equations, Approximation and Convexity, *Cluj-Napoca, Romania*, May 23–29, 2000, 201–207.
- [15] C.-K. ZHONG, J. ZHU and P.-H. ZHAO, An extension of multi-valued contraction mappings and fixed points, *Proc. A. M. S.* **128** (2000), 2439–2444.

ADRIAN PETRUȘEL
DEPARTMENT OF APPLIED MATHEMATICS
"BABEȘ-BOLYAI" UNIVERSITY
STR. M. KOGĂLNICEANU NR. 1
3400 CLUJ-NAPOCA
ROMANIA

E-mail: petrusel@math.ubbcluj.ro

ALINA SÎNTĂMĂRIAN
DEPARTMENT OF APPLIED MATHEMATICS
"BABEȘ-BOLYAI" UNIVERSITY
STR. M. KOGĂLNICEANU NR. 1
3400 CLUJ-NAPOCA
ROMANIA

E-mail: salina@math.ubbcluj.ro

(Received April 17, 2001; revised July 30, 2001)