

Monomials and binomials over finite fields as $\mathcal{R}$ -orthomorphisms

By WUN-SENG CHOU (Taipei) and HARALD NIEDERREITER (Singapore)

Abstract. We give criteria for both monomials and binomials of the form $ax^{(q+1)/2} + bx$ to be $\mathcal{R}$ -orthomorphisms of the finite field F_q of odd order q . We also prove existence theorems for $\mathcal{R}$ -orthomorphisms of this form.

1. Introduction

Throughout this paper, q is a prime power and F_q is the finite field of order q . The polynomial $f(x) \in F_q[x]$ is called a *permutation polynomial* of F_q if the function $\sigma : F_q \longrightarrow F_q$, defined by $\sigma : a \longmapsto f(a)$, is a permutation of F_q . Permutation polynomials of finite fields have been studied extensively (see [4, Chapter 7]). One important and useful class of permutation polynomials is the class of so-called orthomorphisms. Recall that $f(x)$ is an *orthomorphism* of F_q if both $f(x)$ and $f(x) - x$ are permutation polynomials of F_q . Orthomorphisms have interesting applications, for instance to the construction of orthogonal Latin squares (see [9, Chapter 22]) and cryptology (see [8]). Orthomorphisms of F_q are also closely connected with *complete mapping polynomials* of F_q , since $f(x) \in F_q[x]$ is an orthomorphism of F_q if and only if $-f(x)$ is a complete mapping polynomial of F_q . Complete mapping polynomials of F_q were first studied by NIEDERREITER and ROBINSON [6], [7].

In this paper we consider a special class of orthomorphisms. Let $\mathcal{R}$ be a nonempty set of positive integers. Then $f(x)$ is called an *$\mathcal{R}$ -orthomorphism* of F_q if each polynomial $f^{(r)}(x)$ is an orthomorphism of F_q ,

Mathematics Subject Classification: 11T06.

Key words and phrases: permutation polynomials, complete mapping polynomials, orthomorphisms.

where $r \in \mathcal{R}$ and $f^{(r)}(x)$ is the r th iterated composition of $f(x)$ with itself. The definition of $\mathcal{R}$ -orthomorphisms is given explicitly by COHEN, NIEDERREITER, SHPARLINSKI, and ZIEVE [1]. They also present examples of both linearized and sublinearized polynomials that are $\mathcal{R}$ -orthomorphisms. In fact, some special types of $\mathcal{R}$ -orthomorphisms have been used in combinatorial design theory (see [2]).

In Section 2 we study monomials as $\mathcal{R}$ -orthomorphisms. Especially, we consider the monomial $f(x) = ax^{(q+1)/2}$ as a concrete example, where q is odd. We show that for any positive integer k and any sufficiently large $q \equiv 1 \pmod{4}$ there exists an $\mathcal{R}_k$ -orthomorphism of F_q of this form, where $\mathcal{R}_k = \{1, 2, \dots, k\}$. We study the special kind of binomials $f(x) = ax^{(q+1)/2} + bx$ in Section 3. We show in this section that if $\mathcal{R}$ is finite and q is sufficiently large, then there is at least one ordered pair $(a, b) \in F_q^* \times F_q^*$ such that the polynomial $f(x) = ax^{(q+1)/2} + bx$ is an $\mathcal{R}$ -orthomorphism of F_q . Here the lower bound on q is smaller than that in a comparable result in [1, Theorem 3].

2. Monomials as $\mathcal{R}$ -orthomorphisms

In this section, $f(x)$ is a monomial $f(x) = ax^n \in F_q[x]$ with $a \neq 0$. Then, for any positive integer i , we have $f^{(i)}(x) = a^{n^{i-1} + \dots + n + 1} x^{n^i}$. So, $f^{(i)}(x)$ is a permutation polynomial of F_q if and only if $\gcd(n^i, q-1) = 1$ (and thus, $\gcd(n, q-1) = 1$). The following is a criterion for $f(x)$ to be an $\mathcal{R}$ -orthomorphism. By the above remarks, the proof is obvious.

Lemma 2.1. *The monomial $f(x) = ax^n \in F_q[x]$, $a \neq 0$, is an $\mathcal{R}$ -orthomorphism of F_q if and only if $\gcd(n, q-1) = 1$ and for each $m \in \mathcal{R}$, the equation*

$$\frac{x^{n^m} - y^{n^m}}{x - y} = a^{-(n^{m-1} + \dots + n + 1)}$$

has no solution in $F_q \times F_q \setminus \{(\alpha, \alpha) : \alpha \in F_q\}$.

From this lemma, it is trivial that for $n = 1$, $f(x) = ax \in F_q[x]$ is an $\mathcal{R}$ -orthomorphism of F_q if and only if $a \neq 0, 1$ and $\mathcal{R}$ contains no multiple of the (multiplicative) order $\text{ord}(a)$ of a . So, if $\mathcal{R} = \mathcal{R}_k = \{1, 2, \dots, k\}$, then $f(x) = ax$, $a \neq 0$, is an $\mathcal{R}_k$ -orthomorphism of F_q if and only if $1 \leq k < \text{ord}(a)$. For $n > 1$, let $e_m = \gcd(n^{m-1} + \dots + n + 1, q-1)$ and $d_m = \gcd(n-1, \frac{q-1}{e_m})$.

Lemma 2.2. *Let $f(x) = ax^n \in F_q[x]$ with $a \neq 0$ and $n > 1$ and let m be a positive integer. Then $f^{(m)}(x) - x$ has a root in F_q^* if and only if $\text{ord}(a)$ divides $(q-1)/d_m$.*

PROOF. The polynomial $f^{(m)}(x) - x$ has a root in F_q^* if and only if

$$a^{n^{m-1}+n^{m-2}+\dots+n+1}x^{n^m-1} = 1$$

has a root in F_q^* . If g is a primitive element of F_q and $a = g^s$ for some $s > 0$, then the last statement is equivalent to the existence of a positive integer t satisfying

$$1 = g^{s(n^{m-1}+\dots+n+1)+t(n^m-1)} = g^{(n^{m-1}+\dots+n+1)(s+t(n-1))}.$$

Thus, $f^{(m)}(x) - x$ having a root in F_q^* is equivalent to $s + y(n-1) \equiv 0 \pmod{(q-1)/e_m}$ having a solution t , which is equivalent to $d_m \mid s$, and thus $\text{ord}(a)$ dividing $(q-1)/d_m$. $\square$

The above lemma gives a necessary condition for a monomial to be an $\mathcal{R}$ -orthomorphism.

Corollary 2.3. *If $f(x) = ax^n \in F_q[x]$, with $n > 1$ and $a \neq 0$, is an $\mathcal{R}$ -orthomorphism of F_q , then for all $m \in \mathcal{R}$, $\text{ord}(a)$ does not divide $(q-1)/d_m$.*

We omit the proof because it is obvious. In the following, we consider the special kind of monomial $f(x) = ax^{(q+1)/2} \in F_q[x]$, $a \neq 0$, q odd. Moreover, we take $\mathcal{R} = \mathcal{R}_k$ with a positive integer k . We first establish the following result on the iterates $f^{(r)}(x)$.

Lemma 2.4. *Let $f(x) = ax^{(q+1)/2} \in F_q[x]$ with $a \in F_q^*$ and q odd. Then, as functions on F_q , the iterates of f are given by*

$$\begin{aligned} f^{(4s+1)}(x) &= a^{4s+1}x^{(q+1)/2}, \\ f^{(4s+2)}(x) &= a^{(q-1)/2+4s+2}x, \\ f^{(4s+3)}(x) &= a^{(q-1)/2+4s+3}x^{(q+1)/2}, \\ f^{(4(s+1))}(x) &= a^{4(s+1)}x, \end{aligned}$$

for any nonnegative integer s .

PROOF. This is shown by straightforward induction, using the fact that $x^q = x$ as a function on F_q . $\square$

Notice that $f(x)$ is a permutation polynomial of F_q if and only if $q \equiv 1 \pmod{4}$, because we need $\gcd((q+1)/2, q-1) = 1$. From now on in this section, we assume that $q \equiv 1 \pmod{4}$.

It follows from Lemma 2.4 that for an even positive integer m , $f^{(m)}(x) - x$ is a permutation polynomial of F_q if and only if $\text{ord}(a)$ does not divide m if $m \equiv 0 \pmod{4}$, or $m + \frac{q-1}{2}$ if $m \equiv 2 \pmod{4}$. Notice that if $c_m = \gcd(m, q-1)$, then $\gcd(q-1, \frac{q-1}{2} + m) = c_m$ if $q \equiv 1 \pmod{8}$, and $\gcd(q-1, \frac{q-1}{2} + m) = 2c_m$ if $q \equiv 5 \pmod{8}$.

For the consideration of $f^{(m)}(x) - x$ for odd positive integers m , we recall the following result from [4, Theorem 7.11]. We denote by η the quadratic character of F_q , with the convention $\eta(0) = 0$.

Lemma 2.5. *For odd q , the polynomial $x^{(q+1)/2} + bx \in F_q[x]$ is a permutation polynomial of F_q if and only if $\eta(b^2 - 1) = 1$.*

Thus, for an odd positive integer m , it follows from Lemmas 2.4 and 2.5 that $f^{(m)}(x) - x$ is a permutation polynomial of F_q if and only if $\eta(a^{2m} - 1) = 1$.

Theorem 2.6. *Let q be a prime power with $q \equiv 1 \pmod{4}$, let k be a positive integer, and let $\mathcal{R}_k = \{1, 2, \dots, k\}$. Suppose that*

$$q \geq 2^{\lfloor (k-3)/2 \rfloor} (k+1)^2 q^{1/2} + 2^{\lfloor (k-7)/2 \rfloor} (5k^2 + 12) + 1.$$

Then there exists an $a \in F_q^$ such that $f(x) = ax^{(q+1)/2} \in F_q[x]$ is an $\mathcal{R}_k$ -orthomorphism of F_q .*

PROOF. For an odd positive integer m , define $g_m(x) = x^{2m} - 1$. For a positive integer m with $m \equiv 0 \pmod{4}$, define $h_{0,m}(x) = x^{c_m} - 1$; and for $m \equiv 2 \pmod{4}$, define $h_{2,m}(x)$ to be either $h_{2,m}(x) = x^{c_m} - 1$ if $q \equiv 1 \pmod{8}$, or $h_{2,m}(x) = x^{2c_m} - 1$ if $q \equiv 5 \pmod{8}$. Now let $\mathfrak{A}_o$ be the set of roots in F_q of all polynomials of the form $g_m(x)$ for odd integers m with $1 \leq m \leq k$. It is easy to see that

$$|\mathfrak{A}_o| \leq 2 + \sum_{\substack{m=1 \\ m \text{ odd}}}^k (2m-2) \leq \frac{k^2+3}{2}.$$

Also, let $\mathfrak{A}_e$ be the set of roots in F_q of all polynomials $h_{0,m}(x)$ and $h_{2,m}(x)$ for even integers m with $1 \leq m \leq k$. Then it is not difficult to see that

$$|\mathfrak{A}_e \setminus \mathfrak{A}_o| \leq \sum_{\substack{m=1 \\ m \equiv 0 \pmod{4}}}^k (m-2) \leq \frac{k^2}{8}.$$

Let N be the number of elements $a \in F_q^*$ such that $f(x) = ax^{(q+1)/2}$ is an $\mathcal{R}_k$ -orthomorphism. Then

$$\begin{aligned} N &= \frac{1}{2^{\lfloor \frac{k+1}{2} \rfloor}} \sum_{a \in F_q^* \setminus (\mathfrak{A}_o \cup \mathfrak{A}_e)} \prod_{\substack{m=1 \\ m \text{ odd}}}^k (1 + \eta(a^{2m} - 1)) \\ &= \frac{1}{2^{\lfloor \frac{k+1}{2} \rfloor}} \left(\sum_{a \in F_q^*} \prod_{\substack{m=1 \\ m \text{ odd}}}^k (1 + \eta(a^{2m} - 1)) - \sum_{a \in \mathfrak{A}_o \cup \mathfrak{A}_e} \prod_{\substack{m=1 \\ m \text{ odd}}}^k (1 + \eta(a^{2m} - 1)) \right), \end{aligned}$$

and so

$$N \geq \frac{N_1}{2^{\lfloor (k+1)/2 \rfloor}} - \frac{1}{2} |\mathfrak{A}_o \cup \mathfrak{A}_e| \geq \frac{N_1}{2^{\lfloor (k+1)/2 \rfloor}} - \frac{5k^2 + 12}{16}$$

with

$$N_1 := \sum_{a \in F_q^*} \prod_{\substack{m=1 \\ m \text{ odd}}}^k (1 + \eta(a^{2m} - 1)).$$

We can write

$$(2.1) \quad N_1 = q - 1 + \sum_{r=1}^{\lfloor (k+1)/2 \rfloor} \sum_{\substack{1 \leq m_1 < \dots < m_r \leq k \\ m_j \text{ odd}}} \sum_{a \in F_q^*} \eta \left(\prod_{j=1}^r (a^{2m_j} - 1) \right).$$

Consider the innermost sum on the right-hand side of (2.1). If the polynomial $\prod_{j=1}^r (x^{2m_j} - 1)$ is a square in $F_q[x]$, then the corresponding sum is clearly nonnegative. Otherwise by the Weil bound [4, Theorem 5.41],

$$\left| \sum_{a \in F_q^*} \eta \left(\prod_{j=1}^r (a^{2m_j} - 1) \right) \right| < 2q^{1/2} \sum_{j=1}^r m_j.$$

Therefore

$$N_1 > q - 1 - 2q^{1/2} \sum_{r=1}^{\lfloor (k+1)/2 \rfloor} \sum_{\substack{1 \leq m_1 < \dots < m_r \leq k \\ m_j \text{ odd}}} \sum_{j=1}^r m_j$$

$$\begin{aligned}
&= q - 1 - 2^{\lfloor (k+1)/2 \rfloor} q^{1/2} \sum_{\substack{m=1 \\ m \text{ odd}}}^k m \\
&\geq q - 1 - 2^{\lfloor (k+1)/2 \rfloor} q^{1/2} \frac{(k+1)^2}{4}.
\end{aligned}$$

Altogether,

$$N > \frac{q-1}{2^{\lfloor (k+1)/2 \rfloor}} - \frac{(k+1)^2}{4} q^{1/2} - \frac{5k^2+12}{16},$$

and the desired result follows. $\square$

3. $\mathcal{R}$ -orthomorphisms of the form $ax^{(q+1)/2} + bx$

In this section, we consider q odd and polynomials of the form $f(x) = ax^{(q+1)/2} + bx \in F_q[x]$ with $ab \neq 0$. It follows from Lemma 2.5 that $f(x)$ is a permutation polynomial of F_q if and only if $\eta(b^2 - a^2) = 1$. Note that f is a linear function when restricted to the squares in F_q and another linear function when restricted to the nonsquares in F_q . This observation and induction yield the following formulas for the iterates of f as functions on F_q . If $\eta(b+a) = \eta(b-a) = 1$, then for each positive integer n we have

$$f^{(n)}(x) = \frac{(b+a)^n - (b-a)^n}{2} x^{(q+1)/2} + \frac{(b+a)^n + (b-a)^n}{2} x.$$

If $\eta(b+a) = \eta(b-a) = -1$, then for each positive integer n we have

$$f^{(n)}(x) = (b^2 - a^2)^{(n-1)/2} ax^{(q+1)/2} + (b^2 - a^2)^{(n-1)/2} bx$$

whenever n is odd and

$$f^{(n)}(x) = (b^2 - a^2)^{n/2} x$$

whenever n is even. From these formulas, the following criterion is trivial.

Lemma 3.1. *Let $\mathcal{R}$ be a nonempty set of positive integers. Then the polynomial $f(x) = ax^{(q+1)/2} + bx \in F_q[x]$ with $ab \neq 0$ is an $\mathcal{R}$ -orthomorphism of F_q if and only if one of the following two conditions holds:*

- (i) $\eta(b+a) = \eta(b-a) = 1$ and $\eta((b+a)^m - 1)((b-a)^m - 1) = 1$ for all $m \in \mathcal{R}$;

- (ii) $\eta(b+a) = \eta(b-a) = -1$ and for all $m \in \mathcal{R}$, $(b^2 - a^2)^{m/2} \neq 1$ if m is even and $\eta(((b^2 - a^2)^{(m-1)/2}(b+a) - 1)((b^2 - a^2)^{(m-1)/2}(b-a) - 1)) = 1$ if m is odd.

Using this lemma, we have the following counting formula which is closely related to a result of MENDELSON and WOLK [5] (see also EVANS [3]) for the special case where q is a prime.

Corollary 3.2. *If $q \equiv 3 \pmod{4}$, then there are exactly $\frac{(q-3)(q-5)}{4}$ orthomorphisms of F_q of the form $f(x) = ax^{(q+1)/2} + bx \in F_q[x]$ with $ab \neq 0$. If $q \equiv 1 \pmod{4}$, then there are exactly $\frac{(q-5)^2}{4}$ orthomorphisms of F_q of the form $f(x) = ax^{(q+1)/2} + bx \in F_q[x]$ with $ab \neq 0$.*

PROOF. From Lemma 3.1, for $a, b \in F_q^*$, the polynomial $f(x) = ax^{(q+1)/2} + bx$ is an orthomorphism of F_q if and only if $\eta((b-a)(b+a)) = 1$ and $\eta((b-a-1)(b+a-1)) = 1$. Let N be the number of orthomorphisms counted in the corollary. After the substitution $u = b-a$ and $v = b+a$, we see that N is the number of ordered pairs $(u, v) \in F_q \times F_q$ with $u \neq \pm v$, $\eta(uv) = 1$, $\eta((u-1)(v-1)) = 1$. We can restrict u and v to $G_q := F_q \setminus \{0, 1\}$. Then we can write

$$(3.1) \quad N = N_1 - N_2,$$

where N_1 is the number of ordered pairs $(u, v) \in G_q \times G_q$ with $\eta(uv) = 1$, $\eta((u-1)(v-1)) = 1$, and N_2 is the number of ordered pairs $(u, v) \in G_q \times G_q$ with $u = \pm v$, $\eta(uv) = 1$, $\eta((u-1)(v-1)) = 1$. We have

$$\begin{aligned} N_1 &= \frac{1}{4} \sum_{u, v \in G_q} [1 + \eta(uv)][1 + \eta((u-1)(v-1))] \\ &= \frac{(q-2)^2}{4} + \frac{1}{4} \sum_{u, v \in G_q} \eta(uv) + \frac{1}{4} \sum_{u, v \in G_q} \eta((u-1)(v-1)) \\ &\quad + \frac{1}{4} \sum_{u, v \in G_q} \eta(uv)\eta((u-1)(v-1)) \\ &= \frac{(q-2)^2}{4} + \frac{1}{4} \left(\sum_{u \in G_q} \eta(u) \right)^2 + \frac{1}{4} \left(\sum_{u \in G_q} \eta(u-1) \right)^2 \\ &\quad + \frac{1}{4} \left(\sum_{u \in G_q} \eta(u(u-1)) \right)^2 \end{aligned}$$

$$= \frac{(q-2)^2}{4} + \frac{1}{4} + \frac{1}{4} + \frac{1}{4} = \frac{q^2 - 4q + 7}{4},$$

where we used [4, Theorem 5.48] to evaluate the last character sum.

Now we consider N_2 . All ordered pairs $(u, v) \in G_q \times G_q$ with $u = v$ are counted for N_2 and this gives $q - 2$ ordered pairs. If $u = -v$, then the conditions become $\eta(-v^2) = 1$, $\eta(1 - v^2) = 1$. This is only possible if $q \equiv 1 \pmod{4}$, since only then $\eta(-1) = 1$. In this case, it remains to count the number N_3 of $v \in G_q$ with $\eta(1 - v^2) = 1$. We have

$$\begin{aligned} N_3 &= \frac{1}{2} \sum_{v \in G_q} (1 + \eta(1 - v^2)) - \frac{1}{2} = \frac{q-3}{2} + \frac{1}{2} \sum_{v \in G_q} \eta(1 - v^2) \\ &= \frac{q-3}{2} + \frac{1}{2} \sum_{v \in F_q} \eta(1 - v^2) - \frac{1}{2} = \frac{q-5}{2}, \end{aligned}$$

where we again used [4, Theorem 5.48]. Thus, if $q \equiv 1 \pmod{4}$, then

$$N_2 = q - 2 + \frac{q-5}{2} = \frac{3q-9}{2},$$

whereas $N_2 = q - 2$ if $q \equiv 3 \pmod{4}$. Recalling (3.1), we get the claimed result. $\square$

The following theorem shows the existence of $\mathcal{R}$ -orthomorphisms of F_q of the form $f(x) = ax^{(q+1)/2} + bx \in F_q[x]$ for sufficiently large q . The condition on q in this result is less restrictive than that in the comparable result in [1, Theorem 3].

Theorem 3.3. *Let $\mathcal{R}$ be a finite nonempty set of positive integers and q an odd prime power with*

$$q \geq 2^{R+2} \left(2 + \sum_{m \in \mathcal{R}} m \right),$$

where R is the cardinality of $\mathcal{R}$. Then there exists at least one ordered pair $(a, b) \in F_q^* \times F_q^*$ such that the polynomial $f(x) = ax^{(q+1)/2} + bx$ is an $\mathcal{R}$ -orthomorphism of F_q .

PROOF. Let N be the number of ordered pairs $(a, b) \in F_q^* \times F_q^*$ such that the polynomial $f(x) = ax^{(q+1)/2} + bx$ is an $\mathcal{R}$ -orthomorphism

of F_q . Let N_1 be the number of ordered pairs $(a, b) \in F_q \times F_q$ such that $\eta(b+a) = \eta(b-a) = 1$ and

$$\eta(((b+a)^m - 1)((b-a)^m - 1)) = 1 \quad \text{for all } m \in \mathcal{R}.$$

By using only condition (i) in Lemma 3.1, we see that

$$N_1 \leq N + \#\{(a, b) \in F_q \times F_q : ab = 0, \eta(b+a) = \eta(b-a) = 1\},$$

and so

$$(3.2) \quad N \geq N_1 - q + 1.$$

Let $C := \{(a, b) \in F_q \times F_q : b = \pm a\}$ and D be the set of $(a, b) \in F_q \times F_q$ with

$$((b+a)^m - 1)((b-a)^m - 1) = 0 \quad \text{for some } m \in \mathcal{R}.$$

Then

$$\begin{aligned} N_1 &= \frac{1}{2^{R+2}} \sum_{\substack{(a,b) \in F_q \times F_q \\ (a,b) \notin C \cup D}} [1 + \eta(b+a)][1 + \eta(b-a)] \\ &\quad \cdot \prod_{m \in \mathcal{R}} [1 + \eta(((b+a)^m - 1)((b-a)^m - 1))] \\ &\geq \frac{S}{2^{R+2}} - \frac{1}{2} |C \cup D| \end{aligned}$$

with

$$S := \sum_{a,b \in F_q} [1 + \eta(b+a)][1 + \eta(b-a)] \prod_{m \in \mathcal{R}} [1 + \eta(((b+a)^m - 1)((b-a)^m - 1))].$$

By carrying out the substitution $u = b+a$ and $v = b-a$ in the sum S , we

obtain

$$\begin{aligned}
S &= \sum_{u,v \in F_q} (1 + \eta(u))(1 + \eta(v)) \prod_{m \in \mathcal{R}} (1 + \eta((u^m - 1)(v^m - 1))) \\
&= \sum_{u,v \in F_q} (1 + \eta(u))(1 + \eta(v)) \\
&\quad + \sum_{r=1}^R \sum_{\substack{m_1 < m_2 < \dots < m_r \\ m_j \in \mathcal{R}}} \sum_{u,v \in F_q} (1 + \eta(u))(1 + \eta(v)) \prod_{j=1}^r \eta((u^{m_j} - 1)(v^{m_j} - 1)) \\
&= \left(\sum_{u \in F_q} (1 + \eta(u)) \right)^2 \\
&\quad + \sum_{r=1}^R \sum_{\substack{m_1 < m_2 < \dots < m_r \\ m_j \in \mathcal{R}}} \left(\sum_{u \in F_q} (1 + \eta(u)) \prod_{j=1}^r \eta(u^{m_j} - 1) \right)^2 \geq q^2.
\end{aligned}$$

In view of (3.2), this yields

$$N \geq \frac{q^2}{2^{R+2}} - \frac{1}{2}|C| - \frac{1}{2}|D| - q + 1.$$

It is clear that $|C| = 2q - 1$. Furthermore, $|D|$ is the number of $(u, v) \in F_q \times F_q$ with $(u^m - 1)(v^m - 1) = 0$ for some $m \in \mathcal{R}$. Therefore

$$|D| \leq 2q \sum_{m \in \mathcal{R}} m.$$

Altogether, we get

$$N \geq \left(\frac{q}{2^{R+2}} - 2 - \sum_{m \in \mathcal{R}} m \right) q + \frac{3}{2},$$

and the desired result follows. $\square$

Acknowledgements. This paper was partially supported by the National Science Council in Taiwan, ROC, under the grant number 88-2115-M-001-002. The second author is grateful to the Academia Sinica in Taipei for supporting his visit during which this work was initiated. The authors thank an anonymous referee for several useful suggestions on the presentation of the material.

References

- [1] S. D. COHEN, H. NIEDERREITER, I. E. SHPARLINSKI and M. ZIEVE, Incomplete character sums and a special class of permutations, *J. Théorie des Nombres Bordeaux* **13** (2001), 53–63.
- [2] J. DÉNES and P. J. OWENS, Some new Latin power sets not based on groups, *J. Combinatorial Theory Ser. A* **85** (1999), 69–82.
- [3] A. B. EVANS, Orthomorphisms of Z_p , *Discrete Math.* **64** (1987), 147–156.
- [4] R. LIDL and H. NIEDERREITER, Finite Fields, *Cambridge Univ. Press, Cambridge*, 1997.
- [5] N. S. MENDELSON and B. WOLK, A search for a nondesarguesian plane of prime order, Proc. Internat. Conf. on Finite Geometry (Winnipeg, 1984), Lecture Notes in Pure and Applied Math., vol. 103, *Marcel Dekker, New York*, 1985, 199–208.
- [6] H. NIEDERREITER and K. H. ROBINSON, Bol loops of order pq , *Math. Proc. Cambridge Philos. Soc.* **89** (1981), 241–256.
- [7] H. NIEDERREITER and K. H. ROBINSON, Complete mappings of finite fields, *J. Austral. Math. Soc. Ser. A* **33** (1982), 197–212.
- [8] C. P. SCHNORR and S. VAUDENAY, Black box cryptanalysis of hash networks based on multipermutations, Advances in Cryptology – EUROCRYPT '94 (A. De Santis, ed.), Lecture Notes in Computer Science, vol. 950, *Springer, Berlin*, 1995, 47–57.
- [9] J. H. VAN LINT and R. M. WILSON, A Course in Combinatorics, *Cambridge Univ. Press, Cambridge*, 1992.

WUN-SENG CHOU
 INSTITUTE OF MATHEMATICS
 ACADEMIA SINICA
 NANKANG, TAIPEI 11529
 TAIWAN, ROC
E-mail: macws@ccvax.sinica.edu.tw

HARALD NIEDERREITER
 DEPARTMENT OF MATHEMATICS
 NATIONAL UNIVERSITY OF SINGAPORE
 2 SCIENCE DRIVE 2
 SINGAPORE 117543
 REPUBLIC OF SINGAPORE
E-mail: nied@math.nus.edu.sg

(Received October 3, 2001; revised February 12, 2002)