

Another characterization of the gamma function

By DETLEF GRONAU (Graz) and JANUSZ MATKOWSKI (Zielona Góra)

Abstract. The function $\frac{\log \Gamma(x)}{\log x}$ is characterized to be the only convex solution of the functional equation

$$f(x+1) = \frac{\log x}{\log(x+1)}(f(x) + 1), \quad x \in (0, \infty).$$

Some relations to the function $\log \Gamma(x+1)/x^a$, $0 < a \leq 1$ are shown.

0. Introduction

In this paper we examine the behavior of the Euler gamma function Γ in the logarithmically scaled coordinate system. More exactly, we show that the function $g : (0, \infty) \rightarrow \mathbb{R}$ defined by

$$g(x) := \begin{cases} \frac{\log \Gamma(x)}{\log x} & \text{for } x \neq 1, \\ -\gamma & \text{for } x = 1, \end{cases}$$

(where γ is the Euler gamma constant) is increasing, convex, $g(0+) = -1$ and $g(2) = 0$. The main result of our paper states that the function g is the only convex solution of the functional equation

$$f(x+1) = \frac{\log x}{\log(x+1)}(f(x) + 1), \quad x \in (0, \infty). \quad (1)$$

Mathematics Subject Classification: 33B15, 39A13.

Key words and phrases: gamma function, Bohr–Møllerup theorem.

One can weaken the supposition of convexity of the solution in this way that only convexity is supposed in a neighborhood of infinity.

Note that no initial condition is required.


We would like to remark that this characterization of the gamma function is not a consequence of the famous Bohr–Møllerup characterization of the gamma function (see e.g. [2] or [3], p. 288) as the only log-convex solution of the functional equation

$$f(x+1) = x \cdot f(x), \quad x \in (0, \infty); \quad f(1) = 1. \quad (2)$$

And, the more, it cannot be derived from the recent generalization of the Bohr–Møllerup theorem [5] that says that the gamma function is the only solution of (2), which is geometrically convex on a neighborhood of infinity. In these characterizations the initial condition is indispensable.

As an interesting consequence we infer that the function $G(x) = \frac{\log \Gamma(\exp x)}{\log x}$ is strictly increasing and strictly convex on $\mathbb{R}$.

We further consider the functions $\frac{\log \Gamma(x+1)}{\log x^\alpha}$ and $\frac{\log \Gamma(x)}{\log x^\alpha}$ for a fixed real α , $0 < \alpha \leq 1$, relating it with a recent paper of GRABNER *et al.* [4].



1. Some properties of $\log \Gamma(x)/\log x$

The function $g(x) := \frac{\log \Gamma(x)}{\log x}$ is analytic on $(0, \infty)$ with a removable singularity at $x = 1$, $g(1) = -0.577215 \dots = -\gamma$ (where $\gamma = \lim_{n \rightarrow \infty} (\sum_{k=1}^n \frac{1}{k} - \log n)$, the Euler gamma constant). We further have $g(0+) = \lim_{x \rightarrow 0+} g(x) = -1$, and $g'_+(0) = 0$.


To show this the following representations are useful (see [3], p. 287f.):

$$\log \Gamma(x) = -\log x - \gamma x - \sum_{n=1}^{\infty} \left(\log \left(1 + \frac{x}{n} \right) - \frac{x}{n} \right), \quad x > 0;$$

$$\Psi(x) = \frac{d}{dx} \log \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)} = -\frac{1}{x} - \gamma + \sum_{n=1}^{\infty} \frac{x}{(x+n)n}, \quad x > 0.$$

Proposition. *The function $g(x) = \frac{\log \Gamma(x)}{\log x}$ is strictly monotone increasing and strictly convex on $(0, \infty)$.*

PROOF. We show that g' and g'' are positive on $(0, \infty)$. To do this we use asymptotic expansions for $\log \Gamma$, Ψ and Ψ' which will show us, that $g'(x)$ and $g''(x)$ are positive for large x . For smaller x one can see this from the graph of these functions:



The following asymptotic formulas are valid for positive x :

$$\log \Gamma(x) = \left(x - \frac{1}{2} \right) \log x - x + \frac{1}{2} \log(2\pi) + \frac{1}{12x} - \frac{1}{360x^3} + \dots \quad (3)$$

$$\Psi(x) = \log x - \frac{1}{2x} - \frac{1}{12x^2} + \frac{1}{120x^4} - \frac{1}{252x^6} + \dots \quad (4)$$

$$\Psi'(x) = \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3} - \frac{1}{30x^5} + \frac{1}{42x^7} - \dots \quad (5)$$

If we only take a partial sum of one of these series then the error will be less than the first term neglected and has the same sign ([1], p. 257f.).

1. Note that

$$g'(x) = \frac{\Psi(x)}{\log x} - \frac{\log \Gamma(x)}{x(\log x)^2}, \quad x > 0,$$

with a removable singularity at $x = 1$. By (3) and (4) we have

$$\log \Gamma(x) < (x - 1/2x) \log x \quad \text{and} \quad \Psi(x) > \log x - \frac{1}{2x} - \frac{1}{12x^2}.$$

Hence

$$\begin{aligned} g'(x)x(\log x)^2 &= x(\log x)\Psi(x) - \log \Gamma(x) \\ &> x(\log x) \left(\log x - \frac{1}{2x} - \frac{1}{12x^2} \right) - \left(x - \frac{1}{2} \right) \log x > 0 \end{aligned}$$

if $x > 2.7484\dots$.

2. $g''(x) = \frac{\Psi'(x)}{\log x} - \frac{2\Psi(x)}{x(\log x)^2} + \left(\frac{1}{x^2(\log x)^2} + \frac{2}{x^2(\log x)^3} \right) \log \Gamma(x)$ for $x > 0$, again with a removable singularity at $x = 1$. We get

$$\begin{aligned} x^2(\log x)^3 g''(x) &= x^2(\log x)^2 \Psi'(x) + (\log x + 2) \log \Gamma(x) \\ &\quad - 2x \log x \Psi(x). \end{aligned} \quad (6)$$

Here we use the inequalities following from (5), (3) (where $\frac{1}{2} \log(2\pi) = 0.9189\dots$) and (4):

$$\Psi'(x) > \frac{1}{x} + \frac{1}{2x^2}, \quad \log \Gamma(x) > \left(x - \frac{1}{2} \right) \log x - x + 0.9 \quad \text{and} \quad \Psi(x) < \log x.$$

Herewith we get a lower bound for (6) by

$$\left(x - \frac{1}{10} \right) \log x - \frac{10x - 9}{5},$$

which is positive for say $x \geq 6$ (more exactly $x > 5.491776524\dots$). Thus also $g''(x) > 0$ for at least $x > 5.491776524\dots$. $\square$

Remark 1. In the previous proof we found it more appropriate to use computer aided calculations to show the convexity of $g(x)$ for small x than to tackle complicate inequalities. For those who are not convinced by these methods we have a second version of our main result (see Theorem 2 below).

Remark 2. The function $G : \mathbb{R} \rightarrow \mathbb{R}$ defined by $G = g \circ \exp$, i.e.

$$G(x) = \frac{\log \Gamma(\exp x)}{x},$$

as a composition of two strictly increasing and strictly convex functions is again *strictly increasing and strictly convex on $\mathbb{R}$* .

2. The functional equation

The function g satisfies the functional equation

$$f(x+1) = \frac{\log x}{\log(x+1)}(f(x) + 1), \quad x \in (0, \infty). \quad (7)$$

If $f : (0, \infty) \rightarrow \mathbb{R}$ is an arbitrary solution of (7), then (7) with $x = 1$ yields

$$f(2) = 0. \quad (8)$$

Thus, the initial condition (8) is forced by the functional equation (7) itself. Furthermore we have

$$f(n) = \frac{\log \Gamma(n)}{\log n}, \quad n \in \mathbb{N}, \quad n \geq 2.$$

and also

$$f(x+n) = \frac{\log x}{\log(x+n)}f(x) + \frac{\log [(x+n-1) \dots x]}{\log(x+n)}, \quad x \in (0, \infty), \quad n \in \mathbb{N}.$$

Hence also for $x \in (0, \infty)$, $n \in \mathbb{N}$,

$$f(x+n+1) = \frac{\log x}{\log(x+n+1)}f(x) + \frac{\log [(x+n) \dots (x)]}{\log(x+n+1)}, \quad (9)$$

Theorem 1. *The only solution of (7), convex on $(0, \infty)$ is*

$$g(x) = \frac{\log \Gamma(x)}{\log x}.$$

PROOF. Let $f : (0, \infty) \rightarrow \mathbb{R}$ a solution of (7), convex on $(0, \infty)$. Then we have automatically (8) and

$$f(n) = g(n) = \frac{\log(n-1)!}{\log n}, \quad n \in \mathbb{N}, \quad n \geq 2. \quad (10)$$

For $x \in (0, 1]$, $n \in \mathbb{N}$, $n \geq 2$ the convexity condition yields:

$$f(n+1) - f(n) \leq \frac{f(x+n+1) - f(n+1)}{x} \leq f(n+2) - f(n+1), \quad (11)$$

whence

$$\begin{aligned} 0 &\leq \frac{1}{x} [f(x+n+1) - f(n+1) - x(f(n+1) - f(n))] \\ &\leq f(n+2) + f(n) - 2f(n+1). \end{aligned}$$

Hence, applying in turn: relation (9) and (10), multiplication by $\log(x+n+1) > 0$, the monotonicity of $\log$ and $\frac{\log \Gamma}{\log}$, and, finally the Stirling formula which implies that $\log n! < (n + \frac{1}{2}) \log n - n + 1$ (see [1], p. 257), we obtain

$$\begin{aligned} 0 &\leq \frac{1}{x} \left[\log x \cdot f(x) + \log[(x+n) \dots x] \right. \\ &\quad \left. - x \log(x+n+1) \left(\frac{\log n!}{\log(n+1)} - \frac{\log(n-1)!}{\log n} \right) \right] \\ &\leq \log(x+n+1) \left[\frac{\log(n+1)!}{\log(n+2)} + \frac{\log(n-1)!}{\log n} - 2 \frac{\log n!}{\log(n+1)} \right] \\ &\leq \log(n+2) \left[\frac{\log(n+1)!}{\log(n+2)} + \frac{\log(n-1)!}{\log n} - 2 \frac{\log n!}{\log(n+1)} \right] \\ &= \log(n+2) \left[\frac{\log(n+1)!}{\log(n+2)} - \frac{\log n!}{\log(n+1)} \right] \\ &\quad + \log(n+2) \left[\frac{\log(n-1)!}{\log n} - \frac{\log n!}{\log(n+1)} \right] \end{aligned}$$

$$\begin{aligned}
&< \log(n+2) \left[\frac{\log(n+1)!}{\log(n+2)} - \frac{\log n!}{\log(n+1)} \right] \\
&\quad + \log(n+1) \left[\frac{\log(n-1)!}{\log n} - \frac{\log n!}{\log(n+1)} \right] \\
&= \log(n+1)! - \log n! - \frac{\log(n+2)}{\log(n+1)} \log n! + \frac{\log(n+1)}{\log n} \log(n-1)! \\
&= \log(n+1) - \frac{\log(n+1)}{\log n} (\log n! - \log(n-1)!) \\
&\quad + \left(\frac{\log(n+1)}{\log n} - \frac{\log(n+2)}{\log(n+1)} \right) \log n! \\
&= \left[\frac{\log(n+1)}{\log n} - \frac{\log(n+2)}{\log(n+1)} \right] \log n! \\
&< \left[\frac{\log(n+1)}{\log n} - \frac{\log(n+2)}{\log(n+1)} \right] \left[\left(n + \frac{1}{2} \right) \log n - n + 1 \right] =: \theta(n).
\end{aligned}$$

With the aid of the expansions

$$\begin{aligned}
\log(n+1) &= \log n - \log \left(1 - \frac{1}{n+1} \right) = \log n + \sum_{i=1}^{\infty} \frac{1}{i \cdot (n+1)^i}, \\
\log(n+2) &= \log(n+1) + \log \left(1 + \frac{1}{n+1} \right) \\
&= \log(n+1) + \sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i \cdot (n+1)^i}
\end{aligned}$$

we get

$$\begin{aligned}
&\frac{\log(n+1)}{\log n} - \frac{\log(n+2)}{\log(n+1)} \\
&= \frac{1}{\log n} \cdot \sum_{i=1}^{\infty} \frac{1}{i \cdot (n+1)^i} - \frac{1}{\log(n+1)} \cdot \sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i \cdot (n+1)^i} \\
&= \left(\frac{1}{\log n} - \frac{1}{\log(n+1)} \right) \frac{1}{n+1} \\
&\quad + \left(\frac{1}{\log n} + \frac{1}{\log(n+1)} \right) \frac{1}{2(n+1)^2} + \cdots = o \left(\frac{1}{n \log n} \right).
\end{aligned}$$

From this it is easy to see that $\theta(n)$ tends to 0 for $n \rightarrow \infty$. Therefore for $x \in (0, 1]$ we have

$$f(x) = \lim_{n \rightarrow \infty} \left[x \log(x + n + 1) \left(\frac{\log n!}{\log(n+1)} - \frac{\log(n-1)!}{\log n} \right) - \log[(x+n) \dots x] \right] \frac{1}{\log x}$$

that is, f is uniquely determined on $(0, 1]$. According to well known uniqueness theorems on difference equations, we have that f , as a convex solution of (7), is uniquely determined on all of $(0, \infty)$. Since g is also a convex solution of (7), we have $f = g$. $\square$

Theorem 2. *The only solution of (7), convex on a neighborhood of infinity is $g(x) = \frac{\log \Gamma(x)}{\log x}$.*

PROOF. Let $f : (0, \infty) \rightarrow \mathbb{R}$ a solution of (7), convex say for $x > a$ for some positive real a . We can proceed as in the proof of Theorem 1, except that we have suppose in (11) that $n > a$ holds. $\square$

3. Final remarks

In a recent paper [4] P. GRABNER *et al.* showed that the function $x \mapsto \frac{\log \Gamma(x+1)}{x}$ is concave on $(-1, \infty)$ and it is characterized as the only concave solution of its functional equation. In contrary to this the function $\frac{\log \Gamma(x+1)}{\log x} = \frac{\log \Gamma(x)}{\log x} + 1$ is convex on $(0, \infty)$.

In this connection the following question arises. What is the behavior of the function $x \mapsto \frac{\log \Gamma(x+1)}{x^\alpha}$ or $x \mapsto \frac{\log \Gamma(x)}{x^\alpha}$ for a fixed real α ? Here the case $0 < \alpha < 1$ is of interest.

For $\alpha \in (0, 1)$ the plotted graph of these functions looks like convex, at least for small x . Nevertheless it is easy to show that there is a constant c , depending on α , such that these functions are concave for $x > c$.

The function $x \mapsto \frac{\log \Gamma(x)}{x^\alpha}$ e.g. fulfills the functional equation

$$f(x+1) = f(x) \cdot \frac{x^\alpha}{(x+1)^\alpha} + \frac{\log(x)}{(x+1)^\alpha}, \quad x \in (0, \infty). \quad (12)$$

It is routine to show that $x \mapsto \frac{\log \Gamma(x)}{x^\alpha}$ is the only solution of (12), concave in a neighborhood of infinity, together with the side condition $f(1) = 0$. A similar statement can be given for the function $x \mapsto \frac{\log \Gamma(x+1)}{x^\alpha}$.

By this we receive different forms of characterizations of the gamma function.

References

- [1] M. ABRAMOWITZ and I. STEGUN (eds.), Handbook of Mathematical functions, *Dover Publ., New York*, 1970.
- [2] E. ARTIN, Einführung in die Theorie der Gammafunktion, Hamburger Math. Einzelschr., Heft 1, *B.G. Teubner, Leipzig/Berlin*, 1931.
- [3] C. CARATHEODORY, Theory of functions of a complex variable, Vol. I, *Chelsea Publ., New York*, 1964.
- [4] P. J. GRABNER, R. F. TICHY and U. T. ZIMMERMANN, Inequalities for the gamma function with application to permanents, *Discrete Math.* **154** (1996), 53–62.
- [5] D. GRONAU and J. MATKOWSKI, Geometrical convexity and generalizations of the Bohr–Mollerup Theorem on the Gamma function, *Mathematica Pannonica* **4/2** (1993), 1–8.

DETLEF GRONAU
 INSTITUT FÜR MATHEMATIK
 KARL-FRANZENS-UNIVERSITÄT GRAZ
 HEINRICHSTR. 36
 A-8010 GRAZ
 AUSTRIA

E-mail: gronau@uni-graz.at

JANUSZ MATKOWSKI
 INSTITUTE OF MATHEMATICS
 UNIVERSITY OF ZIELONA GÓRA
 PODGÓRNA 50
 PL-65-246 ZIELONA GÓRA
 POLAND

E-mail: jmatk@wsp.zgora.pls

(Received October 24, 2001; revised May 27, 2002)